

Technical Reference

Black Dog Watershed Management Organization Habitat Monitoring Background Summary

In 2002, the Black Dog Watershed Management Organization (BDWMO) created a program for monitoring the habitat quality of strategic water resources in the watershed. The BDWMO lies south of the Minnesota River in the northwest portion of Dakota County. **Figure 1** shows the subwatersheds to the BDWMO's strategic water bodies. The BDWMO began implementing the habitat monitoring program in 2003 and continued the program through 2009. In 2004, based on feedback from the participating cities and to better define the vegetative quality, several improvements were made to the rating system. The BDWMO used this system for the annual habitat monitoring of each strategic water body through 2009. From 2003-2009 Barr staff annually evaluated the habitat quality of each of the following strategic water bodies:

- Crystal Lake (Burnsville)
- Keller Lake (Burnsville)
- Kingsley Lake (Lakeville)
- Lac Lavon (Apple Valley and Burnsville)
- Orchard Lake (Lakeville)
- Sunset Pond (Burnsville)

In 2010, the BDWMO suspended the habitat monitoring program and re-evaluated the program for its effectiveness. Based on feedback obtained from city staff, the BDWMO revised the habitat monitoring program to provide more effective monitoring, more useful and holistic results, and to reduce the monitoring costs. The BDWMO began implementing the revised habitat monitoring program in 2011. Also in 2011, the BDWMO removed Sunset Pond from its list of strategic water bodies.


The revised program includes monitoring habitat quality at one strategic water body per year, such that the BDWMO monitors all five strategic water bodies over a five-year cycle. The 2011 through 2015 reports provided a new baseline for the strategic water bodies—Kingsley Lake (2011), Orchard Lake (2012), Crystal Lake (2013), Lac Lavon (2014), and Keller Lake (2015). This report provides the results of the Lac Lavon 2019 habitat monitoring.

The 2019 Lac Lavon monitoring includes transect, plot, and meandering surveys. Supplemental photographs were taken to document conditions. Private versus public ownership was identified along the entire shoreline. The survey results, along with parcel data, were used to identify possible locations for restoration and preservation. Table 1 of the Technical Memo summarizes the 2019 Lac Lavon monitoring results.

Habitat Quality

The BDWMO's assessment of the BDWMO strategic water bodies provides baseline and ongoing information regarding the habitat quality of the water bodies and a method for detecting change. Habitat quality was evaluated within the following four general zones:

1. **Submergent vegetation zone**—The submergent zone refers to the areas of the water body where water depths are typically 2 to 20 feet (normal maximum rooting depth) and the vegetation is typically submerged or has floating leaves. The vegetation quality within the submergent zone is normally rated as "excellent" when there are: (a) a diverse assemblage of native plant species (more than 14), (b) a moderate plant density or plant occurrence rating, and (c) no exotic species present.
2. **Emergent vegetation zone**—The emergent zone typically refers to the areas of the water body where water depths are less than 2 feet and vegetation grows out of the water. The vegetation quality within the emergent zone is typically rated as "excellent" when there are more than 15 species of native and non-invasive plants present, with few exotic plants present.
3. **Condition of the upland buffer area**—The upland buffer is characterized as the upland area immediately surrounding the water body. An excellent quality buffer should extend upslope at least 25 feet from the wetland edge, consist of native vegetation that is not routinely mowed, and be present continuously around the perimeter of the water body.
4. **Sedimentation and shoreline erosion problems**—The presence of sedimentation may come from erosion on slopes, from storm sewer outfalls, or from other sources. The presence of a regular sediment load to the water body can cause a significant reduction in water quality. Shoreline erosion can be caused by natural forces such as ice and wave action, but can also be human induced (e.g., vegetation removal, grading, runoff, structures, etc.). Identifying and correcting these problems early can prevent habitat degradation.



Vegetation Zones

Appendix C summarizes the overall ratings from 2003 through 2018. **Appendix D** includes the previous management recommendations for water bodies assessed from 2009 through 2018. **Table 2 of the Technical Memo** provides the 2019 management recommendations for Lac Lavoie.

Wildlife Habitat Characteristics

The strategic water bodies within the BDWMO range from shallow wetland systems to deeper lake systems. Some of them support sustainable fisheries, while others may only periodically support fish. All of the water bodies appear to have some potential for supporting waterfowl and shorebirds. To evaluate the wildlife value of these water bodies, it is important to understand the characteristics that will benefit wildlife.

In general, a more diverse assemblage of native plant species will provide a source of food and protective cover for a wider range of wildlife species. Typically, although not always, native plant species do not become established as monocultures to the detriment of other species, as is often the case with many exotic species. As vegetation diversity increases, so does the likelihood that the water body will support a more diverse assemblage of wildlife.

A diverse interspersed of various plant communities also leads to the potential for attracting a wider range of wildlife. For instance, some waterfowl prefer deeper, open water areas while others tend to inhabit the shallow emergent zones. Some furbearers rely heavily on the shallow, emergent zone and upland areas around the water body while others spend most of their time in the deep marsh areas. Amphibians will typically need a permanently inundated water body, but rely on diverse vegetative structure in the upland areas surrounding the water body for critical components of their life cycle. Fish also require permanent inundation to a depth that will not result in freeze-out and where oxygen will not become depleted. A diverse habitat structure is also important for fish.

The upland buffer surrounding these water bodies is important for a number of reasons. A high quality upland buffer will have a diverse vegetative structure dominated by self-sustaining native vegetation. A high quality upland buffer is used by wildlife for shelter, feeding, resting, nesting, and reproduction. In contrast, adjacent upland areas that are maintained in turf grass or paved trails provide little value to wildlife or water quality improvement. Turf grass and trails typically provide feeding and resting grounds only for geese and some species of ducks. Wide and contiguous natural buffers are important as they provide feeding, nesting and safe travel corridors. Upland buffers also help protect the water quality of the water body. Diverse native vegetation helps maintain an open soil structure that promotes infiltration, reduces surface runoff, and increases nutrient uptake.

Wetland Functions and Values Assessment—MNRAM


In addition to the specific habitat parameters described above, the Minnesota Routine Assessment Method for Evaluating Wetland Functions (MNRAM) Version 3.0 was used to evaluate the hydrologic system and ecosystem making up each water resource, first in 2003 and then again in 2006. The results of the 2003 and 2006 MNRAM 3.0 assessments were provided in previous year's reports. Orchard Lake was re-assessed in 2012, Crystal Lake was re-assessed in 2013, Lac Lavon was re-assessed in 2014, Keller Lake was re-assessed in 2015, and Kinsley was re-assessed in 2016 with the more updated MNRAM version 3.4. The results of the 2014 Lac Lavon MNRAM are provided in **Appendix E**. Evaluating each ecosystem with MNRAM is a way to get a detailed picture of the overall health of the watershed and the water resource itself. Instead of just looking at specific parameters that are direct indicators of habitat quality, the MNRAM evaluates many different parameters of the water body and its watershed that contribute to sustaining the wetland functions, which are described in **Appendix F**. In general, the MNRAM assessments compare favorably with the BDWMO habitat vegetation assessment results. This method identifies land use or ecological changes, which might affect the water body in the long term. In addition, the MNRAM assessment provides an independent evaluation of the overall wildlife habitat of the water body.

Appendices


Appendix A

Lac Lavon Aquatic Plant Survey Results


June - August Differences for All Species Lac Lavon - Dakota County, MN




June - August CLP Differences Lac Lavon - Dakota County, MN



June - August EWM Differences Lac Lavon - Dakota County, MN



Maximum Depth of Plant Colonization




Appendix B

Lac Lavon Floristic Quality Assessment Data

2014 Lac Lavon Submergent Vegetation Floristic Quality Index

| Species | Common Name | Coefficient of Conservatism Value (C-value) |
|--|-----------------------|---|
| <i>Ceratophyllum demersum</i> | coontail | 2 |
| <i>Chara sp.</i> | muskgrass | 7 |
| <i>Elodea canadensis</i> | elodea | 4 |
| <i>Myriophyllum spicatum</i> | Eurasian watermilfoil | 0 |
| <i>Najas flexilis</i> | flexuous naiad | 5 |
| <i>Potamogeton crispus</i> | curlyleaf pondweed | 0 |
| <i>Potamogeton foliosus</i> | narrowleaf pondweed | 6 |
| <i>Potamogeton illinoensis</i> | Illinois pondweed | 6 |
| <i>Potamogeton nodosus</i> | longleaf pondweed | 6 |
| <i>Potamogeton pusillus</i> | leafy pondweed | 7 |
| <i>Potamogeton zosteriformis</i> | flatstem pondweed | 6 |
| <i>Ranunculus longirostris</i> | white water crowfoot | 7 |
| <i>Stuckenia pectinatus</i> | sago pondweed | 3 |
| <i>Zosterella dubia</i> | water stargrass | 6 |
| Mean C-value | | 4.6 |
| S (Number of Species of Submergent/Floating-leaf Plants in the Lake) | | 14 |
| Floristic Quality Index (FQI) = (Mean C-value)* (Square Root of S) | | 17.37 |

Lac Lavon 2014 Submergent Vegetation Survey
C-value for each Species




| Species Number | Scientific Name | Common Name | C-value |
|----------------|----------------------------------|-----------------------|---------|
| 1 | <i>Ceratophyllum demersum</i> | coontail | 2 |
| 2 | <i>Chara sp.</i> | muskgrass | 7 |
| 3 | <i>Elodea canadensis</i> | elodea | 4 |
| 4 | <i>Myriophyllum spicatum</i> | Eurasian watermilfoil | 0 |
| 5 | <i>Najas flexilis</i> | flexuous naiad | 5 |
| 6 | <i>Potamogeton crispus</i> | curlyleaf pondweed | 0 |
| 7 | <i>Potamogeton foliosus</i> | narrowleaf pondweed | 6 |
| 8 | <i>Potamogeton illinoensis</i> | Illinois pondweed | 6 |
| 9 | <i>Potamogeton nodosus</i> | longleaf pondweed | 6 |
| 10 | <i>Potamogeton pusillus</i> | leafy pondweed | 7 |
| 11 | <i>Potamogeton zosteriformis</i> | flatstem pondweed | 6 |
| 12 | <i>Ranunculus longirostris</i> | white water crowfoot | 7 |
| 13 | <i>Stuckenia pectinatus</i> | sago pondweed | 3 |
| 14 | <i>Zosterella dubia</i> | water stargrass | 6 |

2019 Lac Lavon Submergent Vegetation Floristic Quality Index

| Species | Common Name | Coefficient of Conservatism Value (C-value) |
|--|-----------------------|---|
| <i>Ceratophyllum demersum</i> | coontail | 2 |
| <i>Chara sp.</i> | muskgrass | 7 |
| <i>Elodea canadensis</i> | elodea | 4 |
| <i>Lemna minor</i> | small duckweed | 5 |
| <i>Spirodela polyrhiza</i> | large duckweed | 5 |
| <i>Myriophyllum spicatum</i> | Eurasian watermilfoil | 0 |
| <i>Potamogeton crispus</i> | curlyleaf pondweed | 0 |
| <i>Potamogeton illinoensis</i> | Illinois pondweed | 6 |
| <i>Potamogeton nodosus</i> | longleaf pondweed | 6 |
| <i>Potamogeton pusillus</i> | leafy pondweed | 7 |
| <i>Ranunculus longirostris</i> | white water crowfoot | 7 |
| <i>Stuckenia pectinatus</i> | sago pondweed | 3 |
| <i>Wolffia columbiana</i> | common watermeal | 5 |
| <i>Zosterella dubia</i> | water stargrass | 6 |
| Mean C-value | | 4.5 |
| S (Number of Species of Submergent/Floating-leaf Plants in the Lake) | | 14 |
| Floristic Quality Index (FQI) = (Mean C-value)* (Square Root of S) | | 16.84 |

Lac Lavon 2019 Submergent Vegetation Survey
C-value for each Species




| Species Number | Scientific Name | Common Name | C-value |
|----------------|--------------------------------|-----------------------|---------|
| 1 | <i>Ceratophyllum demersum</i> | coontail | 2 |
| 2 | <i>Chara sp.</i> | muskgrass | 7 |
| 3 | <i>Elodea canadensis</i> | elodea | 4 |
| 4 | <i>Lemna minor</i> | small duckweed | 5 |
| 5 | <i>Spirodela polyrhiza</i> | large duckweed | 5 |
| 6 | <i>Myriophyllum spicatum</i> | Eurasian watermilfoil | 0 |
| 7 | <i>Potamogeton crispus</i> | curlyleaf pondweed | 0 |
| 8 | <i>Potamogeton illinoensis</i> | Illinois pondweed | 6 |
| 9 | <i>Potamogeton nodosus</i> | longleaf pondweed | 6 |
| 10 | <i>Potamogeton pusillus</i> | leafy pondweed | 7 |
| 11 | <i>Ranunculus longirostris</i> | white water crowfoot | 7 |
| 12 | <i>Stuckenia pectinatus</i> | sago pondweed | 3 |
| 13 | <i>Wolffia columbiana</i> | common watermeal | 5 |
| 14 | <i>Zosterella dubia</i> | water stargrass | 6 |

2014 Lac Lavon Emergent Vegetation Floristic Quality Index

| Species | Common Name | Coefficient of Conservatism Value |
|--|-----------------------|-----------------------------------|
| <i>Ambrosia artemisiifolia</i> | common ragweed | 0 |
| <i>Asclepias incarnata</i> | swamp milkweed | 4 |
| <i>Beckmannia syzigachne</i> | slough grass | 4 |
| <i>Carex stricta</i> | Uptight Sedge | 5 |
| <i>Carex vulpinoidea</i> | fox sedge | 3 |
| <i>Centaureum sp.</i> | knapweed | 0 |
| <i>Cirsium arvense</i> | Canada thistle | 0 |
| <i>Equisetum pratense</i> | meadow horsetail | 6 |
| <i>Erigeron philadelphicus</i> | Philadelphia fleabane | 2 |
| <i>Eupatorium perfoliatum</i> | boneset | 4 |
| <i>Fraxinus pennsylvanica</i> | green ash | 2 |
| <i>Glechoma hederacea</i> | ground ivy | 0 |
| <i>Impatiens capensis</i> | jewelweed | 2 |
| <i>Iris versicolor</i> | harlequin blueflag | 4 |
| <i>Juncus effusus</i> | soft rush | 4 |
| <i>Juncus tenuis</i> | path rush | 1 |
| <i>Lactuca serriola</i> | prickly lettuce | 0 |
| <i>Lotus corniculatus</i> | bird's-foot trefoil | 0 |
| <i>Lycopus uniflorus</i> | northern bugleweed | 5 |
| <i>Lythrum salicaria</i> | purple loosestrife | 0 |
| <i>Melilotus officinalis</i> | sweet clover | 0 |
| <i>Mentha arvensis</i> | wild mint | 3 |
| <i>Onoclea sensibilis</i> | sensitive fern | 4 |
| <i>Parthenocissus vitacea</i> | woodbine | 2 |
| <i>Phalaris arundinacea</i> | reed canarygrass | 0 |
| <i>Plantago major</i> | common plantain | 0 |
| <i>Poa palustris</i> | fowl bluegrass | 5 |
| <i>Populus deltoides</i> | eastern cottonwood | 1 |
| <i>Potamogeton nodosus</i> | longleaf pondweed | 6 |
| <i>Potentilla simplex</i> | cinquefoil | 2 |
| <i>Rhus hirta</i> | smooth sumac | 2 |
| <i>Rumex crispus</i> | curly dock | 0 |
| <i>Salix amygdaloides</i> | peach leaf willow | 5 |
| <i>Salix babylonica</i> | weeping willow | 0 |
| <i>Salix interior</i> | sandbar willow | 2 |
| <i>Schoenoplectus tabernaemontani</i> | softstem bulrush | 4 |
| <i>Solidago canadensis</i> | Canada goldenrod | 1 |
| <i>Solidago gigantea</i> | Late Goldenrod | 3 |
| <i>Taraxacum officinale</i> | common dandelion | 0 |
| <i>Tradescantia sp.</i> | spiderwort | 6 |
| <i>Trifolium repens</i> | white clover | 0 |
| <i>Typha angustifolia</i> | narrowleaf cattail | 0 |
| <i>Urtica dioica</i> | Stinging Nettle | 1 |
| <i>Verbena hastata</i> | blue vervain | 6 |
| <i>Zizia aurea</i> | golden alexanders | 6 |
| Mean C-value | | 2.3 |
| S (Number of Species of Emergent Plants in the Lake) | | 45 |
| Floristic Quality Index (FQI) = (Mean C-value)* (Square Root of S) | | 15.65 |

Lac Lavon 2014 Emergent Vegetation Survey
C-value for each Species



Lac Lavon 2014 Emergent Vegetation Survey

| Species Number | Scientific Name | Common Name | C-value |
|----------------|---------------------------------------|-----------------------|---------|
| 1 | <i>Ambrosia artemisiifolia</i> | common ragweed | 0 |
| 2 | <i>Asclepias incarnata</i> | swamp milkweed | 4 |
| 3 | <i>Beckmannia syzigachne</i> | slough grass | 4 |
| 4 | <i>Carex stricta</i> | Uptight Sedge | 5 |
| 5 | <i>Carex vulpinoidea</i> | fox sedge | 3 |
| 6 | <i>Centaurium sp.</i> | knapweed | 0 |
| 7 | <i>Cirsium arvense</i> | Canada thistle | 0 |
| 8 | <i>Equisetum pratense</i> | meadow horsetail | 6 |
| 9 | <i>Erigeron philadelphicus</i> | Philadelphia fleabane | 2 |
| 10 | <i>Eupatorium perfoliatum</i> | boneset | 4 |
| 11 | <i>Fraxinus pennsylvanica</i> | green ash | 2 |
| 12 | <i>Glechoma hederacea</i> | ground ivy | 0 |
| 13 | <i>Impatiens capensis</i> | jewelweed | 2 |
| 14 | <i>Iris versicolor</i> | harlequin blueflag | 4 |
| 15 | <i>Juncus effusus</i> | soft rush | 4 |
| 16 | <i>Juncus tenuis</i> | path rush | 1 |
| 17 | <i>Lactuca serriola</i> | prickly lettuce | 0 |
| 18 | <i>Lotus corniculatus</i> | bird's-foot trefoil | 0 |
| 19 | <i>Lycopus uniflorus</i> | northern bugleweed | 5 |
| 20 | <i>Lythrum salicaria</i> | purple loosestrife | 0 |
| 21 | <i>Melilotus officinalis</i> | sweet clover | 0 |
| 22 | <i>Mentha arvensis</i> | wild mint | 3 |
| 23 | <i>Onoclea sensibilis</i> | sensitive fern | 4 |
| 24 | <i>Parthenocissus vitacea</i> | woodbine | 2 |
| 25 | <i>Phalaris arundinacea</i> | reed canarygrass | 0 |
| 26 | <i>Plantago major</i> | common plantain | 0 |
| 27 | <i>Poa palustris</i> | fowl bluegrass | 5 |
| 28 | <i>Populus deltoides</i> | eastern cottonwood | 1 |
| 29 | <i>Potamogeton nodosus</i> | longleaf pondweed | 6 |
| 30 | <i>Potentilla simplex</i> | cinquefoil | 2 |
| 31 | <i>Rhus hirta</i> | smooth sumac | 2 |
| 32 | <i>Rumex crispus</i> | curly dock | 0 |
| 33 | <i>Salix amygdaloides</i> | peach leaf willow | 5 |
| 34 | <i>Salix babylonica</i> | weeping willow | 0 |
| 35 | <i>Salix interior</i> | sandbar willow | 2 |
| 36 | <i>Schoenoplectus tabernaemontani</i> | softstem bulrush | 4 |
| 37 | <i>Solidago canadensis</i> | Canada goldenrod | 1 |
| 38 | <i>Solidago gigantea</i> | Late Goldenrod | 3 |
| 39 | <i>Taraxacum officinale</i> | common dandelion | 0 |
| 40 | <i>Tradescantia sp.</i> | spiderwort | 6 |
| 41 | <i>Trifolium repens</i> | white clover | 0 |
| 42 | <i>Typha angustifolia</i> | narrowleaf cattail | 0 |
| 43 | <i>Urtica dioica</i> | Stinging Nettle | 1 |
| 44 | <i>Verbena hastata</i> | blue vervain | 6 |
| 45 | <i>Zizia aurea</i> | golden alexanders | 6 |


2019 Lac Lavon Emergent Vegetation Floristic Quality Index

| Species | Common Name | Coefficient of Conservatism Value |
|---------------------------------------|-----------------------|-----------------------------------|
| <i>Acer ginnala</i> | amur maple | 0 |
| <i>Acer negundo</i> | boxelder | 1 |
| <i>Acer saccharinum</i> | silver maple | 3 |
| <i>Ambrosia artemisiifolia</i> | common ragweed | 0 |
| <i>Asclepias incarnata</i> | swamp milkweed | 4 |
| <i>Asclepias syriaca</i> | common milkweed | 0 |
| <i>Carex comosa</i> | bristly sedge | 4 |
| <i>Carex stricta</i> | Uptight Sedge | 5 |
| <i>Carex vulpinoidea</i> | fox sedge | 3 |
| <i>Centaurium sp.</i> | knapweed | 0 |
| <i>Cirsium arvense</i> | Canada thistle | 0 |
| <i>Cyperus sp.</i> | flat sedge | 3 |
| <i>Echinochloa crus-galli</i> | barnyard grass | 0 |
| <i>Eleocharis erythropoda</i> | red rooted spikerush | 3 |
| <i>Equisetum pratense</i> | meadow horsetail | 6 |
| <i>Erigeron philadelphicus</i> | Philadelphia fleabane | 2 |
| <i>Eupatorium perfoliatum</i> | boneset | 4 |
| <i>Fraxinus pennsylvanica</i> | green ash | 2 |
| <i>Glechoma hederacea</i> | ground ivy | 0 |
| <i>Impatiens capensis</i> | jewelweed | 2 |
| <i>Iris versicolor</i> | harlequin blueflag | 4 |
| <i>Juncus effusus</i> | soft rush | 4 |
| <i>Juncus tenuis</i> | path rush | 1 |
| <i>Juncus torreyi</i> | Torrey's rush | 4 |
| <i>Laportea canadensis</i> | wood nettle | 3 |
| <i>Leersia oryzoides</i> | rice cut grass | 3 |
| <i>Lemna minor</i> | small duckweed | 5 |
| <i>Lycopus uniflorus</i> | northern bugleweed | 5 |
| <i>Lythrum salicaria</i> | purple loosestrife | 0 |
| <i>Onoclea sensibilis</i> | sensitive fern | 4 |
| <i>Panicum virgatum</i> | switchgrass | 2 |
| <i>Parthenocissus vitacea</i> | woodbine | 2 |
| <i>Persicaria amphibia</i> | water smartweed | 4 |
| <i>Phalaris arundinacea</i> | reed canarygrass | 0 |
| <i>Plantago major</i> | common plantain | 0 |
| <i>Poa palustris</i> | fowl bluegrass | 5 |
| <i>Populus deltoides</i> | eastern cottonwood | 1 |
| <i>Potentilla simplex</i> | cinquefoil | 2 |
| <i>Rhamnus cathartica</i> | common buckthorn | 0 |
| <i>Rumex crispus</i> | curly dock | 0 |
| <i>Salix amygdaloides</i> | peach leaf willow | 5 |
| <i>Salix babylonica</i> | weeping willow | 0 |
| <i>Salix interior</i> | sandbar willow | 2 |
| <i>Salix petiolaris</i> | slender willow | 5 |
| <i>Schoenoplectus tabernaemontani</i> | softstem bulrush | 4 |
| <i>Scirpus cyperinus</i> | woolgrass | 3 |
| <i>Solidago gigantea</i> | Late Goldenrod | 3 |
| <i>Taraxacum officinale</i> | common dandelion | 0 |
| <i>Tradescantia sp.</i> | spiderwort | 6 |

2019 Lac Lavon Emergent Vegetation Floristic Quality Index

| Species | Common Name | Coefficient of Conservatism Value |
|--|--------------------|-----------------------------------|
| <i>Trifolium repens</i> | white clover | 0 |
| <i>Typha angustifolia</i> | narrowleaf cattail | 0 |
| <i>Ulmus pumila</i> | Siberian elm | 0 |
| <i>Urtica dioica</i> | Stinging Nettle | 1 |
| <i>Verbena hastata</i> | blue vervain | 6 |
| <i>Zizia aurea</i> | golden alexanders | 6 |
| Mean C-value | | 2.4 |
| S (Number of Species of Emergent Plants in the Lake) | | 55 |
| Floristic Quality Index (FQI) = (Mean C-value)* (Square Root of S) | | 17.80 |

Lac Lavon 2019 Emergent Vegetation Survey
C-value for each Species



Lac Lavon 2019 Emergent Vegetation Survey

| Species Number | Scientific Name | Common Name | C-value |
|----------------|---------------------------------------|-----------------------|---------|
| 1 | <i>Acer ginnala</i> | amur maple | 0 |
| 2 | <i>Acer negundo</i> | boxelder | 1 |
| 3 | <i>Acer saccharinum</i> | silver maple | 3 |
| 4 | <i>Ambrosia artemisiifolia</i> | common ragweed | 0 |
| 5 | <i>Asclepias incarnata</i> | swamp milkweed | 4 |
| 6 | <i>Asclepias syriaca</i> | common milkweed | 0 |
| 7 | <i>Carex comosa</i> | bristly sedge | 4 |
| 8 | <i>Carex stricta</i> | Uptight Sedge | 5 |
| 9 | <i>Carex vulpinoidea</i> | fox sedge | 3 |
| 10 | <i>Centaureum sp.</i> | knapweed | 0 |
| 11 | <i>Cirsium arvense</i> | Canada thistle | 0 |
| 12 | <i>Cyperus sp.</i> | flat sedge | 3 |
| 13 | <i>Echinochloa crus-galli</i> | barnyard grass | 0 |
| 14 | <i>Eleocharis erythropoda</i> | red rooted spikerush | 3 |
| 15 | <i>Equisetum pratense</i> | meadow horsetail | 6 |
| 16 | <i>Erigeron philadelphicus</i> | Philadelphia fleabane | 2 |
| 17 | <i>Eupatorium perfoliatum</i> | boneset | 4 |
| 18 | <i>Fraxinus pennsylvanica</i> | green ash | 2 |
| 19 | <i>Glechoma hederacea</i> | ground ivy | 0 |
| 20 | <i>Impatiens capensis</i> | jewelweed | 2 |
| 21 | <i>Iris versicolor</i> | harlequin blueflag | 4 |
| 22 | <i>Juncus effusus</i> | soft rush | 4 |
| 23 | <i>Juncus tenuis</i> | path rush | 1 |
| 24 | <i>Juncus torreyi</i> | Torrey's rush | 4 |
| 25 | <i>Laportea canadensis</i> | wood nettle | 3 |
| 26 | <i>Leersia oryzoides</i> | rice cut grass | 3 |
| 27 | <i>Lemna minor</i> | small duckweed | 5 |
| 28 | <i>Lycopus uniflorus</i> | northern bugleweed | 5 |
| 29 | <i>Lythrum salicaria</i> | purple loosestrife | 0 |
| 30 | <i>Onoclea sensibilis</i> | sensitive fern | 4 |
| 31 | <i>Panicum virgatum</i> | switchgrass | 2 |
| 32 | <i>Parthenocissus vitacea</i> | woodbine | 2 |
| 33 | <i>Persicaria amphibia</i> | water smartweed | 4 |
| 34 | <i>Phalaris arundinacea</i> | reed canarygrass | 0 |
| 35 | <i>Plantago major</i> | common plantain | 0 |
| 36 | <i>Poa palustris</i> | fowl bluegrass | 5 |
| 37 | <i>Populus deltoides</i> | eastern cottonwood | 1 |
| 38 | <i>Potentilla simplex</i> | cinquefoil | 2 |
| 39 | <i>Rhamnus cathartica</i> | common buckthorn | 0 |
| 40 | <i>Rumex crispus</i> | curly dock | 0 |
| 41 | <i>Salix amygdaloides</i> | peach leaf willow | 5 |
| 42 | <i>Salix babilonica</i> | weeping willow | 0 |
| 43 | <i>Salix interior</i> | sandbar willow | 2 |
| 44 | <i>Salix petiolaris</i> | slender willow | 5 |
| 45 | <i>Schoenoplectus tabernaemontani</i> | softstem bulrush | 4 |
| 46 | <i>Scirpus cyperinus</i> | woolgrass | 3 |
| 47 | <i>Solidago gigantea</i> | Late Goldenrod | 3 |
| 48 | <i>Taraxacum officinale</i> | common dandelion | 0 |
| 49 | <i>Tradescantia sp.</i> | spiderwort | 6 |
| 50 | <i>Trifolium repens</i> | white clover | 0 |
| 51 | <i>Typha angustifolia</i> | narrowleaf cattail | 0 |
| 52 | <i>Ulmus pumila</i> | Siberian elm | 0 |
| 53 | <i>Urtica dioica</i> | Stinging Nettle | 1 |
| 54 | <i>Verbena hastata</i> | blue vervain | 6 |
| 55 | <i>Zizia aurea</i> | golden alexanders | 6 |

2014 Lac Lavon Upland Buffer Vegetation Floristic Quality Index


| Species | Common Name | Coefficient of Conservatism Value (C-value) |
|--------------------------------|-----------------------|--|
| <i>Acer negundo</i> | boxelder | 1 |
| <i>Achillea millefolium</i> | yarrow | 1 |
| <i>Arctium minus</i> | burdock | 0 |
| <i>Asclepias syriaca</i> | common milkweed | 0 |
| <i>Asclepias tuberosa</i> | butterfly weed | 6 |
| <i>Berteroa incana</i> | hoary allysum | 0 |
| <i>Bromus inermis</i> | smooth brome | 0 |
| <i>Carex comosa</i> | bristly sedge | 4 |
| <i>Centaurea biebersteinii</i> | spotted knapweed | 0 |
| <i>Cirsium vulgare</i> | bull thistle | 0 |
| <i>Cornus alba</i> | red osier dogwood | 3 |
| <i>Dactylis glomerata</i> | Orchard Grass | 0 |
| <i>Equisetum arvense</i> | field horsetail | 1 |
| <i>Erigeron philadelphicus</i> | Philadelphia fleabane | 2 |
| <i>Erigeron strigosus</i> | daisy fleabane | 2 |
| <i>Euphorbia esula</i> | leafy spurge | 0 |
| <i>Frangula alnus</i> | glossy buckthorn | 0 |
| <i>Fraxinus pennsylvanica</i> | green ash | 2 |
| <i>Galium aparine</i> | cleavers | 1 |
| <i>Gleditsia triacanthos</i> | honey locust | 0 |
| <i>Lactuca serriola</i> | prickly lettuce | 0 |
| <i>Leonorus cardiaca</i> | motherwort | 0 |
| <i>Leucanthemum vulgare</i> | daisy | 0 |
| <i>Lonicera tatarica</i> | Tatarian honeysuckle | 0 |
| <i>Lotus corniculatus</i> | bird's-foot trefoil | 0 |
| <i>Lythrum salicaria</i> | purple loosestrife | 0 |
| <i>Medicago lupulina</i> | black medic | 0 |
| <i>Melilotus officinalis</i> | sweetclover | 0 |
| <i>Mentha arvensis</i> | wild mint | 3 |
| <i>Nepeta cataria</i> | catnip | 0 |
| <i>Panicum virgatum</i> | switchgrass | 2 |
| <i>Parthenocissus vitacea</i> | woodbine | 2 |
| <i>Phalaris arundinacea</i> | reed canarygrass | 0 |
| <i>Phleum pratense</i> | timothy | 0 |
| <i>Pinus resinosa</i> | red pine | 5 |
| <i>Poa pratensis</i> | Kentucky bluegrass | 0 |
| <i>Populus deltoides</i> | cottonwood | 1 |
| <i>Potentilla simplex</i> | cinquefoil | 2 |
| <i>Rhamnus cathartica</i> | common buckthorn | 0 |
| <i>Rhus hirta</i> | smooth sumac | 2 |
| <i>Rudbeckia hirta</i> | black eyed Susan | 3 |
| <i>Rumex crispus</i> | curly dock | 0 |
| <i>Salix amygdaloides</i> | peach leaf willow | 5 |
| <i>Salix interior</i> | sandbar willow | 2 |

2014 Lac Lavon Upland Buffer Vegetation Floristic Quality Index

| Species | Common Name | Coefficient of Conservatism Value (C-value) |
|--|---------------------|---|
| <i>Salix lucida</i> | shining willow | 5 |
| <i>Salix nigra</i> | black willow | 4 |
| <i>Silene latifolia</i> | bladder campion | 0 |
| <i>Silphium perfoliatum</i> | cup plant | 4 |
| <i>Solanum dulcamara</i> | climbing nightshade | 0 |
| <i>Solidago canadensis</i> | Canada goldenrod | 1 |
| <i>Solidago gigantea</i> | Late Goldenrod | 3 |
| <i>Sorghastrum nutans</i> | Indian grass | 5 |
| <i>Taraxacum officinale</i> | common dandelion | 0 |
| <i>Trifolium hybridum</i> | alsike clover | 0 |
| <i>Trifolium pratense</i> | red clover | 0 |
| <i>Trifolium procumbens</i> | hop clover | 0 |
| <i>Trifolium repens</i> | white clover | 0 |
| <i>Ulmus americana</i> | American elm | 3 |
| <i>Ulmus pumila</i> | Siberian elm | 0 |
| <i>Verbascum thapsus</i> | mullein | 0 |
| <i>Vicia sativa</i> | vetch | 0 |
| <i>Viola</i> sp. * | violet | 6 |
| <i>Vitis riparia</i> | wild grape | 2 |
| Mean C-value | | 1.3 |
| S (Number of Species of Upland Buffer Plants) | | 63 |
| Floristic Quality Index (FQI) = (Mean C-value)* (Square Root of S) | | 10.46 |

* An average C-value was used for this genus, since the species was not verified.

Lac Lavon 2014 Upland Buffer Vegetation Survey
C-value for each Species



Lac Lavon 2014 Upland Buffer Vegetation Survey

| Species Number | Scientific Name | Common Name | C-value |
|----------------|--------------------------------|-----------------------|---------|
| 1 | <i>Acer negundo</i> | boxelder | 1 |
| 2 | <i>Achillea millefolium</i> | yarrow | 1 |
| 3 | <i>Arctium minus</i> | burdock | 0 |
| 4 | <i>Asclepias syriaca</i> | common milkweed | 0 |
| 5 | <i>Asclepias tuberosa</i> | butterfly weed | 6 |
| 6 | <i>Berteroia incana</i> | hoary allysum | 0 |
| 7 | <i>Bromus inermis</i> | smooth brome | 0 |
| 8 | <i>Carex comosa</i> | bristly sedge | 4 |
| 9 | <i>Centaurea biebersteinii</i> | spotted knapweed | 0 |
| 10 | <i>Cirsium vulgare</i> | bull thistle | 0 |
| 11 | <i>Cornus alba</i> | red osier dogwood | 3 |
| 12 | <i>Dactylis glomerata</i> | Orchard Grass | 0 |
| 13 | <i>Equisetum arvense</i> | field horsetail | 1 |
| 14 | <i>Erigeron philadelphicus</i> | Philadelphia fleabane | 2 |
| 15 | <i>Erigeron strigosus</i> | daisy fleabane | 2 |
| 16 | <i>Euphorbia esula</i> | leafy spurge | 0 |
| 17 | <i>Frangula alnus</i> | glossy buckthorn | 0 |
| 18 | <i>Fraxinus pennsylvanica</i> | green ash | 2 |
| 19 | <i>Galium aparine</i> | cleavers | 1 |
| 20 | <i>Gleditsia triacanthos</i> | honey locust | 0 |
| 21 | <i>Lactuca serriola</i> | prickly lettuce | 0 |
| 22 | <i>Leonoros cardiaca</i> | motherwort | 0 |
| 23 | <i>Leucanthemum vulgare</i> | daisy | 0 |
| 24 | <i>Lonicera tatarica</i> | Tatarian honeysuckle | 0 |
| 25 | <i>Lotus corniculatus</i> | bird's-foot trefoil | 0 |
| 26 | <i>Lythrum salicaria</i> | purple loosestrife | 0 |
| 27 | <i>Medicago lupulina</i> | black medic | 0 |
| 28 | <i>Melilotus officinalis</i> | sweetclover | 0 |
| 29 | <i>Mentha arvensis</i> | wild mint | 3 |
| 30 | <i>Nepeta cataria</i> | catnip | 0 |
| 31 | <i>Panicum virgatum</i> | switchgrass | 2 |
| 32 | <i>Parthenocissus vitacea</i> | woodbine | 2 |
| 33 | <i>Phalaris arundinacea</i> | reed canarygrass | 0 |
| 34 | <i>Phleum pratense</i> | timothy | 0 |
| 35 | <i>Pinus resinosa</i> | red pine | 5 |
| 36 | <i>Poa pratensis</i> | Kentucky bluegrass | 0 |
| 37 | <i>Populus deltoides</i> | cottonwood | 1 |
| 38 | <i>Potentilla simplex</i> | cinquefoil | 2 |
| 39 | <i>Rhamnus cathartica</i> | common buckthorn | 0 |
| 40 | <i>Rhus hirta</i> | smooth sumac | 2 |
| 41 | <i>Rudbeckia hirta</i> | black eyed Susan | 3 |
| 42 | <i>Rumex crispus</i> | curly dock | 0 |
| 43 | <i>Salix amygdaloides</i> | peach leaf willow | 5 |
| 44 | <i>Salix interior</i> | sandbar willow | 2 |
| 45 | <i>Salix lucida</i> | shinging willow | 5 |
| 46 | <i>Salix nigra</i> | black willow | 4 |
| 47 | <i>Silene latifolia</i> | bladder campion | 0 |
| 48 | <i>Silphium perfoliatum</i> | cup plant | 4 |
| 49 | <i>Solanum dulcamara</i> | climbing nightshade | 0 |
| 50 | <i>Solidago canadensis</i> | Canada goldenrod | 1 |
| 51 | <i>Solidago gigantea</i> | Late Goldenrod | 3 |
| 52 | <i>Sorghastrum nutans</i> | Indian grass | 5 |
| 53 | <i>Taraxacum officinale</i> | common dandelion | 0 |
| 54 | <i>Trifolium hybridum</i> | alsike clover | 0 |
| 55 | <i>Trifolium pratense</i> | red clover | 0 |
| 56 | <i>Trifolium procumbens</i> | hop clover | 0 |
| 57 | <i>Trifolium repens</i> | white clover | 0 |
| 58 | <i>Ulmus americana</i> | American elm | 3 |
| 59 | <i>Ulmus pumila</i> | Siberian elm | 0 |
| 60 | <i>Verbascum thapsus</i> | mullein | 0 |
| 61 | <i>Vicia sativa</i> | vetch | 0 |
| 62 | <i>Viola sp. *</i> | violet | 6 |
| 63 | <i>Vitis riparia</i> | wild grape | 2 |

2019 Lac Lavon Upland Buffer Vegetation Floristic Quality Index

| Species | Common Name | Coefficient of Conservatism Value (C-value) |
|----------------------------------|--------------------|--|
| <i>Acer negundo</i> | boxelder | 1 |
| <i>Acer rubrum</i> | red maple | 3 |
| <i>Achillea millefolium</i> | yarrow | 1 |
| <i>Ageratina altissima</i> | white snakeroot | 2 |
| <i>Ambrosia artemisiifolia</i> | common ragweed | 0 |
| <i>Ambrosia trifida</i> | great ragweed | 0 |
| <i>Andropogon gerardii</i> | big bluestem | 4 |
| <i>Apocynum cannabinum</i> | dogbane | 3 |
| <i>Arctium minus</i> | burdock | 0 |
| <i>Aronia melanocarpa</i> *** | black chokeberry | 7 |
| <i>Asclepias incarnata</i> | swamp milkweed | 4 |
| <i>Asclepias syriaca</i> | common milkweed | 0 |
| <i>Asclepias tuberosa</i> | butterfly weed | 6 |
| <i>Baptisia alba</i> | white wild indigo | 8 |
| <i>Berteroa incana</i> | hoary allysum | 0 |
| <i>Bouteloua curtipendula</i> | side oats grama | 6 |
| <i>Bromus inermis</i> | smooth brome | 0 |
| <i>Carex comosa</i> | bristly sedge | 4 |
| <i>Carex pensylvanica</i> *** | Pennsylvania sedge | 3 |
| <i>Centaurea biebersteinii</i> | spotted knapweed | 0 |
| <i>Chamecrista fasciculata</i> | partridge pea | 2 |
| <i>Cirsium vulgare</i> | bull thistle | 0 |
| <i>Cornus alba</i> | red osier dogwood | 3 |
| <i>Cornus racemosa</i> | gray dogwood | 2 |
| <i>Dactylis glomerata</i> | Orchard Grass | 0 |
| <i>Daucus carota</i> | Queen Anne's lace | 0 |
| <i>Echnacea purpurea</i> ** | purple coneflower | 10 |
| <i>Equisetum arvense</i> | field horsetail | 1 |
| <i>Erigeron strigosus</i> | daisy fleabane | 2 |
| <i>Euphorbia esula</i> | leafy spurge | 0 |
| <i>Eutrochium maculatum</i> | joe pye weed | 4 |
| <i>Frangula alnus</i> | glossy buckthorn | 0 |
| <i>Fraxinus pennsylvanica</i> | green ash | 2 |
| <i>Galium aparine</i> | cleavers | 1 |
| <i>Geum triflorum</i> | prairie smoke | 7 |
| <i>Glechoma hederacea</i> | ground ivy | 0 |
| <i>Gleditsia triacanthos</i> | honey locust | 0 |
| <i>Hackelia virginiana</i> | Virginia stickseed | 1 |
| <i>Helianthus grosseserratus</i> | sawtooth sunflower | 3 |
| <i>Hemerocallis sp.</i> | day lily | 0 |
| <i>Hosta sp.</i> | hosta | 0 |
| <i>Impatiens capensis</i> | jewelweed | 2 |
| <i>Juglans nigra</i> | black walnut | 4 |
| <i>Laportea canadensis</i> | wood nettle | 3 |

2019 Lac Lavon Upland Buffer Vegetation Floristic Quality Index


| Species | Common Name | Coefficient of Conservatism Value (C-value) |
|--------------------------------|----------------------|--|
| <i>Leonorus cardiaca</i> | motherwort | 0 |
| <i>Leucanthemum vulgare</i> | daisy | 0 |
| <i>Lonicera tatarica</i> | Tatarian honeysuckle | 0 |
| <i>Lotus corniculatus</i> | bird's-foot trefoil | 0 |
| <i>Lythrum salicaria</i> | purple loosestrife | 0 |
| <i>Malus sp.</i> | apple | 0 |
| <i>Medicago lupulina</i> | black medic | 0 |
| <i>Melilotus officinalis</i> | sweetclover | 0 |
| <i>Mentha arvensis</i> | wild mint | 3 |
| <i>Monarda fistulosa</i> | wild bergamot | 3 |
| <i>Nepeta cataria</i> | catnip | 0 |
| <i>Oligoneuron rigidum</i> | stiff goldenrod | 4 |
| <i>Panicum virgatum</i> | switchgrass | 2 |
| <i>Parthenocissus vitacea</i> | woodbine | 2 |
| <i>Phalaris arundinacea</i> | reed canarygrass | 0 |
| <i>Phleum pratense</i> | timothy | 0 |
| <i>Pinus resinosa</i> | red pine | 5 |
| <i>Plantago major</i> | common plantain | 0 |
| <i>Poa pratensis</i> | Kentucky bluegrass | 0 |
| <i>Populus deltoides</i> | cottonwood | 1 |
| <i>Potentilla simplex</i> | cinquefoil | 2 |
| <i>Ratbida pinnata</i> *** | globular coneflower | 4 |
| <i>Rhamnus cathartica</i> | common buckthorn | 0 |
| <i>Rhus hirta</i> | smooth sumac | 2 |
| <i>Ribes americanum</i> | wild black currant | 4 |
| <i>Rosa sp.</i> | rose | 0 |
| <i>Rubus idaeus</i> | red raspberry | 3 |
| <i>Rudbeckia hirta</i> | black eyed Susan | 3 |
| <i>Rumex crispus</i> | curly dock | 0 |
| <i>Salix amygdaloides</i> | peach leaf willow | 5 |
| <i>Salix interior</i> | sandbar willow | 2 |
| <i>Schizachyrium scoparium</i> | little bluestem | 5 |
| <i>Silene latifolia</i> | bladder campion | 0 |
| <i>Silphium perfoliatum</i> | cup plant | 4 |
| <i>Solanum dulcamara</i> | climbing nightshade | 0 |
| <i>Solidago canadensis</i> | Canada goldenrod | 1 |
| <i>Solidago flexicaulis</i> | zigzag goldenrod | 5 |
| <i>Solidago gigantea</i> | Late Goldenrod | 3 |
| <i>Solidago speciosa</i> | showy goldenrod | 5 |
| <i>Sonchus arvensis</i> | sow thistle | 0 |
| <i>Sorghastrum nutans</i> | Indian grass | 5 |
| <i>Taraxacum officinale</i> | common dandelion | 0 |
| <i>Trifolium hybridum</i> | alsike clover | 0 |
| <i>Trifolium pratense</i> | red clover | 0 |

2019 Lac Lavon Upland Buffer Vegetation Floristic Quality Index

| Species | Common Name | Coefficient of Conservatism Value (C-value) |
|--|--------------------|---|
| <i>Ulmus americana</i> | American elm | 3 |
| <i>Ulmus pumila</i> | Siberian elm | 0 |
| <i>Verbascum thapsus</i> | mullein | 0 |
| <i>Verbena hastata</i> | blue vervain | 6 |
| <i>Verbena stricta</i> *** | hoary vervain | 3 |
| <i>Vicia sativa</i> | vetch | 0 |
| <i>Viola</i> sp. * | violet | 6 |
| <i>Vitis riparia</i> | wild grape | 2 |
| <i>Zanthoxylum americanum</i> | common prickly ash | 3 |
| Mean C-value | | 2.0 |
| S (Number of Species of Upland Buffer Plants) | | 97 |
| Floristic Quality Index (FQI) = (Mean C-value)* (Square Root of S) | | 19.80 |

* An average C-value was used for this genus, since the species was not verified.

Lac Lavon 2019 Upland Buffer Vegetation Survey
C-value for each Species



Lac Lavon 2019 Upland Buffer Vegetation Survey

| Species Number | Scientific Name | Common Name | C-value |
|----------------|----------------------------------|----------------------|---------|
| 1 | <i>Acer negundo</i> | boxelder | 1 |
| 2 | <i>Acer rubrum</i> | red maple | 3 |
| 3 | <i>Achillea millefolium</i> | yarrow | 1 |
| 4 | <i>Ageratina altissima</i> | white snakeroot | 2 |
| 5 | <i>Ambrosia artemisiifolia</i> | common ragweed | 0 |
| 6 | <i>Ambrosia trifida</i> | great ragweed | 0 |
| 7 | <i>Andropogon gerardii</i> | big bluestem | 4 |
| 8 | <i>Apocynum cannabinum</i> | dogbane | 3 |
| 9 | <i>Arctium minus</i> | burdock | 0 |
| 10 | <i>Aronia melanocarpa</i> *** | black chokeberry | 7 |
| 11 | <i>Asclepias incarnata</i> | swamp milkweed | 4 |
| 12 | <i>Asclepias syriaca</i> | common milkweed | 0 |
| 13 | <i>Asclepias tuberosa</i> | butterfly weed | 6 |
| 14 | <i>Baptisia alba</i> | white wild indigo | 8 |
| 15 | <i>Berteroa incana</i> | hoary allysum | 0 |
| 16 | <i>Bouteloua curtipendula</i> | side oats grama | 6 |
| 17 | <i>Bromus inermis</i> | smooth brome | 0 |
| 18 | <i>Carex comosa</i> | bristly sedge | 4 |
| 19 | <i>Carex pensylvanica</i> *** | Pennsylvania sedge | 3 |
| 20 | <i>Centaurea biebersteinii</i> | spotted knapweed | 0 |
| 21 | <i>Chamecrista fasciculata</i> | partridge pea | 2 |
| 22 | <i>Cirsium vulgare</i> | bull thistle | 0 |
| 23 | <i>Cornus alba</i> | red osier dogwood | 3 |
| 24 | <i>Cornus racemosa</i> | gray dogwood | 2 |
| 25 | <i>Dactylis glomerata</i> | Orchard Grass | 0 |
| 26 | <i>Daucus carota</i> | Queen Anne's lace | 0 |
| 27 | <i>Echinacea purpurea</i> ** | purple coneflower | 10 |
| 28 | <i>Equisetum arvense</i> | field horsetail | 1 |
| 29 | <i>Erigeron strigosus</i> | daisy fleabane | 2 |
| 30 | <i>Euphorbia esula</i> | leafy spurge | 0 |
| 31 | <i>Eutrochium maculatum</i> | joe pye weed | 4 |
| 32 | <i>Frangula alnus</i> | glossy buckthorn | 0 |
| 33 | <i>Fraxinus pennsylvanica</i> | green ash | 2 |
| 34 | <i>Galium aparine</i> | cleavers | 1 |
| 35 | <i>Geum triflorum</i> | prairie smoke | 7 |
| 36 | <i>Glechoma hederacea</i> | ground ivy | 0 |
| 37 | <i>Gleditsia triacanthos</i> | honey locust | 0 |
| 38 | <i>Hackelia virginiana</i> | Virginia stickseed | 1 |
| 39 | <i>Helianthus grosseserratus</i> | sawtooth sunflower | 3 |
| 40 | <i>Hemerocallis sp.</i> | day lily | 0 |
| 41 | <i>Hosta sp.</i> | hosta | 0 |
| 42 | <i>Impatiens capensis</i> | jewelweed | 2 |
| 43 | <i>Juglens nigra</i> | black walnut | 4 |
| 44 | <i>Laportea canadensis</i> | wood nettle | 3 |
| 45 | <i>Leonorus cardiaca</i> | motherwort | 0 |
| 46 | <i>Leucanthemum vulgare</i> | daisy | 0 |
| 47 | <i>Lonicera tatarica</i> | Tatarian honeysuckle | 0 |
| 48 | <i>Lotus corniculatus</i> | bird's-foot trefoil | 0 |
| 49 | <i>Lythrum salicaria</i> | purple loosestrife | 0 |
| 50 | <i>Malus sp.</i> | apple | 0 |
| 51 | <i>Medicago lupulina</i> | black medic | 0 |
| 52 | <i>Medicago officinalis</i> | sweetclover | 0 |
| 53 | <i>Mentha arvensis</i> | wild mint | 3 |
| 54 | <i>Monarda fistulosa</i> | wild bergamot | 3 |
| 55 | <i>Nepeta cataria</i> | catnip | 0 |
| 56 | <i>Oligoneuron rigidum</i> | stiff goldenrod | 4 |
| 57 | <i>Panicum virgatum</i> | switchgrass | 2 |
| 58 | <i>Parthenocissus vitacea</i> | woodbine | 2 |
| 59 | <i>Phalaris arundinacea</i> | reed canarygrass | 0 |
| 60 | <i>Phleum pratense</i> | timothy | 0 |
| 61 | <i>Pinus resinosa</i> | red pine | 5 |
| 62 | <i>Plantago major</i> | common plantain | 0 |
| 63 | <i>Poa pratensis</i> | Kentucky bluegrass | 0 |
| 64 | <i>Populus deltoides</i> | cottonwood | 1 |
| 65 | <i>Potentilla simplex</i> | cinquefoil | 2 |
| 66 | <i>Ratibida pinnata</i> *** | globular coneflower | 4 |
| 67 | <i>Rhamnus cathartica</i> | common buckthorn | 0 |
| 68 | <i>Rhus hirta</i> | smooth sumac | 2 |
| 69 | <i>Ribes americanum</i> | wild black currant | 4 |
| 70 | <i>Rosa sp.</i> | rose | 0 |
| 71 | <i>Rubus idaeus</i> | red raspberry | 3 |
| 72 | <i>Rudbeckia hirta</i> | black eyed Susan | 3 |
| 73 | <i>Rumex crispus</i> | curly dock | 0 |
| 74 | <i>Salix amygdaloides</i> | peach leaf willow | 5 |
| 75 | <i>Salix interior</i> | sandbar willow | 2 |
| 76 | <i>Schizachyrium scoparium</i> | little bluestem | 5 |
| 77 | <i>Silene latifolia</i> | bladder campion | 0 |
| 78 | <i>Silphium perfoliatum</i> | cup plant | 4 |
| 79 | <i>Solanum dulcamara</i> | climbing nightshade | 0 |
| 80 | <i>Solidago canadensis</i> | Canada goldenrod | 1 |
| 81 | <i>Solidago flexicaulis</i> | zigzag goldenrod | 5 |
| 82 | <i>Solidago gigantea</i> | Late Goldenrod | 3 |
| 83 | <i>Solidago speciosa</i> | showy goldenrod | 5 |
| 84 | <i>Sonchus arvensis</i> | sow thistle | 0 |
| 85 | <i>Sorghastrum nutans</i> | Indian grass | 5 |
| 86 | <i>Taraxacum officinale</i> | common dandelion | 0 |
| 87 | <i>Trifolium hybridum</i> | alsike clover | 0 |
| 88 | <i>Trifolium pratense</i> | red clover | 0 |
| 89 | <i>Ulmus americana</i> | American elm | 3 |
| 90 | <i>Ulmus pumila</i> | Siberian elm | 0 |
| 91 | <i>Verbascum thapsus</i> | mullein | 0 |
| 92 | <i>Verbena hastata</i> | blue vervain | 6 |
| 93 | <i>Verbena stricta</i> *** | hoary vervain | 3 |
| 94 | <i>Vicia sativa</i> | vetch | 0 |
| 95 | <i>Viola sp. *</i> | violet | 6 |
| 96 | <i>Vitis riparia</i> | wild grape | 2 |
| 97 | <i>Zanthoxylum americanum</i> | common prickly ash | 3 |

Community #1

Eggers & Reed Plant Community Type: Shallow Open Water

Percent of AA Occupied by Type: 70

| Spp. # | Scientific Name | Common Name | Cover Class | CC Range | Midpoint CC | Native Status | Rapid FQA Stratum | NWI-GP | NWI-MW | NWI-NCNE | C | p | pC | |
|-----------|-------------------------|--------------------------|----------------|------------|-------------|---------------|----------------------|--------|--------|----------|------|------|-------|-------|
| 1 | Ceratophyllum demersum | Coon's-Tail | 4 | > 25 - 50% | 37.5 | Native | Aquatic | OBL | OBL | OBL | | 2 | 0.6 | 1.2 |
| 2 | Elodea canadensis | Canadian Waterweed | 2 | > 1 - 5% | 3 | Native | Aquatic | OBL | OBL | OBL | | 4 | 0.048 | 0.192 |
| 3 | Spirodela polyrhiza | Common Duckmeat | 1 | > 0 - 1% | 0.5 | Native | Aquatic | OBL | OBL | OBL | | 5 | 0.008 | 0.04 |
| 4 | Potamogeton crispus | Curly Pondweed | 3 | > 5 - 25% | 15 | Introduced | Aquatic | OBL | OBL | OBL | | 0 | 0.24 | 0 |
| 5 | Wolffia columbiana | Columbian Watermeal | 1 | > 0 - 1% | 0.5 | Native | Aquatic | OBL | OBL | OBL | | 5 | 0.008 | 0.04 |
| 6 | Ranunculus longirostris | Long-Beak Water-Crowfoot | 2 | > 1 - 5% | 3 | Native | Aquatic | OBL | OBL | OBL | | 7 | 0.048 | 0.336 |
| 7 | Stuckenia pectinata | Sago False Pondweed | 2 | > 1 - 5% | 3 | Native | Aquatic | OBL | OBL | OBL | | 3 | 0.048 | 0.144 |
| 8 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 9 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 10 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 11 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 12 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 13 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 14 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 15 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 16 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 17 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 18 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 19 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 20 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 21 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 22 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 23 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 24 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 25 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 26 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 27 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 28 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 29 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 30 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 31 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 32 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 33 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 34 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 35 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 36 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 37 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 38 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 39 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 40 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 41 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 42 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 43 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 44 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 45 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 46 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 47 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 48 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 49 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 50 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 51 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 52 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 53 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 54 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 55 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 56 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 57 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 58 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 59 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 60 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |

Community #2

Eggers & Reed Plant Community Type: **Shrub Carr**

Percent of AA Occupied by Type: **5**

| Spp. # | Scientific Name | Common Name | Cover Class | CC Range | Midpoint CC | Native Status | Rapid FQA Stratum | NWI-GP | NWI-MW | NWI-NCNE | C | p | pC |
|-----------|--------------------------------|--------------------------|----------------|------------|-------------|---------------|----------------------|--------|--------|----------|------|--------|---------------|
| 1 | Acer negundo | Ash-Leaf Maple | 2 | > 1 - 5% | 3 | Native | Tree | FAC | FAC | FAC | | 1 | 0.0179 0.0179 |
| 2 | Acer saccharinum | Silver Maple | 2 | > 1 - 5% | 3 | Native | Tree | FAC | FACW | FACW | | 3 | 0.0179 0.0537 |
| 3 | Fraxinus pennsylvanica | Green Ash | 2 | > 1 - 5% | 3 | Native | Tree | FAC | FACW | FACW | | 2 | 0.0179 0.0358 |
| 4 | Laportea canadensis | Canadian Wood-Nettle | 2 | > 1 - 5% | 3 | Native | Herb | FAC | FACW | FACW | | 3 | 0.0179 0.0537 |
| 5 | Lactuca serriola | Prickly Lettuce | 1 | > 0 - 1% | 0.5 | Introduced | Herb | FAC | FACU | FACU | | 0 | 0.003 0 |
| 6 | Parthenocissus inserta | Thicket-Creeper | 1 | > 0 - 1% | 0.5 | Native | Woody Vine | FAC | FACU | FACU | | 2 | 0.003 0.006 |
| 7 | Populus deltoides | Eastern Cottonwood | 2 | > 1 - 5% | 3 | Native | Tree | FAC | FAC | FAC | | 1 | 0.0179 0.0179 |
| 8 | Rhamnus cathartica | European Buckthorn | 3 | > 5 - 25% | 15 | Introduced | Shrub | FACU | FAC | FAC | | 0 | 0.0896 0 |
| 9 | Onoclea sensibilis | Sensitive Fern | 2 | > 1 - 5% | 3 | Native | Herb | FACW | FACW | FACW | | 4 | 0.0179 0.0716 |
| 10 | Urtica dioica | Stinging Nettle | 2 | > 1 - 5% | 3 | Native | Herb | FAC | FACW | FAC | | 1 | 0.0179 0.0179 |
| 11 | Salix amygdaloides | Peach-Leaf Willow | 4 | > 25 - 50% | 37.5 | Native | Tree | FACW | FACW | FACW | | 5 | 0.2239 1.1194 |
| 12 | Salix interior | Sandbar Willow | 4 | > 25 - 50% | 37.5 | Native | Shrub | FACW | FACW | FACW | | 2 | 0.2239 0.4478 |
| 13 | Salix petiolaris | Meadow Willow | 4 | > 25 - 50% | 37.5 | Native | Shrub | OBL | OBL | FACW | | 5 | 0.2239 1.1194 |
| 14 | Verbena hastata | Simpler's-Joy | 2 | > 1 - 5% | 3 | Native | Herb | FACW | FACW | FACW | | 6 | 0.0179 0.1075 |
| 15 | Schoenoplectus tabernaemontani | Soft-Stem Club-Rush | 2 | > 1 - 5% | 3 | Native | Herb | OBL | OBL | OBL | | 4 | 0.0179 0.0716 |
| 16 | Carex vulpinoidea | Common Fox Sedge | 2 | > 1 - 5% | 3 | Native | Herb | FACW | FACW | OBL | | 3 | 0.0179 0.0537 |
| 17 | Lycopus uniflorus | Northern Water-Horehound | 2 | > 1 - 5% | 3 | Native | Herb | OBL | OBL | OBL | | 5 | 0.0179 0.0896 |
| 18 | Rumex crispus | Curly Dock | 1 | > 0 - 1% | 0.5 | Introduced | Herb | FAC | FAC | FAC | | 0 | 0.003 0 |
| 19 | Taraxacum officinale | Common Dandelion | 1 | > 0 - 1% | 0.5 | Introduced | Herb | FACU | FACU | FACU | | 0 | 0.003 0 |
| 20 | | #N/A | 1 | > 0 - 1% | 0.5 | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | 0.003 | #N/A |
| 21 | | #N/A | 1 | > 0 - 1% | 0.5 | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | 0.003 | #N/A |
| 22 | | #N/A | 1 | > 0 - 1% | 0.5 | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | 0.003 | #N/A |
| 23 | | #N/A | 2 | > 1 - 5% | 3 | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | 0.0179 | #N/A |
| 24 | | #N/A | 1 | > 0 - 1% | 0.5 | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | 0.003 | #N/A |
| 25 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 26 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 27 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 28 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 29 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 30 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 31 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 32 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 33 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 34 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 35 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 36 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 37 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 38 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 39 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 40 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 41 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 42 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 43 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 44 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 45 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 46 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 47 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 48 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 49 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 50 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 51 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 52 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 53 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 54 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 55 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 56 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 57 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 58 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 59 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 60 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |

Community #3

Eggers & Reed Plant Community Type: **Fresh Meadow**

Percent of AA Occupied by Type: **5**

| Spp. | Scientific Name | Common Name | Cover | Rapid FQA | | | | | | | | | | |
|------|--------------------------------|--------------------------|-------|------------|-------------|-----------------|---------------|--------|--------|----------|------|------|--------|--------|
| # | | | Class | CC Range | Midpoint CC | Native Status | Stratum | NWI-GP | NWI-MW | NWI-NCNE | C | p | pC | |
| 1 | Ambrosia artemisiifolia | Annual Ragweed | 2 | > 1 - 5% | | 3 Native | Herb | FACU | FACU | FACU | | 0 | 0.0216 | 0 |
| 2 | Asclepias incarnata | Swamp Milkweed | 2 | > 1 - 5% | | 3 Native | Herb | FACW | OBL | OBL | | 4 | 0.0216 | 0.0863 |
| 3 | Carex comosa | Bearded Sedge | 2 | > 1 - 5% | | 3 Native | Herb | OBL | OBL | OBL | | 4 | 0.0216 | 0.0863 |
| 4 | Carex stricta | Uptight Sedge | 2 | > 1 - 5% | | 3 Native | Herb | OBL | OBL | OBL | | 5 | 0.0216 | 0.1079 |
| 5 | Carex vulpinoidea | Common Fox Sedge | 2 | > 1 - 5% | | 3 Native | Herb | FACW | FACW | OBL | | 3 | 0.0216 | 0.0647 |
| 6 | Cirsium arvense | Canadian Thistle | 2 | > 1 - 5% | | 3 Introduced | Herb | FACU | FACU | FACU | | 0 | 0.0216 | 0 |
| 7 | Cyperus esculentus | Chufa | 1 | > 0 - 1% | | 0.5 Introduced | Herb | FACW | FACW | FACW | | 0 | 0.0036 | 0 |
| 8 | Echinochloa crus-galli | Large Barnyard Grass | 2 | > 1 - 5% | | 3 Introduced | Herb | FAC | FACW | FAC | | 0 | 0.0216 | 0 |
| 9 | Eupatorium perfoliatum | Common Boneset | 2 | > 1 - 5% | | 3 Native | Herb | FACW | OBL | FACW | | 4 | 0.0216 | 0.0863 |
| 10 | Impatiens capensis | Spotted Touch-Me-Not | 2 | > 1 - 5% | | 3 Native | Herb | FACW | FACW | FACW | | 2 | 0.0216 | 0.0432 |
| 11 | Iris versicolor | Harlequin Blueflag | 1 | > 0 - 1% | | 0.5 Native | Herb | OBL | OBL | OBL | | 4 | 0.0036 | 0.0144 |
| 12 | Leersia oryzoides | Rice Cut Grass | 2 | > 1 - 5% | | 3 Native | Herb | OBL | OBL | OBL | | 3 | 0.0216 | 0.0647 |
| 13 | Lycopus uniflorus | Northern Water-Horehound | 1 | > 0 - 1% | | 0.5 Native | Herb | OBL | OBL | OBL | | 5 | 0.0036 | 0.018 |
| 14 | Lythrum salicaria | Purple Loosestrife | 3 | > 5 - 25% | | 15 Introduced | Herb | OBL | OBL | OBL | | 0 | 0.1079 | 0 |
| 15 | Onoclea sensibilis | Sensitive Fern | 1 | > 0 - 1% | | 0.5 Native | Herb | FACW | FACW | FACW | | 4 | 0.0036 | 0.0144 |
| 16 | Panicum virgatum | Wand Panic Grass | 2 | > 1 - 5% | | 3 Native | Herb | FAC | FAC | FAC | | 2 | 0.0216 | 0.0432 |
| 17 | Persicaria amphibia | Water Smartweed | 2 | > 1 - 5% | | 3 Native | Aquatic, Herb | OBL | OBL | OBL | | 4 | 0.0216 | 0.0863 |
| 18 | Phalaris arundinacea | Reed Canary Grass | 3 | > 5 - 25% | | 15 Introduced | Herb | FACW | FACW | FACW | | 0 | 0.1079 | 0 |
| 19 | Poa palustris | Fowl Blue Grass | 2 | > 1 - 5% | | 3 Native | Herb | FACW | FACW | FACW | | 5 | 0.0216 | 0.1079 |
| 20 | Rumex crispus | Curly Dock | 1 | > 0 - 1% | | 0.5 Introduced | Herb | FAC | FAC | FAC | | 0 | 0.0036 | 0 |
| 21 | Schoenoplectus tabernaemontani | Soft-Stem Club-Rush | 2 | > 1 - 5% | | 3 Native | Herb | OBL | OBL | OBL | | 4 | 0.0216 | 0.0863 |
| 22 | Scirpus cyperinus | Cottongrass Bulrush | 2 | > 1 - 5% | | 3 Native | Herb | OBL | OBL | OBL | | 3 | 0.0216 | 0.0647 |
| 23 | Solidago gigantea | Late Goldenrod | 3 | > 5 - 25% | | 15 Native | Herb | FAC | FACW | FACW | | 3 | 0.1079 | 0.3237 |
| 24 | Taraxacum officinale | Common Dandelion | 2 | > 1 - 5% | | 3 Introduced | Herb | FACU | FACU | FACU | | 0 | 0.0216 | 0 |
| 25 | Typha angustifolia | Narrow-Leaf Cat-Tail | 4 | > 25 - 50% | | 37.5 Introduced | Herb | OBL | OBL | OBL | | 0 | 0.2698 | 0 |
| 26 | Verbena hastata | Simpler's-Joy | 2 | > 1 - 5% | | 3 Native | Herb | FACW | FACW | FACW | | 6 | 0.0216 | 0.1295 |
| 27 | Zizia aurea | Golden Alexanders | 2 | > 1 - 5% | | 3 Native | Herb | FAC | FAC | FAC | | 6 | 0.0216 | 0.1295 |
| 28 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 29 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 30 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 31 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 32 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 33 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 34 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 35 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 36 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 37 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 38 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 39 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 40 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 41 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 42 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 43 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 44 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 45 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 46 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 47 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 48 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 49 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 50 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 51 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 52 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 53 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 54 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 55 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 56 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 57 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 58 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 59 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |
| 60 | | #N/A | | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A | #N/A |

Metric Summary & Community Assessments

| | Community #1 | Community #2 | Community #3 |
|--------------------------------|--------------------|--------------|--------------|
| Community Type | Shallow Open Water | Shrub Carr | Fresh Meadow |
| wC | 2.0 | 3.3 | 1.6 |
| Numerical Condition Category | 3 | 3 | 3 |
| Condition Category | Fair | Fair | Fair |
| Additional Metrics | | | |
| Native Species Richness | 6 | 15 | 19 |
| Introduced Species Richness | 1 | 4 | 8 |
| Mean C | 3.7 | 2.5 | 2.6 |
| FQI | 9.1 | 9.6 | 11.5 |
| Total Midpoint % Cover | 62.5 | 167.5 | 139 |
| Total Introduced Spp. Cover | 15 | 16.5 | 77.5 |
| Proportion of Introduced Cover | 0.24 | 0.10 | 0.56 |

Overall Assessment

| Community # | Community Type | wC | Condition Category | Numerical Category | Proportion of AA | Proportion x Numerical Category |
|-------------|--------------------|-----|--------------------|--------------------|------------------|---------------------------------|
| 1 | Shallow Open Water | 2.0 | Fair | 3 | 0.7 | 2.1 |
| 2 | Shrub Carr | 3.3 | Fair | 3 | 0.05 | 0.15 |
| 3 | Fresh Meadow | 1.6 | Fair | 3 | 0.05 | 0.15 |

Weighted Average Numerical Category for AA 2
Overall AA Condition Good

Appendix C

2003-2018 Habitat Assessment Monitoring Results

Appendix C: 2003-2009 Habitat Assessment Monitoring Results
Black Dog Watershed Management Organization

| Water Body | Monitoring Year | Approximate Proportion of the Water Body Which is Deep Water Habitat (~ > 20 ft. depth) | Vegetation Quality - Wet Areas | | | | | | | | | | | | | Vegetation Quality - Upland | | | | | | | | | | Erosion/Sedimentation | |
|------------|-----------------|---|--|--|--|---|----------------|-----|-----|---|---|---|---|----------------|--------|---|--|--|--|--|----------------|--------|--|--------------------------|----|-----------------------|-------|
| | | | Submergent Zone Sampling | | | | | | | Vegetated Emergent Zone Sampling | | | | | | Upland Buffer Sampling | | | | | | | | | | | |
| | | | Overall Submergent Vegetative Quality ¹ | Approximate Proportion of Water Body Typically Dominated By Submergent Vegetation (~ 2 - 20 ft. depth) | Average Native Plant Occurrence or Density Rating ^{2,3} | Total Number of Native Species ⁵ | Exotic Species | | | Emergent Zone Vegetative Quality ⁶ | Approximate Proportion of Emergent Zone (0 - 2 ft. depth) Within The Water Body | Approximate Total Percent Vegetative Cover Within The Entire Emergent Zone ⁷ | Total Number of Native Wetland Plant Species ⁸ | Exotic Species | | Overall Upland Buffer Quality ¹⁰ | Unmanicured Buffer Width ¹¹ | Estimated Total Vegetative Cover (Percent Range) ¹² | Total Number of Native Plant Species ¹³ | Buffer Continuity (Percent Surrounding Water Body) ¹⁴ | Exotic Species | | Shoreline Erosion (Percent of Shoreline) ¹⁶ | Sediment Deltas (Yes/No) | | | |
| | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Crystal | 2003 | 15% | Moderate | 80% | 1.5 | 15 | 2 | 1.1 | 1.1 | Moderate | 5% | 26-50% | 18 | 4 | 26-50% | Moderate | <10 ft. | >95% | 16 | 26-50% | 2 | 15-40% | 0-10% | No | | | |
| | 2004 | | Excellent | | 1.2 | 14 | 2 | 1.1 | 2.9 | Excellent | | 26-50% | 16 | 6 | 26-50% | Moderate | <10 ft. | >95% | 16 | 26-50% | 4 | 15-40% | 0-10% | No | | | |
| | 2005 | | Moderate | | 1.2 | 13 | 2 | 1.1 | 2.7 | Excellent | | 26-50% | 16 | 6 | 26-50% | Moderate | <10 ft. | >95% | 17 | 26-50% | 3 | 15-40% | 0-10% | No | | | |
| | 2006 | | Excellent | | 1.0 | 17 | 2 | 1.5 | 3.2 | Excellent | | 26-50% | 18 | 8 | 26-50% | Moderate | <10 ft. | >95% | 17 | 26-50% | 3 | 15-40% | 0-10% | No | | | |
| | 2007 | | Excellent | | 1.5 | 16 | 2 | 1.6 | 3.4 | Excellent | | 26-50% | 22 | 10 | 26-50% | Moderate | <10 ft. | >95% | 15 | 26-50% | 5 | 15-40% | 0-10% | No | | | |
| | 2008 | | Moderate | | 1.3 | 15 | 2 | 1.6 | 2.5 | Excellent | | 26-50% | 21 | 12 | 26-50% | Moderate | <10 ft. | >95% | 15 | 26-50% | 5 | 15-40% | 0-10% | No | | | |
| | 2009 | | Moderate | | 1.3 | 14 | 2 | 1.6 | 2.8 | Excellent | | 26-50% | 20 | 11 | 26-50% | Moderate | <10 ft. | >95% | 15 | 26-50% | 7 | 15-40% | 0-10% | No | | | |
| | Keller | | 2003 | | 0% | Moderate | 90% | 1.9 | 4 | 1 | | 3.2 | 3.2 | Poor | 10% | 51-75% | 5 | 2 | 51-75% | Moderate | <10 ft. | >95% | 7 | 76-100% | 6 | >40% | 0-10% |
| 2004 | | Moderate | 1.7 | 5 | | 1 | | 1.8 | 2.5 | Moderate | 51-75% | 6 | 2 | 51-75% | | Moderate | <10 ft. | >95% | 7 | 76-100% | 6 | >40% | 0-10% | No | | | |
| 2005 | | Moderate | 1.3 | 5 | | 2 | | 1.0 | 1.1 | Moderate | 51-75% | 7 | 2 | 26-50% | | Moderate | <10 ft. | >95% | 8 | 76-100% | 7 | >40% | 0-10% | No | | | |
| 2006 | | Moderate | 2.0 | 5 | | 2 | | 1.8 | 2.5 | Moderate | 51-75% | 8 | 2 | 26-50% | | Moderate | <10 ft. | >95% | 8 | 76-100% | 8 | >40% | 0-10% | No | | | |
| 2007 | | Moderate | 2.1 | 3 | | 2 | | 2.4 | 3.8 | Moderate | 51-75% | 9 | 3 | 26-50% | | Moderate | <10 ft. | >95% | 5 | 76-100% | 9 | >40% | 0-10% | No | | | |
| 2008 | | Moderate | 2.2 | 3 | | 2 | | 2.2 | 2.9 | Moderate | 51-75% | 9 | 3 | 26-50% | | Moderate | <10 ft. | >95% | 6 | 76-100% | 12 | >40% | 0-10% | No | | | |
| 2009 | | Poor | 3.0 | 2 | | 2 | | 2.7 | 3.3 | Moderate | 51-75% | 9 | 4 | 26-50% | | Moderate | <10 ft. | >95% | 4 | 76-100% | 11 | >40% | 0-10% | No | | | |
| Kingsley | | 2003 | 0% | Moderate | | 95% | | 2.7 | 7 | 0 | 0.0 | 0.0 | Excellent | 5% | | 51-75% | 11 | 2 | 0-25% | Excellent | 25-50 ft. | >95% | 15 | 51-75% | 4 | 15-40% | 0-10% |
| | 2004 | Moderate | | 2.7 | 7 | | 0 | 0.0 | 0.0 | Excellent | 51-75% | 11 | 2 | | 0-25% | Excellent | 25-50 ft. | >95% | 15 | 51-75% | 4 | 15-40% | 0-10% | No | | | |
| | 2005 | Moderate | | 2.6 | 7 | | 1 | 1.0 | 1.0 | Excellent | 51-75% | 15 | 6 | | 0-25% | Excellent | 25-50 ft. | >95% | 19 | 76-100% | 2 | 15-40% | 0-10% | No | | | |
| | 2006 | Excellent | | 1.8 ¹⁷ | 13 ¹⁸ | | 1 | 1.0 | 1.0 | Excellent | 51-75% | 15 | 6 | | 0-25% | Excellent | 25-50 ft. | >95% | 19 | 76-100% | 3 | 15-40% | 0-10% | No | | | |
| | 2007 | Excellent | | 1.6 | 13 | | 1 | 1.0 | 1.0 | Excellent | 51-75% | 19 | 6 | | 0-25% | Excellent | 25-50 ft. | >95% | 21 | 76-100% | 4 | 15-40% | 0-10% | No | | | |
| | 2008 | Moderate | | 2.9 | 5 | | 0 | 0.0 | 0.0 | Excellent | 51-75% | 18 | 5 | | 0-25% | Excellent | 25-50 ft. | >95% | 25 | 76-100% | 4 | 15-40% | 0-10% | No | | | |
| | 2009 | Excellent | | 2.0 | 11 | | 1 | 1.0 | 1.0 | Excellent | 51-75% | 16 | 5 | | 0-25% | Excellent | 25-50 ft. | >95% | 23 | 76-100% | 5 | 15-40% | 0-10% | No | | | |
| | Lac Lavon | 2003 | | 25% | Poor | | 70% | 2.0 | 7 | 1 | 1.0 | 1.0 | Poor | | 5% | 0-25% | 14 | 5 | 0-25% | Poor | <10 ft. | <75% | 12 | 0-25% | 17 | >40% | 0-10% |
| 2004 | | Moderate | 0.9 | | 9 | 2 | | 1.6 | 1.9 | Moderate | 0-25% | 15 | 5 | 0-25% | | Poor | <10 ft. | <75% | 12 | 0-25% | 17 | >40% | 0-10% | No | | | |
| 2005 | | Moderate | 2.3 | | 5 | 1 | | 2.0 | 2.0 | Excellent | 0-25% | 20 | 10 | 0-25% | | Poor | <10 ft. | <75% | 12 | 0-25% | 16 | >40% | 0-10% | No | | | |
| 2006 | | Moderate | 1.6 | | 10 ¹⁹ | 2 | | 2.5 | 4.0 | Excellent | 0-25% | 16 | 13 | 0-25% | | Poor | <10 ft. | <75% | 11 | 0-25% | 19 | >40% | 0-10% | No | | | |
| 2007 | | Excellent | 1.8 | | 10 ²⁰ | 3 | | 1.8 | 4.0 | Excellent | 0-25% | 16 | 12 | 0-25% | | Poor | <10 ft. | <75% | 12 | 0-25% | 18 | >40% | 0-10% | No | | | |
| 2008 | | Poor | 1.0 | | 5 | 2 | | 1.0 | 1.0 | Moderate | 0-25% | 14 | 9 | 0-25% | | Poor | <10 ft. | <75% | 9 | 0-25% | 13 | >40% | 0-10% | No | | | |
| 2009 | | Moderate | 1.6 | | 10 | 2 | | 2.5 | 4.0 | Moderate | 0-25% | 13 | 8 | 0-25% | | Poor | <10 ft. | <75% | 9 | 0-25% | 11 | >40% | 0-10% | No | | | |
| Orchard | | 2003 | 20% | | Poor | 75% | | 1.2 | 13 | 1 | 2.3 | 3.4 | Moderate | 5% | | 26-50% | 16 | 5 | 26-50% | Moderate | <10 ft. | >95% | 5 | 26-50% | 5 | >40% | 0-10% |
| | 2004 | Moderate | | 1.2 | 13 | | 1 | 2.3 | 2.3 | Excellent | 26-50% | 17 | 5 | | 26-50% | Moderate | <10 ft. | >95% | 5 | 26-50% | 5 | >40% | 0-10% | No | | | |
| | 2005 | Moderate | | 1.3 | 14 | | 1 | 1.8 | 2.6 | Moderate | 26-50% | 14 | 6 | | 26-50% | Moderate | <10 ft. | >95% | 5 | 26-50% | 5 | >40% | 0-10% | No | | | |
| | 2006 | Moderate | | 1.2 | 13 | | 1 | 1.7 | 3.4 | Excellent | 26-50% | 18 | 9 | | 26-50% | Moderate | <10 ft. | >95% | 5 | 26-50% | 5 | >40% | 0-10% | No | | | |
| | 2007 | Moderate | | 1.3 | 11 | | 1 | 1.9 | 3.3 | Excellent | 26-50% | 18 | 9 | | 26-50% | Moderate | <10 ft. | >95% | 3 | 26-50% | 5 | >40% | 0-10% | No | | | |
| | 2008 | Moderate | | 1.3 | 14 | | 1 | 1.6 | 2.8 | Excellent | 26-50% | 16 | 8 | | 26-50% | Moderate | <10 ft. | >95% | 3 | 26-50% | 7 | >40% | 0-10% | No | | | |
| | 2009 | Moderate | | 1.6 | 11 | | 1 | 1.7 | 2.5 | Excellent | 26-50% | 16 | 8 | | 26-50% | Moderate | <10 ft. | >95% | 3 | 26-50% | 6 | >40% | 0-10% | No | | | |
| | Sunset Pond | 2003 | | 0% | Moderate | | 75% | 3.0 | 11 | 1 | 1.0 | 1.0 | Poor | | 25% | 76 - 100% | 5 | 5 | 76-100% | Moderate | 10-25 ft. | 75-95% | 10 | 51-75% | 15 | 15-40% | 0-10% |
| 2004 | | Excellent | 2.2 | | 11 | 0 | | 0.0 | 0.0 | Poor | 76 - 100% | 4 | 3 | 76-100% | | Moderate | 10-25 ft. | 75-95% | 10 | 51-75% | 18 | 15-40% | 0-10% | Yes | | | |
| 2005 | | Excellent | 2.1 | | 10 | 1 | | 1.0 | 1.0 | Poor | 76 - 100% | 6 | 4 | 76-100% | | Moderate | 10-25 ft. | 75-95% | 9 | 76-100% | 20 | >40% | 0-10% | Yes | | | |
| 2006 | | Moderate | 2.6 | | 11 | 1 | | 1.0 | 1.0 | Poor | 76 - 100% | 7 | 4 | 76-100% | | Moderate | 10-25 ft. | 75-95% | 9 | 76-100% | 19 | >40% | 0-10% | No | | | |
| 2007 | | Excellent | 1.9 | | 12 | 1 | | 1.0 | 1.0 | Moderate | 76-100% | 11 | 6 | 76-100% | | Moderate | 10-25 ft. | 75-95% | 8 | 76-100% | 19 | >40% | 0-10% | No | | | |
| 2008 | | Excellent | 1.8 | | 10 | 1 | | 2.0 | 3.0 | Poor | 76-100% | 10 | 5 | 76-100% | | Moderate | 10-25 ft. | 75-95% | 5 | 76-100% | 15 | >40% | 0-10% | No | | | |
| 2009 | | Moderate | 2.2 | | 11 | 1 | | 3.0 | 3.0 | Poor | 76-100% | 10 | 5 | 76-100% | | Moderate | 10-25 ft. | 75-95% | 6 | 76-100% | 17 | >40% | 0-10% | Yes | | | |

Appendix C: 2003-2009 Habitat Assessment Monitoring Results
Black Dog Watershed Management Organization

The following footnotes pertain to 2003-2009 data.

¹Overall Submergent Vegetative Quality rating is the average of the exotic species density, macrophyte density, and total number of native: >0.66 = Excellent, 0.33-0.66 = Moderate, <0.33 = Poor.

| Overall Submergent Vegetative Quality | Avg. Exotic Species Density | Exotic Species Density/ Occurrence Rating Score | Avg. Macrophyte Density | Avg. Macrophyte Density Rating Score | Total Number of Native Species In Submergent Zone | Species Richness Rating | Total Overall Diversity Score |
|---------------------------------------|-----------------------------|---|----------------------------|--------------------------------------|---|-------------------------|-------------------------------|
| Poor | >2.0 | 0.1 | 0.0 - 1.0 and >3.0 | 0.1 | <9 | 0.1 | < 0.33 |
| Moderate | >0 - 2.0 | 0.5 | 1.0 - 1.5 and > 2.5 to 3.0 | 0.5 | 9-14 | .25-.75 | 0.33 - 0.66 |
| Excellent | 0 | 1.0 | 1.5 to 2.5 | 1.0 | >14 | 1.0 | > 0.66 |

²Plant occurrence ratings are a relative measure of the amount of native submergent vegetation with a scale from 1 to 5; 1 = lowest density (present on only 1 of 4 casts), 5 = highest density (hook full of vegetation on 4 of 4 casts).

³Density data for Crystal, Keller, and Orchard Lakes were collected by Blue Water Science. Numerous sample plots were conducted over the entire water body. A density scale of 1 to 4 was utilized (max = 4) by estimating the amount of vegetation obtained by rake casts and also transforming visual observations.

⁴Maximum exotic plant occurrence ratings represent the worst case scenario of curlyleaf pondweed density early in the growing season and/or Eurasian watermilfoil when it is most prolific later in the growing season.

⁵The Total Number of Native Species within the submergent zone for Crystal, Keller and Orchard Lakes is based on a detailed survey conducted by Blue Water Science; and for Kingsley Lake, Lac Lavon, and Sunset Pond, based on a survey by Barr Engineering and volunteers. The survey of the 3 water bodies conducted by Blue Water Science involved

the sampling of numerous sample plots or stations. The survey for Lac Lavon, Kingsley, and Sunset Pond is based on 3 sampling locations and a visual survey during travels on the water body: <7 = Poor, 7-14 = Moderate, >14 = Excellent.

⁶Emergent Zone Vegetative Quality is the average of the following parameters within the emergent zone: the approximate total percent coverage, the total number of native wetland species, and the percent coverage of exotic species: >0.66 = Excellent, 0.33-0.66 = Moderate, <0.33 = Poor.

| Emergent Zone Vegetative Quality | Percent Cover | Percent Cover Rating Score | | Total Number of Native Wetland Plants | Number of Native Wetland Plants Rating Score | Percent Cover of Exotics | Percent Cover of Exotics Rating Score | Overall Emergent Zone Quality Score |
|----------------------------------|-------------------|----------------------------|--|---------------------------------------|--|--------------------------|---------------------------------------|-------------------------------------|
| Poor | 0-25% | 0.1 | | <or= 5 | 0.1 | 76-100% | 0.1 | < 0.33 |
| Moderate | 76-100% or 26-50% | 0.5 | | >5 - 15 | 0.66 - 0.33 | 26-75% | .33-.66 | 0.33 - 0.66 |
| Excellent | 51-75% | 1.0 | | > 15 | 1.0 | 0-25% | 1.0 | > 0.66 |

⁷Approximate Total Percent Vegetative Cover Within the Entire Emergent Zone (0-2 ft. depth) is estimated based on the three sampling locations and a visual survey during travels around the water body. Estimates are broken into four categories: 0-25%=Poor, 26-50%=Moderate, 51-75%=Excellent, 76-100%=Moderate.

⁸The Total Number of Native Wetland Plant Species within the emergent zone is based on 3 sampling locations and a visual survey during travels on the water body: 0-5 = Poor, 6-15 = Moderate, >15 = Excellent.

⁹Total Exotic Emergent Percent Coverage, out of the entire emergent zone area, is estimated based on the three sampling locations and a visual survey during travels around the water body. Estimates are broken into four categories: 0-25%=Excellen(1.0), 26-50%=Moderate(0.5), 51-75%=Poor(0.0), 76-100%=Poor(0.1)

¹⁰Overall Upland Buffer Quality is determined based on the average of the four upland buffer quality parameters, with the exception of the number of exotic species present and the number of native plant species: >0.66 = Excellent, 0.33-0.66 = Moderate, <0.66 = Poor.

| Overall Upland Buffer Quality | Percent Cover | Percent Cover Rating Score | Exotics Percent Cover Range | Exotics Percent Cover Rating Score | Buffer Width Range | Buffer Width Rating Score | Buffer Continuity Percent Range | Buffer Continuity Rating Score | Overall Upland Buffer QualityScore |
|-------------------------------|---------------|----------------------------|-----------------------------|------------------------------------|--------------------|---------------------------|---------------------------------|--------------------------------|------------------------------------|
| Poor | <75% | 0.1 | >40% | 0.1 | <10 ft. | 0.1 | 0-25% | 0.1 | < 0.33 |
| Moderate | 75-95% | 0.5 | 15-40% | 0.5 | 10-50 ft. | 0.4 - 0.7 | 25-75% | .4-.7 | 0.33 - 0.66 |
| Excellent | >95% | 1.0 | <15% | 1.0 | >50 ft. | 1.0 | 76-100% | 1.0 | > 0.66 |

¹¹Unmanicured (upland) Buffer Width is divided into four categories: Excellent(1.0) = >50 ft, High(0.7) = 25-50 ft, Moderate(0.4) = 10-25 ft, and Low(0.1) = <10 ft.

¹²Estimated Total Vegetative Cover (Percent Range) for upland buffer is the proportion of the ground covered by vegetation within 50 feet of the wetland/upland transition zone. The percent cover is divided into three categories: Excellent(1.0) = >95%, Moderate(0.5) = 75 - 95%, and Poor(0.1) = <75%.

¹³The Total Number of Native Plant Species within the unmanicured upland buffer zone is based on 3 sampling locations and a visual survey.

¹⁴(Upland) Buffer Continuity is a measure of the proportion of the water body surrounded by the unmanicured, native upland buffer. This measure is divided into four categories: Excellent(1.0) = 76 - 100%, High(0.7) = 51 - 75%, Medium(0.4) = 26 - 50%, and Low(0.1) = 0 - 25%.

¹⁵Upland buffer exotic species "Percent of Total Coverage" is the percent cover of exotic species within the unmanicured upland buffer, which is divided into three categories: Excellent(1.0) = <15%, Moderate(0.5) = 15 - 40%, and Poor(0.1) = >40%.

¹⁶The presence of shoreline erosion is determined by the approximate percentage of the shoreline affected and is divided into the following three categories: 0 - 10%, 11 - 25%, 26 - 100%.

¹⁷The 2006 plant occurrence rating is lower (has improved), when compared to past assessment years primarily due to the low occurrence of additional plants found during a more detailed survey of the lake. The more detailed plant survey was conducted to better understand the extent of curlyleaf pondweed.

¹⁸The number of plant species documented in 2006, when compared to past assessment years, increased primarily due to additional plants found during a more detailed survey of the lake. The more detailed plant survey was conducted to better understand the extent of curlyleaf pondweed.

¹⁹The number of native submergent plant species documented in 2006, was incorrectly represented as 11 in the 2006 annual report. The actual number of native submergent plant species documented in 2006 was 10.

²⁰Native plant species were noted by the Minnesota Department of Natural Resouces during an October 25, 2007 macropyte survey and used in the 2007 annual report.

Rating Code:

Poor

Moderate

Excellent

Table 1: Orchard Lake 2012 and 2017 Habitat Assessment Monitoring Results Black Dog Watershed Management Organization

| Monitoring Year | Submergent Zone | | | | | | | | |
|-----------------|---|--|--|--|---|--|-------------------------|---|--|
| | Approximate Proportion of the Water Body Which is Deep Water Habitat (~ > 20 ft. depth) | Overall Submergent Zone Quality ¹ | Approximate Proportion of Water Body Typically Dominated By Submergent Vegetation (~ 2 - 20 ft. depth) | Native Species | | Mean Coefficient of Conservatism Value | Exotic Species | | |
| | | | | Average Native Plant Density Rating ^{2,3} | Total Number of Native Species ⁵ | | Total Number of Species | Average Exotic Plant Density Rating ^{2, 3} | Maximum Exotic Plant Density Rating ⁴ |
| 2012 | 20% | Moderate | 75% | 2.0 (Moderate) | 13 (High) | 5.4 (Moderate) | 1 | 1.7 (Moderate) | 3.0 (Poor) |
| 2017 | 20% | High | 75% | 1.2 (Excellent) | 16 (Excellent) | 5.2 (Moderate) | 2 | 1.1 (Moderate) | 1.5 (Moderate) |

| Monitoring Year | Emergent Zone | | | | | | |
|-----------------|--|---|---|---|--|-------------------|---|
| | Overall Emergent Zone Quality ⁶ | Approximate Proportion of Emergent Zone (0 - 2 ft. depth) Within The Water Body | Approximate Total Percent Vegetative Cover Within The Entire Emergent Zone ⁷ | Total Number of Native Wetland Plant Species ⁸ | Mean Coefficient of Conservatism Value | Exotic Species | |
| | | | | | | Number of Species | Total Exotic Emergent Percent Coverage ⁹ |
| 2012 | Moderate | 5% | 26-50% (Moderate) | 43 (Excellent) | 3.1 (Moderate) | 12 | 51-75% (Moderate) |
| 2017 | Moderate | 15% | 51-75% (High) | 50 (Excellent) | 2.7 (Poor) | 13 | 51-75% (Moderate) |

| Monitoring Year | Upland Buffer | | | | | | | | Erosion/Sedimentation | |
|-----------------|---|--|--|--|--|--|-------------------|---|--|--------------------------|
| | Overall Upland Buffer Quality ¹⁰ | Unmanicured Buffer Width ¹¹ | Estimated Total Vegetative Cover (Percent Range) ¹² | Total Number of Native Plant Species ¹³ | Mean Coefficient of Conservatism Value | Buffer Continuity (Percent Surrounding Water Body) ¹⁴ | Exotic Species | | Shoreline Erosion (Percent of Shoreline) ¹⁶ | Sediment Deltas (Yes/No) |
| | | | | | | | Number of Species | Percent of Total Coverage ¹⁵ | | |
| 2012 | Poor | <10 ft. (Poor) | >95% (High) | 19 (Moderate) | 1.6 (Poor) | 0-25% (Poor) | 20 | >40% (Poor) | 0-10% | No |
| 2017 | Moderate | <10 ft. (Poor) | >95% (High) | 25 (High) | 1.9 (Poor) | 0-25% (Poor) | 21 | >40% (Poor) | 0-10% | No |

Table 1: Orchard Lake 2017 Habitat Assessment Monitoring Results Black Dog Watershed Management Organization

The following changes were made to the 2011 - 2017 monitoring and analysis:

- Monitor one or two water bodies per year. Kingsley Lake in 2011 - Conduct a meandering survey of submergent, emergent, and upland buffer zones rather than monitoring of plot locations. Orchard Lake in 2012, Crystal Lake in 2013, Lac Lavon in 2014, Keller Lake in 2015, Kingsley Lake in 2016, Orchard Lake in 2017 - Conduct a meandering survey of submergent, emergent, and upland buffer zones. In addition, the emergent and upland buffer plot locations were evaluated.
- Changes were made in 2011 through 2017 to the calculations to include floristic quality as part of the assessment. These changes include adding a rating of "High" to the categories to accommodate MPCA ratings for floristic quality. These changes included adding a Rating Code:

Poor **Moderate** **High or Excellent**

The following footnotes pertain to 2011 through 2017 data:

¹**Overall Submergent Zone Quality** rating is the average of the rating scores for the following parameters: average exotic plant density, average native plant density, total number of native species, and C-value rating: >0.80 = Excellent, 0.67-0.80 = High, 0.33-0.66 = Moderate, <0.33 = Poor.

| Overall Submergent Zone Quality | Avg. Exotic Plant Density | Exotic Plant Density Rating Score | Avg. Native Plant Density | Avg. Native Plant Density Rating Score | Total Number of Native Species In Submergent Zone | Species Richness Rating Score | Mean Coefficient of Conservatism Value (C-Value) | C-Value Rating (using MPCA values, 2007) | Total Overall Submergent Zone Quality Score |
|---------------------------------|---------------------------|-----------------------------------|---------------------------|--|---|-------------------------------|--|--|---|
| Poor | >2.0 | 0.1 | > 1.75 | 0.1 | <7 | 0.1 | 0 - <3 | 0.10 | < 0.33 |
| Moderate | >1.0 - 2.0 | 0.5 | 1.25 - 1.75 | 0.5 | >7 - <9 | 0.5 | >3 - <6 | 0.50 | 0.33 - 0.66 |
| High | >0 - 1.0 | 0.75 | | | >9 - <14 | 0.75 | >6 - <9 | 0.75 | 0.67 - 0.80 |
| Excellent | 0 | 1.0 | 1.0 to 1.25 | 1.0 | >14 | 1.0 | >9 - 10 | 1.00 | > 0.80 |

²Plant density ratings are a relative measure of the total amount of submergent vegetation covering the submergent zone, with a scale from 1 to 4 according to MN DNR methodology. The rating system is based on a 1 to 3 scale. Therefore the density results were converted to match the rating system.

³Density data for Orchard Lake were collected by Blue Water Science using a stratified line transect survey throughout the lake.

⁴Maximum exotic plant density ratings represent the worst case scenario of curlyleaf pondweed density early in the growing season and/or Eurasian watermilfoil when it is most prolific later in the growing season.

⁵The Total Number of Native Species within the submergent zone for Orchard Lake was collected by Blue Water Science using a stratified line transect survey. The additional category of "High" was added in 2011 through 2017 and values were adjusted to: <7 = Poor, 7-9 = Moderate, 9-14 = High, >14 = Excellent.

⁶**Overall Emergent Zone Quality** is the average of the rating scores for the following parameters within the emergent zone: the total percent coverage, the total number of native wetland plant species, the percent coverage of exotic species, and the C-Value Rating: >0.80 = Excellent, 0.67-0.80 = High, 0.33-0.66 = Moderate, <0.33 = Poor.

| Overall Emergent Zone Quality | Percent Cover | Percent Cover Rating Score | Total Number of Native Wetland Plant Species | Number of Native Wetland Plant Species Rating Score | Percent Cover of Exotics | Percent Cover of Exotics Rating Score | Mean Coefficient of Conservatism Value (C-Value) | C-Value Rating (using MPCA values, 2007) | Overall Emergent Zone Quality Score |
|-------------------------------|-------------------|----------------------------|--|---|--------------------------|---------------------------------------|--|--|-------------------------------------|
| Poor | 0-25% | 0.1 | < or= 5 | 0.1 | 76-100% | 0.1 | 0 - <3 | 0.10 | < 0.33 |
| Moderate | 76-100% or 26-50% | 0.5 | 6 - 10 | 0.33 | 51-75% | 0.33 | >3 - <6 | 0.50 | 0.33 - 0.66 |
| High | 51-75% | 1.0 | 11 - 15 | 0.66 | 26-50% | 0.66 | >6 - <9 | 0.75 | 0.67 - 0.80 |
| Excellent | 51-75% | 1.0 | > 15 | 1.0 | 0-25% | 1.0 | >9 - 10 | 1.00 | > 0.80 |

Table 1: Orchard Lake 2017 Habitat Assessment Monitoring Results Black Dog Watershed Management Organization

⁷Approximate Total Percent Vegetative Cover Within the Entire Emergent Zone (0-2 ft. depth) is estimated based on the three sampling locations and a visual survey during travels around the water body. Estimates are broken into the following categories: 0-25%=Poor, 26-50%=Moderate, 51-75%=High and Excellent, 76-100%=Moderate.

⁸The Total Number of Native Wetland Plant Species within the emergent zone is based on 3 sampling locations, a meandering visual survey during travels on the water body, and walking along the shoreline: 0-5 = Poor, 6-10 = Moderate, 11-15 = High, and >15 = Excellent.

⁹Total Exotic Emergent Percent Coverage, out of the entire emergent zone area, is estimated based on two plot locations, a meandering visual survey during travels on the water body, and walking along the shoreline. Estimates are broken into four categories: 0-25%=Excellent (1.0), 26-50%=High (0.66), 51-75%=Moderate (0.33), 76-100%=Poor (0.1)

¹⁰**Overall Upland Buffer Quality** is determined based on the average of the six upland buffer quality parameter rating scores: >0.80 = Excellent, 0.67-0.80 = High, 0.33-0.66 = Moderate, <0.33 = Poor.

| Overall Upland Buffer Quality | Percent Cover | Percent Cover Rating Score | Exotics Percent Cover Range | Exotics Percent Cover Rating Score | Buffer Width Range | Buffer Width Rating Score | Buffer Continuity Percent Range | Buffer Continuity Rating Score | Mean Coefficient of Conservatism Value (C-Value) | C-Value Rating (using MPCA values, 2007) | Number of Native Species | Number of Native Species Rating Score | Overall Upland Buffer Quality Score |
|--------------------------------------|----------------------|-----------------------------------|------------------------------------|---|---------------------------|----------------------------------|--|---------------------------------------|---|---|---------------------------------|--|--|
| Poor | <75% | 0.1 | >40% | 0.1 | <10 ft. | 0.1 | 0-25% | 0.1 | 0 - <3 | 0.10 | <5 | 0.1 | < 0.33 |
| Moderate | 75-95% | 0.5 | 15-40% | 0.5 | 10-25 ft. | 0.4 | 25-50% | 0.4 | >3 - <6 | 0.50 | 5-20 | 0.33 | 0.33 - 0.66 |
| High | >95% | 1.0 | <15% | 1.0 | 25-50 ft. | 0.7 | 51-75% | 0.7 | >6 - <9 | 0.75 | 20-30 | 0.66 | 0.67 - 0.80 |
| Excellent | >95% | 1.0 | <15% | 1.0 | >50 ft. | 1.0 | 76-100% | 1.0 | >9 - 10 | 1.00 | >30 | 1.0 | > 0.80 |

¹¹Unmanicured (upland) Buffer Width is divided into four categories: Excellent (1.0) = >50 ft, High (0.7) = 25-50 ft, Moderate (0.4) = 10-25 ft, and Low (0.1) = <10 ft.

¹²Estimated Total Vegetative Cover (Percent Range) for upland buffer is the proportion of the ground covered by vegetation within 50 feet of the wetland/upland transition zone. The percent cover is divided into three categories: High and Excellent (1.0) = >95%, Moderate (0.5) = 75 - 95%, and Poor (0.1) = <75%.

¹³The Total Number of Native Plant Species within the unmanicured upland buffer zone is based on two plot locations and a meandering visual survey along the shoreline.

¹⁴(Upland) Buffer Continuity is a measure of the proportion of the water body surrounded by the unmanicured, native upland buffer. This measure is divided into four categories: Excellent (1.0) = 76 - 100%, High (0.7) = 51 - 75%, Medium (0.4) = 26 - 50%, and Low (0.1) = 0 - 25%.

¹⁵Upland buffer exotic species "Percent of Total Coverage" is the percent cover of exotic species within the unmanicured upland buffer, which is divided into three categories: High and Excellent (1.0) = <15%, Moderate (0.5) = 15 - 40%, and Poor (0.1) = >40%.

¹⁶The presence of shoreline erosion is determined by the approximate percentage of the shoreline affected and is divided into the following three categories: 0 - 10%, 11 - 25%, 26 - 100%.

Table 1: Crystal Lake 2013 and 2018 Habitat Assessment Monitoring Results Black Dog Watershed Management Organization

| Monitoring Year | Submergent Zone Sampling | | | | | | | | |
|-----------------|---|--|--|--|---|--|-------------------------|---|--|
| | Approximate Proportion of the Water Body Which is Deep Water Habitat (~ > 20 ft. depth) | Overall Submergent Zone Quality ¹ | Approximate Proportion of Water Body Typically Dominated By Submergent Vegetation (~ 2 - 20 ft. depth) | Native Species | | Mean Coefficient of Conservatism Value | Exotic Species | | |
| | | | | Average Native Plant Density Rating ^{2,3} | Total Number of Native Species ⁵ | | Total Number of Species | Average Exotic Plant Density Rating ^{2, 3} | Maximum Exotic Plant Density Rating ⁴ |
| 2013 | 15% | High | 80% | 1.2 (Excellent) | 18 (Excellent) | 4.9 (Moderate) | 2 | 1.8 (Moderate) | 2.2 (Poor) |
| 2018 | 15% | High | 80% | 1.2 (Excellent) | 15 (Excellent) | 5.0 (Moderate) | 2 | 1.2 (Moderate) | 1.4 (Moderate) |

| Monitoring Year | Vegetated Emergent Zone Sampling | | | | | | |
|-----------------|--|---|---|---|--|-------------------|---|
| | Overall Emergent Zone Quality ⁶ | Approximate Proportion of Emergent Zone (0 - 2 ft. depth) Within The Water Body | Approximate Total Percent Vegetative Cover Within The Entire Emergent Zone ⁷ | Total Number of Native Wetland Plant Species ⁸ | Mean Coefficient of Conservatism Value | Exotic Species | |
| | | | | | | Number of Species | Total Exotic Emergent Percent Coverage ⁹ |
| 2013 | High | 5% | 26-50% (Moderate) | 36 (Excellent) | 3.0 (Moderate) | 10 | 26-50% (High) |
| 2018 | High | 5% | 26-50% (Moderate) | 50 (Excellent) | 3.3 (Moderate) | 9 | 26-50% (High) |

| Monitoring Year | Upland Buffer Sampling | | | | | | | | Erosion/Sedimentation | |
|-----------------|---|--|--|--|--|--|-------------------|---|--|--------------------------|
| | Overall Upland Buffer Quality ¹⁰ | Unmanicured Buffer Width ¹¹ | Estimated Total Vegetative Cover (Percent Range) ¹² | Total Number of Native Plant Species ¹³ | Mean Coefficient of Conservatism Value | Buffer Continuity (Percent Surrounding Water Body) ¹⁴ | Exotic Species | | Shoreline Erosion (Percent of Shoreline) ¹⁶ | Sediment Deltas (Yes/No) |
| | | | | | | | Number of Species | Percent of Total Coverage ¹⁵ | | |
| 2013 | Moderate | <10 ft. (Poor) | >95% (High) | 39 (Excellent) | 2.6 (Poor) | 26-50% (Moderate) | 16 | 15-40% (Moderate) | 0-10% | No |
| 2018 | Moderate | <10 ft. (Poor) | >95% (High) | 54 (Excellent) | 2.7 (Poor) | 26-50% (Moderate) | 20 | 15-40% (Moderate) | 0-10% | No |

Table 1: Crystal Lake 2018 Habitat Assessment Monitoring Results Black Dog Watershed Management Organization

The following changes were made to the 2011 - 2018 monitoring and analysis:

- Monitor one or two water bodies per year. Kingsley Lake in 2011 and 2016, Orchard Lake in 2012 and 2017, Crystal Lake in 2013 and 2018, Lac Lavon in 2014, Keller Lake in 2015 - Conduct a meandering survey of submergent, emergent, and upland buffer zones. In addition, the emergent and upland buffer plot locations were evaluated.
- Changes were made in 2011 through 2018 to the calculations to include floristic quality as part of the assessment. These changes include adding a rating of "High" to the categories to accommodate MPCA ratings for floristic quality. These changes included adding a Rating Code:

| | | |
|------|----------|-------------------|
| Poor | Moderate | High or Excellent |
|------|----------|-------------------|

The following footnotes pertain to 2011 through 2018 data:

¹**Overall Submergent Zone Quality** rating is the average of the rating scores for the following parameters: average exotic plant density, average native plant density, total number of native species, and C-value rating: >0.80 = Excellent, 0.67-0.80 = High, 0.33-0.66 = Moderate, <0.33 = Poor.

| Overall Submergent Zone Quality | Avg. Exotic Plant Density | Exotic Plant Density Rating Score | Avg. Native Plant Density | Avg. Native Plant Density Rating Score | Total Number of Native Species In Submergent Zone | Species Richness Rating Score | Mean Coefficient of Conservatism Value (C-Value) | C-Value Rating (using MPCA values, 2007) | Total Overall Submergent Zone Quality Score |
|---------------------------------|---------------------------|-----------------------------------|---------------------------|--|---|-------------------------------|--|--|---|
| Poor | >2.0 | 0.1 | > 1.75 | 0.1 | <7 | 0.1 | 0 - <3 | 0.10 | < 0.33 |
| Moderate | >1.0 - 2.0 | 0.5 | 1.25 - 1.75 | 0.5 | >7 - <9 | 0.5 | >3 - <6 | 0.50 | 0.33 - 0.66 |
| High | >0 - 1.0 | 0.75 | | | >9 - <14 | 0.75 | >6 - <9 | 0.75 | 0.67 - 0.80 |
| Excellent | 0 | 1.0 | 1.0 to 1.25 | 1.0 | >14 | 1.0 | >9 - 10 | 1.00 | > 0.80 |

²Plant density ratings are a relative measure of the total amount of submergent vegetation covering the submergent zone, with a scale from 1 to 3.

³Density data for Orchard Lake were collected by Blue Water Science using a stratified line transect survey throughout the lake.

⁴Maximum exotic plant density ratings represent the worst case scenario of curlyleaf pondweed density early in the growing season and/or Eurasian watermilfoil when it is most prolific later in the growing season.

⁵The Total Number of Native Species within the submergent zone for Orchard Lake was collected by Blue Water Science using a stratified line transect survey. The additional category of "High" was added in 2011 through 2018 and values were adjusted to: <7 = Poor, 7-9 = Moderate, 9-14 = High, >14 = Excellent.

⁶**Overall Emergent Zone Quality** is the average of the rating scores for the following parameters within the emergent zone: the total percent coverage, the total number of native wetland plant species, the percent coverage of exotic species, and the C-Value Rating: >0.80 = Excellent, 0.67-0.80 = High, 0.33-0.66 = Moderate, <0.33 = Poor.

| Overall Emergent Zone Quality | Percent Cover | Percent Cover Rating Score | Total Number of Native Wetland Plant Species | Number of Native Wetland Plant Species Rating Score | Percent Cover of Exotics | Percent Cover of Exotics Rating Score | Mean Coefficient of Conservatism Value (C-Value) | C-Value Rating (using MPCA values, 2007) | Overall Emergent Zone Quality Score |
|-------------------------------|-------------------|----------------------------|--|---|--------------------------|---------------------------------------|--|--|-------------------------------------|
| Poor | 0-25% | 0.1 | < or= 5 | 0.1 | 76-100% | 0.1 | 0 - <3 | 0.10 | < 0.33 |
| Moderate | 76-100% or 26-50% | 0.5 | 6 - 10 | 0.33 | 51-75% | 0.33 | >3 - <6 | 0.50 | 0.33 - 0.66 |
| High | 51-75% | 1.0 | 11 - 15 | 0.66 | 26-50% | 0.66 | >6 - <9 | 0.75 | 0.67 - 0.80 |
| Excellent | 51-75% | 1.0 | > 15 | 1.0 | 0-25% | 1.0 | >9 - 10 | 1.00 | > 0.80 |

Table 1: Crystal Lake 2018 Habitat Assessment Monitoring Results Black Dog Watershed Management Organization

⁷Approximate Total Percent Vegetative Cover Within the Entire Emergent Zone (0-2 ft. depth) is estimated based on the three sampling locations and a visual survey during travels around the water body. Estimates are broken into the following categories: 0-25%=Poor, 26-50%=Moderate, 51-75%=High and Excellent, 76-100%=Moderate.

⁸The Total Number of Native Wetland Plant Species within the emergent zone is based on 3 sampling locations, a meandering visual survey during travels on the water body, and walking along the shoreline: 0-5 = Poor, 6-10 = Moderate, 11-15 = High, and >15 = Excellent.

⁹Total Exotic Emergent Percent Coverage, out of the entire emergent zone area, is estimated based on two plot locations, a meandering visual survey during travels on the water body, and walking along the shoreline. Estimates are broken into four categories: 0-25%=Excellent (1.0), 26-50%=High (0.66), 51-75%=Moderate (0.33), 76-100%=Poor (0.1)

¹⁰**Overall Upland Buffer Quality** is determined based on the average of the six upland buffer quality parameter rating scores: >0.80 = Excellent, 0.67-0.80 = High, 0.33-0.66 = Moderate, <0.33 = Poor.

| Overall Upland Buffer Quality | Percent Cover | Percent Cover Rating Score | Exotics Percent Cover Range | Exotics Percent Cover Rating Score | Buffer Width Range | Buffer Width Rating Score | Buffer Continuity Percent Range | Buffer Continuity Rating Score | Mean Coefficient of Conservatism Value (C-Value) | C-Value Rating (using MPCA values, 2007) | Number of Native Species | Number of Native Species Rating Score | Overall Upland Buffer Quality Score |
|--------------------------------------|----------------------|-----------------------------------|------------------------------------|---|---------------------------|----------------------------------|--|---------------------------------------|---|---|---------------------------------|--|--|
| Poor | <75% | 0.1 | >40% | 0.1 | <10 ft. | 0.1 | 0-25% | 0.1 | 0 - <3 | 0.10 | <5 | 0.1 | < 0.33 |
| Moderate | 75-95% | 0.5 | 15-40% | 0.5 | 10-25 ft. | 0.4 | 25-50% | 0.4 | >3 - <6 | 0.50 | 5-20 | 0.33 | 0.33 - 0.66 |
| High | >95% | 1.0 | <15% | 1.0 | 25-50 ft. | 0.7 | 51-75% | 0.7 | >6 - <9 | 0.75 | 20-30 | 0.66 | 0.67 - 0.80 |
| Excellent | >95% | 1.0 | <15% | 1.0 | >50 ft. | 1.0 | 76-100% | 1.0 | >9 - 10 | 1.00 | >30 | 1.0 | > 0.80 |

¹¹Unmanicured (upland) Buffer Width is divided into four categories: Excellent (1.0) = >50 ft, High (0.7) = 25-50 ft, Moderate (0.4) = 10-25 ft, and Low (0.1) = <10 ft.

¹²Estimated Total Vegetative Cover (Percent Range) for upland buffer is the proportion of the ground covered by vegetation within 50 feet of the wetland/upland transition zone. The percent cover is divided into three categories: High and Excellent (1.0) = >95%, Moderate (0.5) = 75 - 95%, and Poor (0.1) = <75%.

¹³The Total Number of Native Plant Species within the unmanicured upland buffer zone is based on two plot locations and a meandering visual survey along the shoreline.

¹⁴(Upland) Buffer Continuity is a measure of the proportion of the water body surrounded by the unmanicured, native upland buffer. This measure is divided into four categories: Excellent (1.0) = 76 - 100%, High (0.7) = 51 - 75%, Medium (0.4) = 26 - 50%, and Low (0.1) = 0 - 25%.

¹⁵Upland buffer exotic species "Percent of Total Coverage" is the percent cover of exotic species within the unmanicured upland buffer, which is divided into three categories: High and Excellent (1.0) = <15%, Moderate (0.5) = 15 - 40%, and Poor (0.1) = >40%.

¹⁶The presence of shoreline erosion is determined by the approximate percentage of the shoreline affected and is divided into the following three categories: 0 - 10%, 11 - 25%, 26 - 100%.

Table 1: Keller Lake 2015 Habitat Assessment Monitoring Results Black Dog Watershed Management Organization

| Submergent Zone | | | | | | | | |
|---|--|--|--|---|--|-------------------------|--|--|
| Approximate Proportion of the Water Body Which is Deep Water Habitat (~ > 20 ft. depth) | Overall Submergent Zone Quality ¹ | Approximate Proportion of Water Body Typically Dominated By Submergent Vegetation (~ 2 - 20 ft. depth) | Native Species | | Mean Coefficient of Conservatism Value | Exotic Species | | |
| | | | Average Native Plant Density Rating ^{2,3} | Total Number of Native Species ⁵ | | Total Number of Species | Average Exotic Plant Density Rating ^{2,3} | Maximum Exotic Plant Density Rating ⁴ |
| 0% | Poor | 90% | 1.3 (Moderate) | 2 (Poor) | 1.5 (Poor) | 2 | 1.8 (Moderate) | 2.2 |

| Emergent Zone | | | | | | |
|--|---|---|---|--|-------------------|---|
| Overall Emergent Zone Quality ⁶ | Approximate Proportion of Emergent Zone (0 - 2 ft. depth) Within The Water Body | Approximate Total Percent Vegetative Cover Within The Entire Emergent Zone ⁷ | Total Number of Native Wetland Plant Species ⁸ | Mean Coefficient of Conservatism Value | Exotic Species | |
| | | | | | Number of Species | Total Exotic Emergent Percent Coverage ⁹ |
| High | 10% | 51-75% (High) | 28 (Excellent) | 2.3 (Poor) | 8 | 26-50% (High) |

| Upland Buffer | | | | | | | | Erosion/Sedimentation | |
|---|--|--|--|--|--|-------------------|---|--|--------------------------|
| Overall Upland Buffer Quality ¹⁰ | Unmanicured Buffer Width ¹¹ | Estimated Total Vegetative Cover (Percent Range) ¹² | Total Number of Native Plant Species ¹³ | Mean Coefficient of Conservatism Value | Buffer Continuity (Percent Surrounding Water Body) ¹⁴ | Exotic Species | | Shoreline Erosion (Percent of Shoreline) ¹⁶ | Sediment Deltas (Yes/No) |
| | | | | | | Number of Species | Percent of Total Coverage ¹⁵ | | |
| Moderate | 25-50 ft. (High) | >95% (High) | 20 (Moderate) | 1.6 (Poor) | 76-100% (Excellent) | 10 | >40% (Poor) | 0-10% | No |

Table 1: Keller Lake 2015 Habitat Assessment Monitoring Results Black Dog Watershed Management Organization

The following changes were made to the 2011 - 2015 monitoring and analysis:

- Monitor one or two water bodies per year. Kingsley Lake only in 2011 - Conduct a meandering survey of submergent, emergent, and upland buffer zones rather than monitoring of plot locations. Orchard Lake only in 2012, Crystal Lake only in 2013, Lac Lavon only in 2014, Keller Lake only in 2015 - Conduct a meandering survey of submergent, emergent, and upland buffer zones. In addition, the emergent and upland buffer plot locations were evaluated.
- Changes were made in 2011 through 2015 to the calculations to include floristic quality as part of the assessment. These changes include adding a rating of "High" to the categories to accommodate MPCA ratings for floristic quality. These changes included adding a Rating Code:

| | | |
|------|----------|-------------------|
| Poor | Moderate | High or Excellent |
|------|----------|-------------------|

The following footnotes pertain to 2015 data:

¹**Overall Submergent Zone Quality** rating is the average of the rating scores for the following parameters: average exotic plant density, average native plant density, total number of native species, and C-value rating: >0.80 = Excellent, 0.67-0.80 = High, 0.33-0.66 = Moderate, <0.33 = Poor.

| Overall Submergent Zone Quality | Avg. Exotic Plant Density | Exotic Plant Density Rating Score | Avg. Native Plant Density | Avg. Native Plant Density Rating Score | Total Number of Native Species In Submergent Zone | Species Richness Rating Score | Mean Coefficient of Conservatism Value (C-Value) | C-Value Rating (using MPCA values, 2007) | Total Overall Submergent Zone Quality Score |
|---------------------------------|---------------------------|-----------------------------------|---------------------------|--|---|-------------------------------|--|--|---|
| Poor | >2.0 | 0.1 | > 1.75 | 0.1 | <7 | 0.1 | 0 - <3 | 0.10 | < 0.33 |
| Moderate | >1.0 - 2.0 | 0.5 | 1.25 - 1.75 | 0.5 | >7 - <9 | 0.5 | >3 - <6 | 0.50 | 0.33 - 0.66 |
| High | >0 - 1.0 | 0.75 | | | >9 - <14 | 0.75 | >6 - <9 | 0.75 | 0.67 - 0.80 |
| Excellent | 0 | 1.0 | 1.0 to 1.25 | 1.0 | >14 | 1.0 | >9 - 10 | 1.00 | > 0.80 |

²Plant density ratings are a relative measure of the total amount of submergent vegetation covering the submergent zone, with a scale from 1 to 3 utilizing a 6-tined hook; 1 = light density (plant species found on only 1 tine), 2 = moderate density (plant species found on 2 to 4 tines), 3 = heavy density (plant species found on 5 or 6 tines).

³Density data for Keller Lake were collected by Blue Water Science using a point intercept survey throughout the lake.

⁴Maximum exotic plant density ratings represent the worst case scenario of curlyleaf pondweed density early in the growing season and/or Eurasian watermilfoil when it is most prolific later in the growing season.

⁵The Total Number of Native Species within the submergent zone for Keller Lake was collected by Blue Water Science using a point intercept survey. The additional category of "High" was added in 2011 through 2015 and values were adjusted to: <7 = Poor, 7-9 = Moderate, 9-14 = High, >14 = Excellent.

⁶**Overall Emergent Zone Quality** is the average of the rating scores for the following parameters within the emergent zone: the total percent coverage, the total number of native wetland plant species, the percent coverage of exotic species, and the C-Value Rating: >0.80 = Excellent, 0.67-0.80 = High, 0.33-0.66 = Moderate, <0.33 = Poor.

| Overall Emergent Zone Quality | Percent Cover | Percent Cover Rating Score | Total Number of Native Wetland Plant Species | Number of Native Wetland Plant Species Rating Score | Percent Cover of Exotics | Percent Cover of Exotics Rating Score | Mean Coefficient of Conservatism Value (C-Value) | C-Value Rating (using MPCA values, 2007) | Overall Emergent Zone Quality Score |
|-------------------------------|-------------------|----------------------------|--|---|--------------------------|---------------------------------------|--|--|-------------------------------------|
| Poor | 0-25% | 0.1 | < or= 5 | 0.1 | 76-100% | 0.1 | 0 - <3 | 0.10 | < 0.33 |
| Moderate | 76-100% or 26-50% | 0.5 | 6 - 10 | 0.33 | 51-75% | 0.33 | >3 - <6 | 0.50 | 0.33 - 0.66 |
| High | 51-75% | 1.0 | 11 - 15 | 0.66 | 26-50% | 0.66 | >6 - <9 | 0.75 | 0.67 - 0.80 |
| Excellent | 51-75% | 1.0 | > 15 | 1.0 | 0-25% | 1.0 | >9 - 10 | 1.00 | > 0.80 |

Table 1: Keller Lake 2015 Habitat Assessment Monitoring Results Black Dog Watershed Management Organization

⁷Approximate Total Percent Vegetative Cover Within the Entire Emergent Zone (0-2 ft. depth) is estimated based on the three sampling locations and a visual survey during travels around the water body. Estimates are broken into the following categories: 0-25%=Poor, 26-50%=Moderate, 51-75%=High and Excellent, 76-100%=Moderate.

⁸The Total Number of Native Wetland Plant Species within the emergent zone is based on 3 sampling locations, a meandering visual survey during travels on the water body, and walking along the shoreline: 0-5 = Poor, 6-10 = Moderate, 11-15 = High, and >15 = Excellent.

⁹Total Exotic Emergent Percent Coverage, out of the entire emergent zone area, is estimated based on two plot locations, a meandering visual survey during travels on the water body, and walking along the shoreline. Estimates are broken into four categories: 0-25%=Excellent (1.0), 26-50%=High (0.66), 51-75%=Moderate (0.33), 76-100%=Poor (0.1)

¹⁰**Overall Upland Buffer Quality** is determined based on the average of the six upland buffer quality parameter rating scores: >0.80 = Excellent, 0.67-0.80 = High, 0.33-0.66 = Moderate, <0.33 = Poor.

| Overall Upland Buffer Quality | Percent Cover | Percent Cover Rating Score | Exotics Percent Cover Range | Exotics Percent Cover Rating Score | Buffer Width Range | Buffer Width Rating Score | Buffer Continuity Percent Range | Buffer Continuity Rating Score | Mean Coefficient of Conservatism Value (C-Value) | C-Value Rating (using MPCA values, 2007) | Number of Native Species | Number of Native Species Rating Score | Overall Upland Buffer Quality Score |
|--------------------------------------|----------------------|-----------------------------------|------------------------------------|---|---------------------------|----------------------------------|--|---------------------------------------|---|---|---------------------------------|--|--|
| Poor | <75% | 0.1 | >40% | 0.1 | <10 ft. | 0.1 | 0-25% | 0.1 | 0 - <3 | 0.10 | <5 | 0.1 | < 0.33 |
| Moderate | 75-95% | 0.5 | 15-40% | 0.5 | 10-25 ft. | 0.4 | 25-50% | 0.4 | >3 - <6 | 0.50 | 5-20 | 0.33 | 0.33 - 0.66 |
| High | >95% | 1.0 | <15% | 1.0 | 25-50 ft. | 0.7 | 51-75% | 0.7 | >6 - <9 | 0.75 | 20-30 | 0.66 | 0.67 - 0.80 |
| Excellent | >95% | 1.0 | <15% | 1.0 | >50 ft. | 1.0 | 76-100% | 1.0 | >9 - 10 | 1.00 | >30 | 1.0 | > 0.80 |

¹¹Unmanicured (upland) Buffer Width is divided into four categories: Excellent (1.0) = >50 ft, High (0.7) = 25-50 ft, Moderate (0.4) = 10-25 ft, and Low (0.1) = <10 ft.

¹²Estimated Total Vegetative Cover (Percent Range) for upland buffer is the proportion of the ground covered by vegetation within 50 feet of the wetland/upland transition zone. The percent cover is divided into three categories: High and Excellent (1.0) = >95%, Moderate (0.5) = 75 - 95%, and Poor (0.1) = <75%.

¹³The Total Number of Native Plant Species within the unmanicured upland buffer zone is based on two plot locations and a meandering visual survey along the shoreline.

¹⁴(Upland) Buffer Continuity is a measure of the proportion of the water body surrounded by the unmanicured, native upland buffer. This measure is divided into four categories: Excellent (1.0) = 76 - 100%, High (0.7) = 51 - 75%, Medium (0.4) = 26 - 50%, and Low (0.1) = 0 - 25%.

¹⁵Upland buffer exotic species "Percent of Total Coverage" is the percent cover of exotic species within the unmanicured upland buffer, which is divided into three categories: High and Excellent (1.0) = <15%, Moderate (0.5) = 15 - 40%, and Poor (0.1) = >40%.

¹⁶The presence of shoreline erosion is determined by the approximate percentage of the shoreline affected and is divided into the following three categories: 0 - 10%, 11 - 25%, 26 - 100%.

Table 1: Kingsley Lake 2011 and 2016 Habitat Assessment Monitoring Results Black Dog Watershed Management Organization

| Monitoring Year | Submergent Zone | | | | | | | | |
|-----------------|---|--|--|--|---|--|-------------------------|--|--|
| | Approximate Proportion of the Water Body Which is Deep Water Habitat (~ > 20 ft. depth) | Overall Submergent Zone Quality ¹ | Approximate Proportion of Water Body Typically Dominated By Submergent Vegetation (~ 2 - 20 ft. depth) | Native Species | | Mean Coefficient of Conservatism Value | Exotic Species | | |
| | | | | Average Native Plant Density Rating ^{2,3} | Total Number of Native Species ⁵ | | Total Number of Species | Average Exotic Plant Density Rating ^{2,3} | Maximum Exotic Plant Density Rating ⁴ |
| 2011 | 0% | High | 70% | 1.4 (Moderate) | 18 (Excellent) | 5.8 (Moderate) | 0 | 0.0 (Excellent) | 0.0 (Excellent) |
| 2016 | 0% | High | 70% | 1.4 (Moderate) | 20 (Excellent) | 5.7 (Moderate) | 1 | <1.0 (High) | <1.0 (High) |

| Monitoring Year | Emergent Zone | | | | | | |
|-----------------|--|---|---|---|--|-------------------|---|
| | Overall Emergent Zone Quality ⁶ | Approximate Proportion of Emergent Zone (0 - 2 ft. depth) Within The Water Body | Approximate Total Percent Vegetative Cover Within The Entire Emergent Zone ⁷ | Total Number of Native Wetland Plant Species ⁸ | Mean Coefficient of Conservatism Value | Exotic Species | |
| | | | | | | Number of Species | Total Exotic Emergent Percent Coverage ⁹ |
| 2011 | High | 30% | 51-75% (High) | 22 (Excellent) | 3.3 (Moderate) | 4 | 26-50% (High) |
| 2016 | High | 30% | 51-75% (High) | 31 (Excellent) | 3.8 (Moderate) | 4 | 26-50% (High) |

| Monitoring Year | Upland Buffer | | | | | | | | Erosion/Sedimentation | |
|-----------------|---|--|--|--|--|--|-------------------|---|--|--------------------------|
| | Overall Upland Buffer Quality ¹⁰ | Unmanicured Buffer Width ¹¹ | Estimated Total Vegetative Cover (Percent Range) ¹² | Total Number of Native Plant Species ¹³ | Mean Coefficient of Conservatism Value | Buffer Continuity (Percent Surrounding Water Body) ¹⁴ | Exotic Species | | Shoreline Erosion (Percent of Shoreline) ¹⁶ | Sediment Deltas (Yes/No) |
| | | | | | | | Number of Species | Percent of Total Coverage ¹⁵ | | |
| 2011 | High | 25-50 ft. (High) | >95% (High) | 45 (Excellent) | 2.2 (Poor) | 76-100% (Excellent) | 25 | 15-40% (Moderate) | 0-10% | No |
| 2016 | High | 25-50 ft. (High) | >95% (High) | 59 (Excellent) | 2.2 (Poor) | 76-100% (Excellent) | 26 | 15-40% (Moderate) | 0-10% | No |

Table 1: Kingsley Lake 2016 Habitat Assessment Monitoring Results Black Dog Watershed Management Organization

The following changes were made to the 2011 - 2016 monitoring and analysis:

- Monitor one or two water bodies per year. Kingsley Lake in 2011 - Conduct a meandering survey of submergent, emergent, and upland buffer zones rather than monitoring of plot locations. Orchard Lake in 2012, Crystal Lake in 2013, Lac Lavon in 2014, Keller Lake in 2015, Kingsley Lake in 2016 - Conduct a meandering survey of submergent, emergent, and upland buffer zones. In addition, the emergent and upland buffer plot locations were evaluated.
- Changes were made in 2011 through 2016 to the calculations to include floristic quality as part of the assessment. These changes include adding a rating of "High" to the categories to accommodate MPCA ratings for floristic quality. These changes included adding a Rating Code:

| | | |
|------|----------|-------------------|
| Poor | Moderate | High or Excellent |
|------|----------|-------------------|

The following footnotes pertain to 2011 through 2016 data:

¹**Overall Submergent Zone Quality** rating is the average of the rating scores for the following parameters: average exotic plant density, average native plant density, total number of native species, and C-value rating: >0.80 = Excellent, 0.67-0.80 = High, 0.33-0.66 = Moderate, <0.33 = Poor.

| Overall Submergent Zone Quality | Avg. Exotic Plant Density | Exotic Plant Density Rating Score | Avg. Native Plant Density | Avg. Native Plant Density Rating Score | Total Number of Native Species In Submergent Zone | Species Richness Rating Score | Mean Coefficient of Conservatism Value (C-Value) | C-Value Rating (using MPCA values, 2007) | Total Overall Submergent Zone Quality Score |
|---------------------------------|---------------------------|-----------------------------------|---------------------------|--|---|-------------------------------|--|--|---|
| Poor | >2.0 | 0.1 | > 1.75 | 0.1 | <7 | 0.1 | 0 - <3 | 0.10 | < 0.33 |
| Moderate | >1.0 - 2.0 | 0.5 | 1.25 - 1.75 | 0.5 | >7 - <9 | 0.5 | >3 - <6 | 0.50 | 0.33 - 0.66 |
| High | >0 - 1.0 | 0.75 | | | >9 - <14 | 0.75 | >6 - <9 | 0.75 | 0.67 - 0.80 |
| Excellent | 0 | 1.0 | 1.0 to 1.25 | 1.0 | >14 | 1.0 | >9 - 10 | 1.00 | > 0.80 |

²Plant density ratings are a relative measure of the total amount of submergent vegetation covering the submergent zone, with a scale from 1 to 3 utilizing a 6-tined hook; 1 = light density (plant species found on only 1 tine), 2 = moderate density (plant species found on 2 to 4 tines), 3 = heavy density (plant species found on 5 or 6 tines).

³Density data for Kingsley Lake were collected by Barr using a meander survey throughout the lake.

⁴Maximum exotic plant density ratings represent the worst case scenario of curlyleaf pondweed density early in the growing season and/or Eurasian watermilfoil when it is most prolific later in the growing season.

⁵The Total Number of Native Species within the submergent zone for Kingsley Lake was collected by Barr using a meander survey.

The additional category of "High" was added in 2011 through 2016 and values were adjusted to: <7 = Poor, 7-9 = Moderate, 9-14 = High, >14 = Excellent.

⁶**Overall Emergent Zone Quality** is the average of the rating scores for the following parameters within the emergent zone: the total percent coverage, the total number of native wetland plant species, the percent coverage of exotic species, and the C-Value Rating: >0.80 = Excellent, 0.67-0.80 = High, 0.33-0.66 = Moderate, <0.33 = Poor.

| Overall Emergent Zone Quality | Percent Cover | Percent Cover Rating Score | Total Number of Native Wetland Plant Species | Number of Native Wetland Plant Species Rating Score | Percent Cover of Exotics | Percent Cover of Exotics Rating Score | Mean Coefficient of Conservatism Value (C-Value) | C-Value Rating (using MPCA values, 2007) | Overall Emergent Zone Quality Score |
|-------------------------------|-------------------|----------------------------|--|---|--------------------------|---------------------------------------|--|--|-------------------------------------|
| Poor | 0-25% | 0.1 | < or= 5 | 0.1 | 76-100% | 0.1 | 0 - <3 | 0.10 | < 0.33 |
| Moderate | 76-100% or 26-50% | 0.5 | 6 - 10 | 0.33 | 51-75% | 0.33 | >3 - <6 | 0.50 | 0.33 - 0.66 |
| High | 51-75% | 1.0 | 11 - 15 | 0.66 | 26-50% | 0.66 | >6 - <9 | 0.75 | 0.67 - 0.80 |
| Excellent | 51-75% | 1.0 | > 15 | 1.0 | 0-25% | 1.0 | >9 - 10 | 1.00 | > 0.80 |

Table 1: Kingsley Lake 2016 Habitat Assessment Monitoring Results Black Dog Watershed Management Organization

⁷Approximate Total Percent Vegetative Cover Within the Entire Emergent Zone (0-2 ft. depth) is estimated based on the three sampling locations and a visual survey during travels around the water body. Estimates are broken into the following categories: 0-25%=Poor, 26-50%=Moderate, 51-75%=High and Excellent, 76-100%=Moderate.

⁸The Total Number of Native Wetland Plant Species within the emergent zone is based on 3 sampling locations, a meandering visual survey during travels on the water body, and walking along the shoreline: 0-5 = Poor, 6-10 = Moderate, 11-15 = High, and >15 = Excellent.

⁹Total Exotic Emergent Percent Coverage, out of the entire emergent zone area, is estimated based on two plot locations, a meandering visual survey during travels on the water body, and walking along the shoreline. Estimates are broken into four categories: 0-25%=Excellent (1.0), 26-50%=High (0.66), 51-75%=Moderate (0.33), 76-100%=Poor (0.1)

¹⁰**Overall Upland Buffer Quality** is determined based on the average of the six upland buffer quality parameter rating scores: >0.80 = Excellent, 0.67-0.80 = High, 0.33-0.66 = Moderate, <0.33 = Poor.

| Overall Upland Buffer Quality | Percent Cover | Percent Cover Rating Score | Exotics Percent Cover Range | Exotics Percent Cover Rating Score | Buffer Width Range | Buffer Width Rating Score | Buffer Continuity Percent Range | Buffer Continuity Rating Score | Mean Coefficient of Conservatism Value (C-Value) | C-Value Rating (using MPCA values, 2007) | Number of Native Species | Number of Native Species Rating Score | Overall Upland Buffer Quality Score |
|--------------------------------------|----------------------|-----------------------------------|------------------------------------|---|---------------------------|----------------------------------|--|---------------------------------------|---|---|---------------------------------|--|--|
| Poor | <75% | 0.1 | >40% | 0.1 | <10 ft. | 0.1 | 0-25% | 0.1 | 0 - <3 | 0.10 | <5 | 0.1 | < 0.33 |
| Moderate | 75-95% | 0.5 | 15-40% | 0.5 | 10-25 ft. | 0.4 | 25-50% | 0.4 | >3 - <6 | 0.50 | 5-20 | 0.33 | 0.33 - 0.66 |
| High | >95% | 1.0 | <15% | 1.0 | 25-50 ft. | 0.7 | 51-75% | 0.7 | >6 - <9 | 0.75 | 20-30 | 0.66 | 0.67 - 0.80 |
| Excellent | >95% | 1.0 | <15% | 1.0 | >50 ft. | 1.0 | 76-100% | 1.0 | >9 - 10 | 1.00 | >30 | 1.0 | > 0.80 |

¹¹Unmanicured (upland) Buffer Width is divided into four categories: Excellent (1.0) = >50 ft, High (0.7) = 25-50 ft, Moderate (0.4) = 10-25 ft, and Low (0.1) = <10 ft.

¹²Estimated Total Vegetative Cover (Percent Range) for upland buffer is the proportion of the ground covered by vegetation within 50 feet of the wetland/upland transition zone. The percent cover is divided into three categories: High and Excellent (1.0) = >95%, Moderate (0.5) = 75 - 95%, and Poor (0.1) = <75%.

¹³The Total Number of Native Plant Species within the unmanicured upland buffer zone is based on two plot locations and a meandering visual survey along the shoreline.

¹⁴(Upland) Buffer Continuity is a measure of the proportion of the water body surrounded by the unmanicured, native upland buffer. This measure is divided into four categories: Excellent (1.0) = 76 - 100%, High (0.7) = 51 - 75%, Medium (0.4) = 26 - 50%, and Low (0.1) = 0 - 25%.

¹⁵Upland buffer exotic species "Percent of Total Coverage" is the percent cover of exotic species within the unmanicured upland buffer, which is divided into three categories: High and Excellent (1.0) = <15%, Moderate (0.5) = 15 - 40%, and Poor (0.1) = >40%.

¹⁶The presence of shoreline erosion is determined by the approximate percentage of the shoreline affected and is divided into the following three categories: 0 - 10%, 11 - 25%, 26 - 100%.

Appendix D

2003–2018 Recommended and Completed Management Actions

**Table D-1: 2009 Recommended and Completed Management Actions
Black Dog Watershed Management Organization Habitat Monitoring**

| Strategic Water Body | Problem Identified | Recommendation | Proposed Action | Benefits | Implementation Period | Completed 2004-2009 Actions Which May Improve Wildlife Habitat and/or Water Quality |
|----------------------|---|--|---|--|----------------------------|--|
| Crystal | Unmanicured, native vegetation in adjacent upland and emergent zone is narrow and not continuous, limiting wildlife benefits. | 1. Increase width and continuity of native upland buffer and emergent zone. | Conduct an educational workshop and lakescaping demonstration project. Assist lakeshore owners with funding and obtaining any needed MnDNR permits for potential upland buffer and emergent zone enhancements. | Inform/show lakeshore property owners how a native upland buffer and native emergent zone can improve functions and values of the lake and improve aesthetics. | Spring - Fall | 2009: Operation of the ferric chloride treatment system halted due to low water levels. The City of Burnsville harvested curlyleaf pondweed. In late 2009, the City of Burnsville treated 14 acres of buckthorn within Crystal West Park. In 2009 and 2008, garlic mustard within the upland buffer was removed/pulled. 2004-2008: The BDV/MO resumed and continued operation of the ferric chloride treatment system. The City of Burnsville: 1) excavated/enhanced four stormwater treatment ponds (including West Buck Hill Park), which reduced the phosphorus loading into the lake, and 2) conducted annual harvesting of Eurasian watermilfoil and curlyleaf pondweed. The City of Lakeville excavated/enhanced the Bluebill stormwater treatment pond. |
| | Purple loosestrife is present. | 2. Continue to control and manage purple loosestrife. | Control and manage. For large stands of purple loosestrife, contact the MnDNR to request a release of purple loosestrife-controlling beetles. For a few small colonies of purple loosestrife, hand pull or dig the plants out before they go to seed. | Increase/maintain wildlife habitat. | Spring - Fall | |
| | Curlyleaf pondweed is present. | 3. Control curlyleaf pondweed | Control by harvesting or chemical treatment. | Maintain wildlife habitat. | Late Spring | |
| | Eurasian watermilfoil is present. | 4. Control Eurasian watermilfoil. | Control by chemical treatment. | Maintain wildlife habitat. | Summer | |
| Keller Lake | Unmanicured, native vegetation in adjacent upland is narrow and not continuous, limiting wildlife benefits. | 1. Increase width and continuity of native upland buffer. | Conduct an educational workshop and lakescaping demonstration project. Assist lakeshore owners with funding of potential upland buffer enhancements. | Inform/show lakeshore property owners how a native upland buffer can improve functions and values of the lake and improve aesthetics. | Spring - Fall | In 2010 the City of Apple Valley may construct Whitney Pond for stormwater treatment within the Keller Lake watershed. 2009: Due to low water levels, operation of the ferric chloride treatment system halted and no harvesting of curlyleaf pondweed was conducted. The City of Burnsville stabilized approximately one hundred feet of shoreline on the southeast edge of the lake. Logs were interlaced and secured along the shoreline and red-osier dogwood live stakes were installed along the eroding banks. 2004 - 2008: The Cities of Apple Valley and Burnsville partnered to conduct annual harvesting of curlyleaf pondweed. 2005: The City of Apple Valley excavated and enhanced Redwood Pond, which will decrease phosphorus loading into Keller Lake. Also, in 2010 the City of Apple Valley may construct Whitney Pond for stormwater treatment within the Keller Lake watershed. |
| | Purple loosestrife is present. | 2. Continue to control and manage purple loosestrife. | Control and manage. For large stands of purple loosestrife, contact the MnDNR to request a release of purple loosestrife-controlling beetles. For a few small colonies of purple loosestrife, hand pull or dig the plants out before they go to seed. | Increase/maintain wildlife habitat. | Spring - Fall | |
| | Curlyleaf pondweed dominates the lake in late spring-early summer. | 3. Continue control of curlyleaf pondweed. | Control as recommended by the MnDNR. Since the MnDNR designates Keller Lake as a "Natural Environment", a special permit is needed to chemically treat the lake. | Maintain wildlife habitat. | Summer | |
| | Eurasian watermilfoil is present. | 4. Control Eurasian watermilfoil. | Control as recommended by the MnDNR. Since the MnDNR designates Keller Lake as a "Natural Environment", a special permit is needed to chemically treat the lake. | Maintain wildlife habitat. | Summer | |
| Kingsley Lake | Curlyleaf pondweed is present. | 1. Conduct a detailed late spring macrophyte survey to ascertain densities and extent of coverage. | Consider control measures, dependent on results of an detailed early growing season survey. | Maintain wildlife habitat. | Late Spring | 2005 - 2008: Annually, the City of Lakeville and members of the Kingsley Lake Homeowner's Association removed purple loosestrife plants and common buckthorn from portions of the lake and the upland buffer surrounding the lake. On March 6, 2008, soil sediment samples were collected on Kingsley Lake by Blue Water Science (BWS) and the City of Lakeville. Based on the results of the soil analysis, the BWS report stated that "curlyleaf pondweed is not expected to produce heavy growth conditions (where plants top out in a solid canopy) in Kingsley Lake." However, since curlyleaf pondweed may typically die-off prior to the early-June habitat assessment, the peak density and percent total coverage of curlyleaf pondweed is uncertain. To date, it is unclear if curlyleaf pondweed densities and percent coverage have been relatively consistent or increasing within the lake over the last few years. In 2008, a Kingsley lakeshore resident, inspired by the Blue Thumb program, commenced shoreline stabilization utilizing native plants. |
| | Common buckthorn dominates portions of the upland buffer. | 2. Conduct an evaluation of common buckthorn, followed by removal. | Remove buckthorn. Volunteer groups and contractors can effectively remove buckthorn by pulling, cutting, and treating stumps with herbicide. | Increase wildlife habitat. | Open | |
| | Purple loosestrife is present. | 3. Continue to control and manage purple loosestrife. | Control and manage. For large stands of purple loosestrife, contact the MnDNR to request a release of purple loosestrife-controlling beetles. For a few small colonies of purple loosestrife, hand pull or dig the plants out before they go to seed. | Increase/maintain wildlife habitat. | Spring - Fall | |
| | Hybrid cattail and reed canary grass are present. | 4. Control hybrid cattail and reed canary grass. | Control hybrid cattail and reed canary grass now before colonies become more abundant. The herbicide Rodeo™ can be used to effectively control both invasive emergent species. | Increase/maintain wildlife habitat. | Spring-Summer | |
| Lac Lavon | Eurasian watermilfoil dominates portions of the lake. | 1. Continue to manage Eurasian watermilfoil. | Control by chemical treatment as recommended by MnDNR. | Increase/maintain wildlife habitat and water quality | Spring-Summer | 2006: The Cities of Burnsville and Apple Valley and the lake homeowners partnered to fund a fluridone treatment for control of Eurasian watermilfoil. The treatment is expected to provide control of Eurasian watermilfoil for three years, while allowing native plant species to rebound. The cities have continued to inform the MnDNR of the ongoing treatments and the MnDNR proposes to continue aquatic plant surveys to study the effects of whole-lake fluridone treatments. However, no MnDNR macrophyte survey was conducted in 2008. |
| | Curlyleaf pondweed is present. | 2. Monitor presence of curlyleaf pondweed. | Control if increased occurrence and subsequent midsummer die off threatens water quality) | Identify the problem before it becomes difficult to treat. | Spring | |
| | Unmanicured, native vegetation in adjacent upland is narrow and not continuous, limiting wildlife benefits. | 3. Increase width/creation of native upland buffer. | Conduct an educational workshop and lakescaping demonstration project. Assist lakeshore owners with funding of potential upland buffer enhancements. | Inform/show lakeshore property owners of how a native upland buffer can improve functions and values of the lake and improve aesthetics. | Spring - Fall | |
| Orchard Lake | Curlyleaf pondweed dominates the lake in late spring-early summer. | 1. Continue curlyleaf pondweed control measures. | Restore sustainable native communities | Increase wildlife habitat. | Spring - Fall | 2009: The City of Lakeville conducted herbicide treatment for curlyleaf pondweed within the northeast bay (~20 acres). The herbicide treatment resulted in lake-wide control of curlyleaf pondweed. 2004-2008: The City of Lakeville provided lakeshore owners with shoreline restoration information. However, to date, no plans have been made for potential future shoreline restoration projects. Annually, the City of Lakeville harvested approximately 70 acres of curlyleaf pondweed. 2007: A small area of lakeshore, near the boat launch, was restored using native plants. |
| | Unmanicured, native vegetation in adjacent upland is narrow and not continuous, limiting wildlife benefits. | 2. Increase width and continuity of native upland buffer. | Control and manage | Increase/maintain wildlife habitat and water quality. | Late Spring - Early summer | |
| | | | Conduct an educational workshop and lakescaping demonstration project. Assist lakeshore owners with funding of potential upland buffer enhancements. | Inform/show lakeshore property owners how a native upland buffer can improve functions and values of the lake and improve aesthetics. | Spring - Fall | |
| | | | Restore sustainable native communities | Increase wildlife habitat. | Spring - Fall | |
| Sunset Pond | Purple loosestrife is present. | 3. Conduct a detailed evaluation of purple loosestrife, followed by removal/control. | Control and manage by hand-pulling if only a few plants are present or introduce beetles if numerous plants are present. | Increase/maintain wildlife habitat. | Spring - Summer | In 2009, as in past years, the City of Burnsville actively managed the restored native buffer adjacent to the pond, the surrounding prairie restoration area, and portions of the emergent zone. Specifically, in 2007 through 2009 the City of Burnsville conducted spot spraying of invasive vegetation, such as reed canary grass, thistle, and purple loosestrife. A prescription burn, inter-seeding of prairie species, and buckthorn removal were conducted in 2008 to increase the plant diversity in the upland area. |
| | Extensive algal bloom | 1. Reduce phosphorus loading into the pond. | Construct/install: catch basin sumps, prefabricated treatment devices (e.g. Stormceptor), infiltration facilities within the watershed, or other more conventional methods. Conduct more frequent street sweepings. | Improve wildlife habitat, fishery habitat, and aesthetics/recreation. | Open | |
| | Maintained turf grass remains within portions of the upland buffer. | 2. Enhance/maintain upland buffer. | Continue restoring sustainable native communities | Improve wildlife habitat and water quality. | Spring - Fall | |
| | Exotic species are dominant in emergent zone, and include narrow-leaf cattail, hybrid cattail, and reed canary grass. | 3. Manage exotic species within emergent zone. | Selective herbicide treatments to reduce presence of exotic species | Allow for the establishment of more diverse native species that provide better wildlife values. | Spring - Fall | |
| | Presence of curlyleaf pondweed observed in 2003 and 2005 through 2008. | 4. Conduct a late spring macrophyte survey to ascertain densities and extent of coverage. | consider control measures dependent on the results of an early growing season survey. | Maintain wildlife habitat. Reduce down-stream phosphorus loading. | Late Spring | |
| | The southern portion of the pond is shallow (1 to 3 feet deep). | 5. Create a "navigation channel". | Excavate and remove sediment. | Improve wildlife habitat, fishery habitat, and aesthetics/recreation. | Winter | |

**Table 2: 2011 Recommended and Completed Management Actions for Kingsley Lake
Black Dog Watershed Management Organization Habitat Monitoring**

| Problem Identified | Recommendation | Proposed Action | Benefits | Implementation Period | Completed 2004-2009 Actions Which May Improve Wildlife Habitat and/or Water Quality |
|---|---|--|---|-----------------------|--|
| Curlyleaf pondweed is present in some years. | Conduct a detailed late spring macrophyte survey to ascertain densities and extent of coverage. | Consider control measures, dependent on results of a detailed early growing season survey. | Maintain wildlife habitat. | Late Spring | <p>2005 - 2008: Annually, the City of Lakeville and members of the Kingsley Lake Homeowner's Association removed purple loosestrife plants and common buckthorn from portions of the lake and the upland buffer surrounding the lake. Purple loosestrife beetles were released by the MnDNR prior to 2002. Follow up monitoring by the MnDNR indicates that beetles are present at a population that the MnDNR feels is appropriate for biological control. On March 6, 2008, soil sediment samples were collected on Kingsley Lake by Blue Water Science (BWS) and the City of Lakeville. Based on the results of the soil analysis, the BWS report stated that "curlyleaf pondweed is not expected to produce heavy growth conditions (where plants top out in a solid canopy) in Kingsley Lake." However, since curlyleaf pondweed may typically die-off prior to the early-June habitat assessment, the peak density and percent total coverage of curlyleaf pondweed is uncertain. To date, it is unclear if curlyleaf pondweed densities and percent coverage have been relatively consistent or increasing within the lake over the last few years. In 2008, a Kingsley Lake lakeshore resident, inspired by the Blue Thumb program, commenced shoreline stabilization utilizing native plants.</p> |
| Common buckthorn dominates portions of the upland buffer. | Conduct an evaluation of common buckthorn, followed by removal. | Remove buckthorn. Volunteer groups and contractors can effectively remove buckthorn by pulling, cutting, and treating stumps with herbicide. | Increase wildlife habitat. | Open | |
| Purple loosestrife is present. | Continue to control and manage purple loosestrife. | Control and manage. For a few small colonies of purple loosestrife, hand pull or dig the plants out before they go to seed. Continue to request monitoring from the MnDNR to assure beetles are present and at appropriate populations for biological control. | Increase/maintain wildlife habitat. | Spring - Fall | |
| Hybrid cattail and reed canary grass are present. | Control hybrid cattail and reed canary grass. | Control hybrid cattail and reed canary grass now before colonies become more abundant. The herbicide Rodeo™ can be used to effectively control both invasive emergent species. | Increase/maintain wildlife habitat. | Spring-Summer | |
| Stormwater drainage from impervious surfaces is directed into the lake. | Redirect stormwater for infiltration prior to discharge. | Install a rainwater garden or other suitable method for infiltration. | Improve water quality | Open | |
| Bare soil on steep slope could cause erosion and sedimentation into lake. | Vegetate hillslope. | Plant vegetation suited for steep slopes along hillside to prevent erosion. | Improve water quality | Open | |
| Upland buffer areas lacking naturalized vegetation. | Improve the shoreline with a naturalized upland buffer. | Rather than manicured turf grass, gravel, and managed plantings with bare soil, the shoreline could be vegetated with native grasses and wildflowers. A landscape architect could create inviting spaces and views for restaurant customers to enjoy. | Increase wildlife habitat and Improve water quality | Open | |
| Emergent and upland buffer areas contain non-native invasive vegetation. | Replace non-native invasive vegetation with native vegetation. | Treat non-native invasive vegetation and then seed with an appropriate BWSR seed mix. | Increase/maintain wildlife habitat. | Spring-Summer | |

**Table 2: 2012 Recommended and Completed Management Actions for Orchard Lake
Black Dog Watershed Management Organization Habitat Monitoring**

| Problem Identified | Recommendation | Proposed Action | Benefits | Implementation Period | Completed 2004-2012 Actions Which May Improve Wildlife Habitat and/or Water Quality |
|---|--|--|---|----------------------------|---|
| Curlyleaf pondweed dominates the lake in late spring-early summer. | Continue curlyleaf pondweed control measures. | Continue to control and manage. See Figure 3 for locations of curlyleaf pondweed. | Increase wildlife habitat, improve water quality, vegetative diversity, aesthetics, and recreation. | Late Spring - Early summer | <p>1999 through 2012: The City of Lakeville conducts aquatic vegetation monitoring twice/year.</p> <p>2009 through 2012: The City of Lakeville conducted annual herbicide treatment for curlyleaf pondweed.</p> <p>2004 through 2008: Annually, the City of Lakeville harvested approximately 70 acres of curlyleaf pondweed.</p> <p>2010: Adjacent to the southwest end of the lake, an aeration system was installed in Orchard Pond to precipitate out phosphorus and improve water quality flowing into Orchard Lake.</p> <p>2004 through 2012: The City of Lakeville annually provides lakeshore owners with shoreline restoration information and encourages homeowners to take advantage of the Blue Thumb restoration program. One shoreline resident started a restoration project in 2012.</p> <p>2007: A small area of lakeshore, near the boat launch, was restored using native plants.</p> <p>2002: Purple loosestrife beetles were released by the MNDNR. Follow up monitoring indicates that beetles are present at a population that the MNDNR feels is appropriate for biological control of purple loosestrife plants.</p> |
| Upland buffer areas lacking naturalized vegetation within publicly owned properties. | Increase width and continuity of native upland buffer. | <p>To expand on the shoreline restoration that was done near the boat launch in 2007, the adjacent upland buffer could also be restored to naturalized native vegetation and not mowed (Potential Restoration Area #1 as shown in Appendix A and Figure 5).</p> <p>In the Wayside Park Area, non-native invasive vegetation including common buckthorn, vetch, spotted knapweed, and cattails could be removed and replaced with native vegetation. The naturalized upland buffer could be widened (Potential Restoration Area #2 as shown in Appendix A and Figure 5).</p> <p>At the beach area, there is a timber wall which is currently being used for fishing. A shoreline restoration could be done in this area (Potential Restoration Area #3 as shown in Appendix A and Figure 5).</p> <p>On the northwest side of the lake, one property owned by the City of Lakeville (adjacent to residential shoreline properties) could be restored to naturalized vegetation and provide an example for adjacent residential landowners for shoreline and upland buffer restoration (Potential Restoration Area #4 as shown in Appendix A and Figure 5).</p> | Increase wildlife habitat. Improve water quality. Improve vegetative diversity and aesthetics. | Spring - Fall | |
| Upland buffer areas lacking naturalized vegetation. Most of the residential properties have turf grass up the the lakeshore edge. | Increase width and continuity of native upland buffer. | Restore sustainable native communities. Rather than manicured turf grass, sand, and bare soil, the shoreline could be vegetated with native grasses and wildflowers. A native upland buffer can improve functions and values of the lake and improve aesthetics (Potential Restoration Area #5 as shown in Appendix A and Figure 5). | Increase wildlife habitat. Improve water quality. Improve vegetative diversity and aesthetics. | Spring - Fall | |
| Purple loosestrife is present. | Continue to control and manage purple loosestrife. | Control and manage. For a few small colonies of purple loosestrife, hand pull or dig the plants out before they go to seed. Continue to request monitoring from the MNDNR to assure beetles are present and at appropriate populations for biological control (See Figures 3 and 5 for location of purple loosestrife). | Increase wildlife habitat. Improve vegetative diversity. | Spring - Fall | |

Table 2: 2013 Recommended and Completed Management Actions for Crystal Lake
Black Dog Watershed Management Organization Habitat Monitoring

| Problem Identified | Recommendation | Proposed Action | Benefits | Implementation Period | Completed Actions Which May Improve Wildlife Habitat and/or Water Quality |
|---|--|--|---|----------------------------|---|
| Curlyleaf pondweed dominates the lake in late spring-early summer. | Continue curlyleaf pondweed control measures. | Continue to control and manage. See Blue Water Science report for locations of curlyleaf pondweed. | Increase wildlife habitat, improve water quality, vegetative diversity, aesthetics, and recreation. | Late Spring - Early summer | <p>1999 through 2013: The City of Burnsville conducts aquatic vegetation monitoring twice/year.</p> <p>2003 through 2013: The City of Burnsville conducted annual harvesting of curlyleaf pondweed.</p> <p>2004-2008:</p> <ul style="list-style-type: none">-The BDWMO operated the ferric chloride treatment system.-The City of Burnsville: 1) excavated/enhanced four stormwater treatment ponds (including West Buck Hill Park), which reduced the phosphorus loading into the lake, and 2) conducted annual harvesting of Eurasian watermilfoil and curlyleaf pondweed.-The City of Lakeville excavated/enhanced the Bluebill stormwater treatment pond. <p>In 2009 and 2008, garlic mustard within the upland buffer was removed/pulled.</p> <p>In late 2009, the City of Burnsville treated 14 acres of buckthorn within Crystal West Park.</p> |
| Upland buffer areas lacking naturalized vegetation within publicly owned properties. | Increase width and continuity of native upland buffer. | The width and density of naturalized shoreline buffer at the location of Emergent Plot #1 near the swimming area has improved significantly since 2009. The adjacent upland buffer could also be restored to naturalized native vegetation and not mowed (Potential Restoration Areas #1 through 4 as shown in Figure 4 and photos). | Increase wildlife habitat. Improve water quality. Improve vegetative diversity and aesthetics. | Spring - Fall | |
| Upland buffer areas lacking naturalized vegetation. Most of the residential properties have turf grass up the the lakeshore edge. | Increase width and continuity of native upland buffer. | Restore sustainable native communities. Rather than manicured turf grass, sand, and bare soil, the shoreline could be vegetated with native grasses and wildflowers. A native upland buffer can improve functions and values of the lake and improve aesthetics (Potential Restoration Area #5 as shown in Figure 4 and photos). | Increase wildlife habitat. Improve water quality. Improve vegetative diversity and aesthetics. | Spring - Fall | |
| Purple loosestrife is present. | Continue to control and manage purple loosestrife. | Continue to control. For a few small colonies of purple loosestrife, hand pull or dig the plants out before they go to seed. | Increase wildlife habitat. Improve vegetative diversity. | Spring - Fall | |
| Eurasian watermilfoil is present. | Control Eurasian watermilfoil. | Control by chemical treatment. | Maintain wildlife habitat. | Summer | |

**Table 2: 2014 Recommended and Completed Management Actions for Lac Lavon
Black Dog Watershed Management Organization Habitat Monitoring**

| Problem Identified | Recommendation | Proposed Action | Benefits | Implementation Period | Completed Actions Which May Improve Wildlife Habitat and/or Water Quality |
|---|---|--|---|----------------------------|--|
| Curlyleaf pondweed dominates the lake in late spring-early summer. | Continue curlyleaf pondweed control measures. | Continue to control and manage. See Macrophyte Survey Results for locations of curlyleaf pondweed. | Increase wildlife habitat, improve water quality, vegetative diversity, aesthetics, and recreation. | Late Spring - Early summer | Aquatic plant surveys were conducted by Barr in 2013 and 2014. |
| Eurasian watermilfoil is present. | Control Eurasian watermilfoil. | Control by chemical treatment. See Macrophyte Survey Results for locations of Eurasian watermilfoil | Maintain wildlife habitat. | Summer | In 2006, the cities of Burnsville and Apple Valley and the lake homeowners partnered to fund a fluridone treatment for control of Eurasian watermilfoil. Aquatic plant surveys were conducted by Barr in 2013 and 2014. |
| Purple loosestrife is present. | Continue to control and manage purple loosestrife. | Continue to control. For a few small colonies of purple loosestrife, hand pull or dig the plants out before they go to seed. See Macrophyte Survey Results for locations of purple loosestrife | Increase wildlife habitat. Improve vegetative diversity. | Spring - Fall | Purple loosestrife removal on shallow island areas was completed by the cities of Apple Valley and Burnsville in 2011. |
| Upland buffer areas lacking naturalized vegetation within publicly owned properties. | Increase width and continuity of native upland buffer. | Expand native prairie planting to include area to the east, which is dominated by knapweed. This could become a tall grass prairie. Potential Restoration Area #1 | Increase wildlife habitat. Improve water quality. Improve vegetative diversity and aesthetics. | Spring - Fall | In 2013, the city of Burnsville installed a native prairie planting converting a sand beach and turf grass to prairie and wetland vegetation. |
| Upland buffer areas in city parks contain non-native invasive vegetation such as buckthorn, Siberian elm, leafy spurge, and spotted kanpweed. | Continue to control and manage non-native invasive vegetation | Continue to control and manage non-native invasive vegetation Potential Restoration Area #2 | Increase wildlife habitat. Improve vegetative diversity and aesthetics | Spring - Fall | Some invasive species control for Canada thistle and knapweed was conducted on the new native planting area in 2014. In 2010, the city of Apple Valley released about 150 spotted knapweed seedhead boring weevils in Lac Lavon Park in Apple Valley. Continued management of the vegetation communities and shoreline restoration activities will help to maintain and improve wildlife habitat, vegetation diversity, aesthetics, and recreation |
| Impervious surfaces and turf grass in the Apple Valley park near the fishing pier can collect pollutants in stormwater and flow directly into the lake, decreasing water quality. | Increase areas of naturalized vegetation to slow down and pretreat stormwater prior to entering the lake. | Strategically create buffer strips with naturalized vegetation adjacent to impervious surfaces to slow down and pretreat stormwater prior to entering the lake. Potential Restoration Area #3 | Improve water quality | Spring - Fall | |
| Upland buffer areas lacking naturalized vegetation. Most of the residential properties have turf grass or sand up to the lakeshore edge. | Increase width and continuity of native upland buffer. | Restore sustainable native communities. Rather than manicured turf grass, sand, and bare soil, the shoreline could be vegetated with native grasses and wildflowers. A native upland buffer can improve functions and values of the lake and improve aesthetics. Potential Restoration Area #4 | Increase wildlife habitat. Improve water quality. Improve vegetative diversity and aesthetics. | Spring - Fall | One raingarden was installed in the backyard of a shoreline property owner on Highview Drive in Apple Valley through the Blue Thumb program. The establishment of shoreline restoration projects (especially contiguous) on residential properties in the future will help balance out the differences in upland buffer habitat between city owned property and residential property. |

**Table 2: 2015 Recommended and Completed Management Actions for Keller Lake
Black Dog Watershed Management Organization Habitat Monitoring**

| Problem Identified | Recommendation | Proposed Action | Benefits | Implementation Period | Completed Actions Which May Improve Wildlife Habitat and/or Water Quality |
|---|---|---|---|------------------------------|--|
| Curlyleaf pondweed dominates the lake in late spring-early summer. | Continue curlyleaf pondweed control measures. | Continue to control and manage. See Appendix A Aquatic Plant Survey for locations of curlyleaf pondweed. | Increase wildlife habitat, improve water quality, vegetative diversity, aesthetics, and recreation. | Late Spring - Early summer | Aquatic plant surveys have been conducted by Blue Water Science 1998-2015. Iron dosing occurred from 1999 through 2008. Mechanical harvesting is conducted each year since 2004. |
| Eurasian watermilfoil is present. | Control Eurasian watermilfoil. | Continue to monitor. Control as recommended by the MnDNR. Since the MnDNR designates Keller Lake as a "Natural Environment Lake", chemical treatment is not allowed. | Maintain wildlife habitat. | Summer | Aquatic plant surveys have been conducted by Blue Water Science 1998-2015. |
| The inlet coming from the stormwater pond at the south end of Keller Lake is surrounded by bare soil or sparse vegetation. | Re-vegetated bare areas to prevent soil erosion into Keller Lake. | Seed or plant bare areas with native vegetation. Potential Restoration Area #1 | Improve water quality and vegetative diversity. | Spring or Fall | |
| Shoreline fishing traffic is causing bare soil areas along the shoreline. | Re-vegetated bare areas to prevent soil erosion into Keller Lake. | Create designated stone walkways for fishing access. Potential Restoration Area #2 | Improve water quality, vegetative diversity, and aesthetics. | Spring - Fall | |
| The southern public park is littered with trash and other dumped items especially near the shoreline. | Clean up the litter. | Organize a neighborhood clean-up project to pick up trash and other dumped items along the south shoreline of the lake. Potential Restoration Area #3 | Improve aesthetics. Potentially prevent harm to wildlife. Prevent migration of trash into lake. | Spring - Fall | |
| Upland buffer areas in city parks contain non-native invasive vegetation such as buckthorn and garlic mustard. | Continue to control and manage non-native invasive vegetation | Continue to control and manage non-native invasive vegetation. Pull garlic mustard within the City of Burnsville property at the north end of the lake. Continue to remove and treat new growth of buckthorn in city parks. Potential Restoration Area #4 | Increase wildlife habitat. Improve vegetative diversity and aesthetics | Spring - Fall | Buckthorn appears to have been previously removed in the park along the southern shoreline. |
| Upland buffer areas lacking naturalized vegetation. Some of the residential properties have narrow buffers with turf grass close to the lakeshore edge. | Increase width and continuity of native upland buffer. | Restore sustainable native communities. Manicured turf grass near the shoreline could be vegetated with native grasses and wildflowers. A native upland buffer can improve functions and values of the lake and improve aesthetics. Potential Restoration Area #5 | Increase wildlife habitat. Improve water quality. Improve vegetative diversity and aesthetics. | Spring - Fall | Most residential properties allow a narrow width of naturalized vegetation to prevent soil erosion, however a wider buffer of native vegetation could help improve wildlife habitat, vegetative diversity, and aesthetics. |

**Table 2: 2016 Recommended and Completed Management Actions for Kingsley Lake
Black Dog Watershed Management Organization Habitat Monitoring**

| Problem Identified | Recommendation | Proposed Action | Benefits | Implementation Period | Completed Actions Which May Improve Wildlife Habitat and/or Water Quality |
|---|--|--|---|----------------------------|--|
| Curlyleaf pondweed is present in some years. | Continue to monitor | Consider control measures, if densities and locations increase to an extent of concern. See Appendix A Aquatic Plant Survey for locations of curlyleaf pondweed. | Increase wildlife habitat, improve water quality, vegetative diversity, aesthetics, and recreation. | Late Spring - Early summer | On March 6, 2008, soil sediment samples were collected on Kingsley Lake by Blue Water Science (BWS) and the City of Lakeville. Based on the results of the soil analysis, the BWS report stated that “curlyleaf pondweed is not expected to produce heavy growth conditions (where plants top out in a solid canopy) in Kingsley Lake.” |
| Common buckthorn dominates portions of the upland buffer. | Conduct an evaluation of common buckthorn, followed by removal. | Remove buckthorn. Volunteer groups and contractors can effectively remove buckthorn by pulling, cutting, and treating stumps with herbicide. See Figure 4, Potential Restoration Area #1 | Increase wildlife habitat. Improve vegetative diversity and aesthetics | Spring - Fall | From 2005-2008, the City of Lakeville and members of the Kingsley Lake Association removed common buckthorn from portions of the lake and the upland buffer surrounding the lake. |
| Purple loosestrife is present. | Continue to control and manage purple loosestrife. | Control and manage. For a few small colonies of purple loosestrife, hand pull or dig the plants out before they go to seed. See Figure 4 for purple loosestrife locations. | Increase/maintain wildlife habitat. | Spring - Fall | From 2005-2008, the City of Lakeville and members of the Kingsley Lake Association removed purple loosestrife plants from portions of the lake and the upland buffer surrounding the lake. Purple loosestrife beetles were released by the MnDNR prior to 2002. Follow up monitoring by the MnDNR indicates that beetles are present at a population that the MnDNR feels is appropriate for biological control. |
| Stormwater drainage from impervious surfaces is directed into the lake. | Redirect stormwater for infiltration prior to discharge. | Install a rainwater garden, pervious pavement, or other suitable method for infiltration. See Figure 4, Potential Restoration Area #2. | Improve water quality | Open | |
| Bare soil on steep slope could cause erosion and sedimentation into lake. | Re-vegetate bare areas to prevent soil erosion into Kingsley Lake. | Plant vegetation suited for steep slopes along hillside to prevent erosion. See Figure 4, Potential Restoration Area #3 | Improve water quality | Spring - Fall | |
| Upland buffer areas lacking naturalized vegetation. | Increase width and continuity of native upland buffer. | Rather than manicured turf grass, gravel, and managed plantings with bare soil, the shoreline could be vegetated with native grasses and wildflowers. See Figure 4, Potential Restoration Areas #4 through 7. See Appendix G for examples of improvements. See also island shoreline areas becoming bare from YMCA camper overuse (Figure 4, Potential Restoration Areas 10 and 11). | Improve water quality, increase wildlife habitat. Improve vegetative diversity and aesthetics. | Spring - Fall | In 2008, a Kingsley Lake lakeshore resident, inspired by the Blue Thumb program, commenced shoreline stabilization utilizing native plants. |
| Emergent zone and upland buffer areas contain non-native invasive vegetation. | Continue to control and manage non-native invasive vegetation, including, but not limited to reed canary grass, hybrid cattail, and yellow iris. | Treat non-native invasive vegetation and then seed with an appropriate BWSR seed mix. See Figure 4, Potential Restoration Areas #8 and 9. Remove yellow iris (See Appendix A for locations of yellow iris). The MN DNR may require a permit for cattail treatment and yellow iris removal if below the OHW. Dense reed canary grass is located at Plot 2b as shown of Figure 3 . Dense hybrid cattail is located at Plot 1b as shown on Figure 3 . | Increase/maintain wildlife habitat. | Spring-Summer | |

**Table 2: 2017 Recommended and Completed Management Actions for Orchard Lake
Black Dog Watershed Management Organization Habitat Monitoring**

| Problem Identified | Recommendation | Proposed Action | Benefits | Implementation Period | Completed Actions Which May Improve Wildlife Habitat and/or Water Quality |
|--|---|--|---|----------------------------|--|
| Curlyleaf pondweed is common in early spring | Continue to monitor, control, and manage. | Continue to treat curlyleaf pondweed where growth is predicted to be heavy. See Appendix A Aquatic Plant Survey for more details. | Increase wildlife habitat, improve water quality, vegetative diversity, aesthetics, and recreation. | Late Spring - Early summer | From 1999-2017, the City of Lakeville contracts Blue Water Science to conduct aquatic plant surveys twice per year. Curlyleaf pondweed was harvested annually from 2004-2009. Herbicide treatments were conducted annually from 2009-2012 and 2015-2017. |
| Purple loosestrife is present. | Continue to control and manage purple loosestrife. | Control and manage. For a few small colonies of purple loosestrife, hand pull or dig the plants out before they go to seed. See Figure 4 for purple loosestrife locations. | Increase/maintain wildlife habitat. | Spring - Fall | Purple loosestrife beetles were released by the MnDNR prior to 2002. Follow up monitoring by the MnDNR indicates that beetles are present at a population that the MnDNR feels is appropriate for biological control. |
| Stormwater drainage from impervious surfaces is directed into the lake. | Redirect stormwater for infiltration prior to discharge. | Install a rainwater garden, pervious pavement, or other suitable method for infiltration and establish a naturalized upland buffer. See Figure 4 and Site Photos, Potential Restoration Area #6. | Improve water quality | Open | Two raingardens were completed on 175th St W. In 2010, adjacent to the southwest end of the lake, an aeration system was installed in Orchard Pond to precipitate out phosphorus and improve water quality flowing into Orchard Lake. |
| Bare soil along shoreline could cause erosion and sedimentation into lake. | Re-vegetate bare areas to prevent soil erosion into Orchard Lake. | Improve soil and plant vegetation along shoreline to prevent erosion. Establish a canoe and kayak access at Wayside Park. See Figure 4 and Site Photos, Potential Restoration Area #4 and #5. | Improve water quality | Spring - Fall | The City of Lakeville removed a dilapidated timber wall and attempted a shoreline restoration south of the beach, however, the soil was too poor for the plantings to become established. North of the beach, a concrete wall was built to prevent shoreline erosion. |
| Upland buffer areas lacking naturalized vegetation. | Increase width and continuity of native upland buffer. | Rather than manicured turf grass the shoreline could be vegetated with native grasses and wildflowers. See Figure 4 and Site Photos, Potential Restoration Areas #1-3, 7 and 8. See Appendix G for examples of improvements. | Improve water quality, increase wildlife habitat. Improve vegetative diversity and aesthetics. | Spring - Fall | 2004 through 2012: The City of Lakeville annually provides lakeshore owners with shoreline restoration information and encourages homeowners to take advantage of the Blue Thumb restoration program. Two residential shoreline restoration projects have been completed. One is located north of the beach area and one is on 175th St. W. 2007: A small area of lakeshore, near the boat launch, was restored using native plants. |

**Table 2: 2018 Recommended and Completed Management Actions for Crystal Lake
Black Dog Watershed Management Organization Habitat Monitoring**

| Problem Identified | Recommendation | Proposed Action | Benefits | Implementation Period | Completed Actions Which May Improve Wildlife Habitat and/or Water Quality |
|---|--|---|---|----------------------------|---|
| Curlyleaf pondweed dominates the lake in late spring-early summer. | Continue curlyleaf pondweed control measures. | Continue to control and manage. See Appendix A Blue Water Science report for locations of curlyleaf pondweed. | Increase wildlife habitat, improve water quality, vegetative diversity, aesthetics, and recreation. | Late Spring - Early summer | 1999 through 2018: The City of Burnsville conducts aquatic vegetation monitoring twice/year. 2003 through 2018: The City of Burnsville conducted annual harvesting of curlyleaf pondweed. |
| Eurasian watermilfoil is present. | Control Eurasian watermilfoil. | Control by chemical treatment. See Appdendix A Blue Water Science report for locations of Eurasian watermilfoil. | Increase wildlife habitat, improve water quality, vegetative diversity, aesthetics, and recreation. | Summer | |
| Common and glossy buckthorn are present | Control common and glossy buckthorn | Remove buckthorn. Volunteer groups and contractors can effectively remove buckthorn by pulling, cutting, and treating stumps with herbicide. See Appendix H for buckthorn management guidelines. See Appendix I for locations of buckthorn. | Increase wildlife habitat. Improve vegetative diversity and aesthetics | Fall | In 2009, the City of Burnsville treated 14 acres of buckthorn within Crystal Lake West Park (Appendix I). |
| Garlic mustard is present | Control garlic mustard | Organize a volunteer neighborhood group to pull garlic mustard. See Appendix I for locations of garlic mustard. | Increase wildlife habitat. Improve vegetative diversity and aesthetics | Spring | In 2008 and 2009, the City of Burnsville removed garlic mustard within the upland buffer (Appendix I) |
| Purple loosestrife is present. | Continue to control and manage purple loosestrife. | Control and manage. For a few small colonies of purple loosestrife, hand pull or dig the plants out before they go to seed. See Appendix I for locations of purple loosestrife. | Increase wildlife habitat. Improve vegetative diversity. | Spring - Fall | Purple loosestrife beetles were released by the MnDNR prior to 2002. Follow up monitoring by the MnDNR indicates that beetles are present at a population that the MnDNR feels is appropriate for biological control. |
| Bare soil areas are present along shoreline in Crystal Lake West Park area. | Re-vegetate bare soil areas to prevent soil erosion into Crystal Lake and create designated stone walkways for fishing access. | Exposed soil along the shoreline of Crystal Lake West Park Area could be re-vegetated to prevent shoreline erosion. Strategically located stones could provide fishing access to prevent disturbance of vegetation after it is established. (Potential Restoration Area #1 as shown in Figure 4 and photos) | Improve water quality and prevent erosion. | Spring - Fall | |
| Timber retaining wall in Tyecke Park area is in poor condition. | Repair timber retaining wall to prevent soil erosion into Crystal Lake. | Steep slopes in the Tyecke Park area are well protected with mature naturalized vegetation, however a timber retaining wall along the shoreline may need to be repaired or replaced to prevent slope destabilization and erosion. (Potential Restoration Area #2 as shown in Figure 4 and photos) | Prevent erosion | Winter | |
| Shoreline areas lacking naturalized vegetation within publicly owned beach area. Some areas have mowed turf grass close to the lakeshore edge. | Increase width and continuity of native upland buffer. | The upland buffer near the location of Plot #1C and shoreline to the south, and north of the beach area could be restored to naturalized native vegetation and not mowed (Potential Restoration Areas #3 and 4 as shown in Figure 4 and photos). | Increase wildlife habitat. Improve water quality. Improve vegetative diversity and aesthetics. | Spring - Fall | The width and density of naturalized shoreline buffer at the location of Emergent Plot #1B near the beach area has improved significantly since 2009. |
| Shoreline areas lacking naturalized vegetation within residential properties. Most of the residential properties have turf grass up the the lakeshore edge. | Increase width and continuity of native upland buffer. | Rather than manicured turf grass, the shoreline could be vegetated with native grasses and wildflowers. (Potential Restoration Area #5 as shown in Figure 4 and photos). | Increase wildlife habitat. Improve water quality. Improve vegetative diversity and aesthetics. | Spring - Fall | Six residential property owners have completed shoreline restortion projects using either City of Burnsville or Dakota Soil and Water Conservation District grants. |

Appendix E

2014 Lac Lavon MNRAM 3.4 Wetland Functional Assessment Results

Wetland Functional Assessment Summary

| Wetland Functional Assessment Summary | | | | | Maintenance of Hydrologic Regime | Flood/Stormwater/Attenuation | Downstream Water Quality | Maintenance of Wetland Water Quality | Shoreline Protection |
|---------------------------------------|---|--|---|--|----------------------------------|---------------------------------|-------------------------------|---|---------------------------------------|
| Wetland Name | Hydrogeomorphology | | | | | | | | |
| Lac Lavon | Depressional/Isolated (no discernable inlets or outlets), Lacustrine Fringe (edge of deepwater areas)/Shoreland | | | | 0.65 | 0.52 | 0.63 | 0.69 | 0.36 |
| | | | | | Moderate | Moderate | Moderate | High | Moderate |
| Additional Information | | | | | | | | | |
| Wetland Name | Maintenance of Characteristic Wildlife Habitat Structure | Maintenance of Characteristic Fish Habitat | Maintenance of Characteristic Amphibian Habitat | Aesthetics/Recreation/Education/Cultural | Commercial Uses | Ground-Water Interaction | Wetland Restoration Potential | Wetland Sensitivity to Stormwater and Urban Development | Additional Stormwater Treatment Needs |
| Lac Lavon | 0.50 | 0.68 | 0.05 | 0.76 | 0.00 | Combination Discharge, Recharge | 0.00 | 0.50 | 0.69 |
| | Moderate | High | Low | High | Not Applicable | | Not Applicable | Moderate | High |

Wetland Community Summary

| | | Vegetative Diversity/Integrity | | | | | | | |
|--------------|------------------|--------------------------------|-------------|---------------------------------|--------------------|-----------------------------|------------------------|------------------------|---------------------------------|
| Wetland Name | Location | Community | | | Wetland Proportion | Individual Community Rating | Highest Wetland Rating | Average Wetland Rating | Weighted Average Wetland Rating |
| | | Cowardin Classification | Circular 39 | Plant Community | | | | | |
| Lac Lavon | 19-114-21-11-001 | L2UBGh | Type 5 | Shallow, Open Water Communities | 95 | 0.5 | 0.50 | 0.50 | 0.50 |
| | | | | | | | Moderate | Moderate | Moderate |
| | | PSS1B | Type 6 | Shrub Carr | 5 | 0.5 | 0.50 | 0.50 | 0.50 |
| | | | | | | | Moderate | Moderate | Moderate |
| | | | | | 100 | | 0.50 | 0.50 | 0.50 |

☒ Denotes incomplete calculation data.

Management Classification Report for Lac Lavon

ID: 4

DWMO Strategic Waterbodies

DAKOTA County
Minnesota (Shakopee) Watershed, #33
Corps Bank Service Area 9

Based on the MnRAM data input from field and office review and using the classification settings as shown below, this wetland is classified as **Manage 1**

| Functional rank of this wetland based on MnRAM data | Functional Category | Self-defined classification value settings for this management level |
|---|---|--|
| Moderate | Vegetative Diversity/Integrity | High |
| Moderate | Habitat Structure (wildlife) | High |
| Low | Amphibian Habitat | Moderate |
| High | Fish Habitat | High |
| Moderate | Shoreline Protection | Moderate |
| High | Aesthetic/Cultural/Rec/Ed and Habitat | High / Moderate |
| Moderate | Stormwater/Urban Sensitivity and Vegetative Diversity | High / Moderate |
| High | Wetland Water Quality and Vegetative Diversity | High / Moderate |
| Moderate | Characteristic Hydrology and Vegetative Diversity | High / Moderate |
| Moderate | Flood/Stormwater Attenuation* | - |
| Not Applicable | Commercial use* | High |
| Moderate | Downstream Water Quality* | - |

The critical function that caused this wetland to rank as **Manage 1** was **Maintenance of Characteristic Fish Habitat**

Details of the formula for this action are shown below:

Maintenance of Characteristic Fish Habitat
$$\frac{[Q46*2)+Q24+Q18+Q20R+Q28+Q30+Q31+Q33R]}{9}$$

| Question | Value | Description |
|----------|-------|---------------------------------------|
| 18 | 1 | Sediment delivery |
| 20 | 1 | Stormwater runoff |
| 24 | 0.425 | Adjacent area Management |
| 28 | 1 | Nutrient loading |
| 30 | 0.1 | Shoreline rooted vegetation (%cover) |
| 31 | 0.1 | Shoreline wetland in-water width |
| 33 | 0.5 | Shoreline erosion potential |
| 46 | 1 | Fish habitat quality |

* The classification value settings for these functions are not adjustable

Management Classification Report for Lac Lavon

ID: 4

DWMO Strategic Waterbodies

DAKOTA County
Minnesota (Shakopee) Watershed, #33
Corps Bank Service Area 9

This report was printed on: Thursday, November 20, 2014

* The classification value settings for these functions are not adjustable

MnRAM Site Assessment Report

Thursday, November 20, 2014

Wetland: Lac Lavon

Project: BDWMO Strategic Waterbodies

Wetland ID: 4, Township 114, Section 11, Range 21, , ,

DAKOTA County, Minnesota (Shakopee) Watershed, Corps Bank Service Area #9

Assessment Purpose: Inventory

A site visit was made to this wetland on 7/2/2014 by KSW. Site conditions were Normal. This wetland is estimated to cover 60 acres.

This report reflects conditions on the ground at the date of the assessment and, unless noted or implicit in the standard questions, does not reflect speculation on the future or past conditions.

This wetland is located in or near the city of Lakeville

General Features

Hydrogeomorphology

The maximum water depth at this site is 120 inches, with 95 percent inundated. With an immediate drainage area of 185 acres, it is doubtful that this wetland is sustainable given its small catchment area.

As a Depressional/Isolated wetland, this site has no discernable inlets or outlets. As such, it is valued for its ability to store water, especially if located lower in the watershed. If it does not already have invasive species in the plant community, its lack of connection to upstream sites with such species may protect it.

As a Lacustrine Fringe wetland, this site located at the edge of deepwater areas and may be considered shoreland. As such, it protects from possible erosive wave effects and may be used as a spawning area for fish.

This wetland has been drained or altered 0% from its original size of 60 acres.

Soils

The soils in the immediate wetland area are primarily Pits, gravel. The adjacent upland, to about 500 feet, is Wadena loam, 2 to 6 percent slopes.

Vegetation and Upland Buffer

The extent of vegetation in this wetland is about 50 percent and the naturalized buffer width averages 5 feet. Vegetated buffers around wetlands provide multiple benefits including wildlife habitat, erosion protection, and a reduction in surface water runoff.

This buffer provides very little, if any, protection of water quality or habitat for wildlife.

As a shoreline wetland, this site has the potential to protect from erosion and provide spawning and nursery habitat for fish and wildlife. Wetlands located in areas with strong currents and wave action have the greatest potential for protecting shoreline. Shorelines composed of sandy or erodible soils will benefit the most from shoreline wetland protection.

Special Features

- F Public park, forest, trail or recreation area.
- K Local Shoreland Management Plan area.
- M Shoreland area identified in a zoning ordinance.

Vegetative Communities

The following plant communities were observed:

(See Appendix A for details on the Dominant Species per plant community)

Shallow, Ow Communities Type 5, L2UBGh. This community had a vegetative index of moderate and comprised 95 percent of the entire area.

Shrub-carr Type 6, PSS1B. This community had a vegetative index of moderate and comprised 5 percent of the entire area.

The highest rated community was the Shallow Marsh community rated at 0.5. Averaging all the communities together, the Vegetative Diversity and Integrity of this wetland is Moderate. A more accurate look uses a weighted average; using this method, this site shows a Moderate Vegetative Diversity and Integrity.

The weighted average provides the best measure for an entire wetland. Plant communities at this site are, overall, of average quality. Individual community ratings should be examined to provide a complete picture of possible high-value communities or smaller-but-poor-quality segments that might degrade the site over time.

Functional Ratings

| <i>Function</i> | <i>Rating</i> | <i>Comment</i> |
|---------------------------------------|---------------|---|
| Vegetative Diversity | Moderate | Moderate-functioning vegetative communities indicate a presence of native wetland species with substantial non-native or invasive species. |
| Additional stormwater treatment needs | High | Because the maintenance of wetland water quality index is high, no additional treatment is called for. |
| Maintenance of Hydrologic Regime | Moderate | There has been some degree of human alteration of the wetland hydrology, either by outlet control or by altering immediate watershed conditions. However, the wetland retains some of the hydrologic regime similar to the original wetland type, either in part of the wetland or overall to some extent. Because of the interference (whether active or inadvertant), some characteristic vegetative communities have likely been affected, as also have the functions of flood attenuation, water quality and groundwater interaction. |
| Flood/Stormwater/Attenuation | Moderate | The wetland provides some flood storage and/or flood wave attenuation. It may have either an altered or unrestricted outlet, disturbed wetland soils, thin or little emergent vegetation (with channels) or it may be situated high in a watershed with a low proportion of impervious surfaces, moderate runoff volumes, loamy upland soils, and one or more other wetlands present within the subwatershed. |

| | | |
|--|----------------|--|
| Downstream Water Quality | Moderate | This wetland has some ability and opportunity to protect downstream resources. The ability of the wetland to remove sediment from stormwater is determined by emergent vegetation and overland flow characteristics. A high nutrient removal rating indicates dense vegetation and sheet flow to maximize nutrient uptake and residence time within the wetland. The opportunity for a wetland to protect a valuable water resource diminishes with distance from the wetland so wetlands with valuable waters within 0.5 miles downstream have the greatest opportunity to provide protection, as do those that receive more (and less-treated) runoff. |
| Maintenance of Wetland Water Quality | High | Wetland water quality is high, indicating little need for additional treatment. As long as upland land use and existing buffer conditions do not change, this wetland can be expected to sustain current characteristics. |
| Shoreline Protection | Moderate | This fringe site provides some protection against erosive action. Reducing the amount of buffer that is manicured would further protect the adjacent water resource, as would increasing the buffer width. |
| Maintenance of Characteristic Wildlife Habitat Structure | Moderate | The site provides good habitat and is relatively accessible to wildlife, although it may be somewhat isolated on the landscape and lack the rich vegetative community and complex structure that would support a wider range of wildlife. |
| Maintenance of Characteristic Fish Habitat | High | The site has a direct connection to spawning or nursery habitat, or may provide refuge or shade for native species of fish. Low amounts of sediment mean that eggs are not smothered; good water quality supports fish health. |
| Maintenance of Characteristic Amphibian Habitat | Low | Predatory fish are always present and winter habitat unsuitable as site often freezes to the bottom. High inputs of untreated stormwater or unfiltered runoff contribute to poor water quality and reproductive conditions. |
| Aesthetics/Recreation /Education/Cultural | High | Regardless of actual integrity, the site is accessible and valued by significant populations of people. Its value is enhanced by not being visibly altered by human influences such as trash or roads. There is a high evidence it is used for multiple recreational activities. |
| Wetland restoration potential | Not Applicable | Because restoration would affect permanent structures or infrastructure (houses, roads, septic systems), this site is not suitable for restoration. |
| Wetland Sensitivity to Stormwater and Urban Development | Moderate | This wetland is moderately sensitive to stormwater; Floodplain forests, fresh wet meadows dominated by reed canary grass, shallow and deep marshes dominated by cattail, reed canary grass, giant reed or purple loosestrife, and shallow, open water communities with low to moderate vegetative diversity. |

Appendix A: Dominant Species By Plant Community

| | Wetland Type | Plant Community | Dominant Species | Percent Cover |
|------|--------------|-------------------------|---------------------------|---------------|
| L2UB | Type 5 | Shallow, Ow Communities | Sago pondweed | >3-<10% |
| | | | American pondweed | 0-3% |
| | | | Water stargrass | >3-<10% |
| | | | Limp white water crowfoot | >3-<10% |
| | | | Leafy pondweed | >10-25% |
| | | | Illinois pondweed | >3-<10% |
| | | | Flat-stemmed pondweed | 0-3% |
| | | | Eurasian water milfoil | >10-25% |
| | | | Curly pondweed | >10-25% |
| | | | Canadian elodea | >3-<10% |
| | | | Common coontail | >25-50% |
| | | | Flexuous naiad | 0-3% |
| PSS1 | Type 6 | Shrub-carr | Black willow | 0-3% |
| | | | Common mint | 0-3% |
| | | | Common dandelion | 0-3% |
| | | | Common boneset | 0-3% |
| | | | Canada thistle | 0-3% |
| | | | American slough grass | 0-3% |
| | | | Blue vervain | 0-3% |
| | | | Cottonwood | 0-3% |
| | | | Curly dock | 0-3% |
| | | | Bristly sedge | 0-3% |
| | | | Prickly lettuce | 0-3% |
| | | | Tussock sedge | 0-3% |
| | | | Swamp milkweed | 0-3% |
| | | | Stinging nettle | 0-3% |
| | | | Spotted touch-me-not | 0-3% |
| | | | Soft stem bulrush | 0-3% |
| | | | Sensitive fern | 0-3% |
| | | | Sandbar willow | >25-50% |
| | | | Reed canary grass | >10-25% |
| | | | Common ragweed | 0-3% |
| | | | Purple loosestrife | 0-3% |
| | | | Yarrow | 0-3% |
| | | | Peach-leaved willow | >10-25% |
| | | | Northern bugleweed | 0-3% |
| | | | Northern blue flag | 0-3% |
| | | | Narrow-leaved cattail | 0-3% |

| | |
|-------------------|---------|
| Green ash | 0-3% |
| Golden alexanders | 0-3% |
| Giant goldenrod | 0-3% |
| Fox sedge | 0-3% |
| Fowl bluegrass | >10-25% |
| Red maple | 0-3% |

MnRAM: Site Response Record

For Wetland Lac Lavon

Location: 19-114-21-11-001

BDWMO Strategic Waterbodies

Plant Community: Shallow, Open Water C

Cowardin Classification: L2UBGh
Circular 39: Type 5

Plant Community: Shrub Carr

Cowardin Classification: PSS1B
Circular 39: Type 6

- 4 Listed, rare, special species?
- 5 Rare community or habitat?
- 6 Pre-European-settlement condition?

Hydrogeomorphology / topography:

7 Depressional/Isolated, Lacustrine

- 8-1 Maximum water depth 120 inch
- 8-2 % inundated 95%
- 9 Immediate drainage--local WS 185 acres
- 10 Estimated size/existing site: (see #66)

11-Upland Soil Wadena loam, 2 to 6 percent slopes

11-Wetland Soil Pits, gravel

- 12 Outlet for flood control
- 13 Outlet for hydro regime
- 14 Dominant upland land use
- 15 Wetland soil condition
- 16 Vegetation (% cover)
- 17 Emerg. veg flood resistance
- 18 Sediment delivery
- 19 Upland soils (soil group)
- 20 Stormwater runoff
- 21 Subwatershed wetland density
- 22 Channels/sheet flow

23 Adjacent buffer width

Adjacent area management

- 24-A Full
- 24-B Manicured
- 24-C Bare

Adjacent area diversity/structure

- 25-A Native
- 25-B Mixed

25-C Sparse

Adjacent area slope

- 26-A Gentle
- 26-B Moderate
- 26-C Steep

- 27 Downstream sens./WQ protect.
- 28 Nutrient loading

29 Shoreline wetland?

Shoreline Wetland

- 30 Rooted veg., % cover
- 31 Wetland in-water width
- 32 Emerg. veg. erosion resistance
- 33 Erosion potential of site
- 34 Upslope veg./bank protection
- 35 Rare wildlife?
- 36 Scarce/Rare/S1/S2 community
- 37 Vegetative cover
- 38 Veg. community interspersed
- 39 Wetland detritus
- 40 Interspersion on landscape
- 41 Wildlife barriers

Amphibian-breeding potential

- 42 Hydroperiod adequacy
- 43 Fish presence
- 44 Overwintering habitat
- 45 Wildlife species (list)
- 46 Fish habitat quality
- 47 Fish species (list)
- 48 Unique/rare opportunity
- 49 Wetland visibility
- 50 Proximity to population
- 51 Public ownership
- 52 Public access
- 53 Human influence on wetland
- 54 Human influence on viewshed
- 55 Spatial buffer
- 56 Recreational activity potential

57 Commercial crop--hydro impact

Groundwater-specific questions

- 58 Wetland soils Recharge
- 59 Subwatershed land use Discharge
- 60 Wetland size/soil group Recharge
- 61 Wetland hydroperiod Discharge
- 62 Inlet/Outlet configuration Recharge
- 63 Upland topo relief Discharge

Additional information

- 64 Restoration potential
- 65 LO affected by restoration
- 66 Existing size
- Restorable size
- Potential new wetland
- 67 Average width of pot. buffer
- 68 Ease of potential restoration
- 69 Hydrologic alterations
- 70 Potential wetland type
- 71 Stormwater sensitivity
- 72 Additional treatment needs

Watershed Minnesota (Shakopee)

WS# 33 Service Area: 9

For functional ratings, please run the Summary tab report.

This report printed on: 11/20/2014

Appendix F

Descriptions of MNRAM Wetland Functions

Appendix D

Descriptions of MNRAM Wetland Functions

6.0 Functional Rating Formulas

GENERAL NOTE: Some questions are not applicable to particular wetlands and will be scored N/A. In these cases, rather than count N/A as zero, an alternate equation is provided that eliminates the question from the formula altogether. Because not every question has N/A as an option, formulas that do not include N/A-option questions have only one configuration.

Formulas with a “reverse rating” (marked as “R”) take the actual response and “flip” its value for the calculation, so that a question response of “A” high (value of 1.0) will be calculated as low (value of 0.1). In such a formula, medium ratings stay medium.

6.1 VEGETATIVE DIVERSITY/INTEGRITY

Table 3: Vegetative Diversity/Integrity Summary

The functional rating is based primarily on the diversity of vegetation within the wetland in comparison to an undisturbed condition for that wetland type. An exceptional rating results from one of the following conditions: 1) highly diverse wetlands with virtually no non-native species, 2) rare or critically impaired wetland communities in the watershed, or 3) the presence or previous siting of rare, threatened, or endangered plant species. A high rating indicates the presence of diverse, native wetland species and a lack of non-native or invasive species. Wetlands that rate low are primarily dominated by non-native and/or invasive species.

This table may be used when calculating Vegetative Diversity/Integrity Functional Index manually. It shows four options for calculating and presenting floristic data. If you are entering data directly into the MnRAM 3.0 database, this table does not apply.

| | 3A Proportion of Wetland | 3B Individual Community Scores | 3C Highest Quality | 3D Non-Weighted Average | 3E Weighted Average |
|---------------------------------|---|---|-----------------------------------|--|---|
| Community #1 | T | A | | A | A |
| Community #2 | U | B | | B | B |
| Community #3 | V | C | | C | C |
| Community #4 | W | D | | D | D |
| Community #5 | X | E | | E | E |
| Community #6 | Y | F | | F | F |
| Community #7 | Z | G | | G | G |
| Wetland Rating Value | 1.0 | | Highest Value | $(A+B+C+D+E+F+G)/7 = \text{Ave.}$ | $(A*T)+(B*U)+(C*V)+(D*W)+(E*X)+(F*Y)+(G*Z) = \text{Wt. Ave.}$ |

If any questions #4-6 are answered yes and/or if any of the Special Features b, d, or i have been selected, enter Exceptional for the functional index. If not, compute the contribution to vegetative diversity and integrity by each plant community by doing the following: multiply the ranking for each community (Question #3b) by its total proportion in Question 3a (percent of total). Then, the functional index for the entire wetland can be calculated four ways (as follows) and should be utilized according to the scope of the project:

3b) Individual Community Scores: maintain raw data as recorded.

3c) Highest Quality Community: report the highest-functioning community.

3d) Non-Weighted Average Quality of all Communities: straight average

3e) Weighted Average Quality Based on Percentage of Each Community: multiply each community rating by its percentage, then add all together.

| Vegetative Diversity/ Integrity | | | | | |
|------------------------------------|---------------------------------|--|---|-----------------------------------|--|
| | 3a. Proportion of Wetland | 3b. Individual Community Scores | 3c. Highest Rated Community Quality | 3d. Non- Weighted Average | 3e. Weighted Average |
| Community #1 | T | A | If Spec. Features b, d or i are checked then rate Exceptional (2); if either question 4, 5, or 6 are Yes, then rate Exceptional (2); else: | | |
| Community #2 | U | B | | | |
| Community #3 | V | C | | | |
| Community #4 | W | D | | | |
| Community #5 | X | E | | | |
| Community #6 | Y | F | | | |
| Community #7 | Z | G | | | |
| Overall Wetland Value Rating | 1.0 | | : Highest Value of A-G | : (A+B+C+ D+E+F+G)/7 = Ave. | :(A*T)+(B* U)+(C*V)+ (D*W)+(E* X)+(F*Y)+(G*Z) = Wt. Ave. |

6.2 MAINTENANCE OF CHARACTERISTIC HYDROLOGIC REGIME

A wetland's hydrologic regime or hydroperiod is the seasonal pattern of the wetland water level that is like a hydrologic signature of each wetland type. It defines the rise and fall of a wetland's surface and subsurface water. The constancy of the seasonal patterns from year to year ensures a reasonable stability for the wetland²³. The ability of the wetland to maintain a hydrologic regime characteristic of the wetland type is evaluated based upon wetland soil and vegetation characteristics, land use within the wetland, land use within the upland watershed contributing to the wetland, and wetland outlet configuration. Maintenance of the hydrologic regime is important for maintaining a characteristic vegetative community, and is closely associated with other functions including flood attenuation, water quality and groundwater interaction.

Measures the degree of human alteration of the wetland hydrology, either by outlet control or by altering immediate watershed conditions. Each parameter is weighted equally.

| MnRAM # | Excel # | Variable Description | Type of Interaction |
|---------|---------|--|---------------------|
| 13 | E17 | Outlet—natural hydrologic regime | Controlling |
| 14 | E18 | Dominant upland land use | Compensatory |
| 15 | E19 | Soil condition/wetland | Compensatory |
| 20 | F24 | Stormwater runoff/pretreatment-Reverse | Compensatory |

Hydrologic Regime Index = (13+14+15+20)/4

6.3 FLOOD AND STORMWATER STORAGE/ATTENUATION

A wetland's ability to provide flood storage and/or flood wave attenuation is dependent on many characteristics of the wetland and contributing watershed. Characteristics of the subwatershed that affect the wetlands ability to provide flood storage and attenuation include: soil types, land use and resulting stormwater runoff volume, sediment delivery from the subwatershed, and the abundance of wetlands and waterbodies in the subwatershed. Wetland characteristics which affect the wetland's ability to store and or attenuate stormwater include: condition of wetland soils; presence, extent, and type of wetland vegetation; presence and connectivity of channels; and most importantly outlet configuration. Higher rated wetlands will have an unaltered or restricted outlet, undisturbed wetland soils, dense emergent vegetation without channels, a high proportion of impervious surfaces in the subwatershed, large runoff volumes, clayey upland soils, and few wetlands present within the subwatershed.

This formula is based on the Surface Water Storage Functional Capacity Index scoring concept and equation²⁴. The formula was altered with the addition of three surface flow characteristics and two stormwater runoff parameters (Stormwater Runoff Quality/Quantity and Subwatershed Wetland Density) along with the removal of two parameters (Soil Porosity and Subsurface Outlet,

²³ Mitsch and Gosselink, 2000

²⁴ Lee et al., 1997

which is already characterized in another parameter). This index is comprised of 5 primary processes, which are weighted equally; included in each major process are one to three characteristics that equally contribute to that process.

1. **Outlet Characteristics:** Outlet characteristics
2. **Upland Watershed:** Upland land use, Upland soils,
3. **Wetland Condition/Land Use:** Wetland land use, sediment delivery
4. **Runoff Characteristics:** Stormwater runoff quality/quantity, subwatershed wetland density
5. **Surface Flow Characteristics:** Flow-through emergent vegetation density, surface flow characteristics

Flood and Stormwater Storage Index Computation:

Entire Formula: Outlet for flood retention{ 12 } + (Dominant upland use-RR{ 14 }+ Upland soils{ 19 })/2 + (Soil condition{ 15 } + Sediment delivery{ 18 })/2 + Stormwater runoff pretreat&det{ 20 } + Subwatershed wetland density { 21 })/2 + (Percent emergent vegetative cover{ 16 } + Flow-through emergent vegetative roughness{ 17 } + Channels/sheet flow{ 22 })/3)/5.

1. If 12=0, then: $((14+19)/2+(15+18)/2+(20+21)/2+(16+17+22)/3)/4$
2. If 12>0, then: $(12+(14+19)/2+(15+18)/2+(20+21)/2+(16+17+22)/3)/5$

No changes to the formula are necessary if 16=0.

Flood and Stormwater Storage/Attenuation Variables

| MnRAM # | Excel # | Variable Description | Type of Interaction |
|---------|---------|--------------------------------------|----------------------|
| 12 | E16 | Outlet—flood attenuation | Controlling—optional |
| 14 | F18 | Dominant upland land use-RR | Compensatory |
| 19 | E23 | Upland soils | Compensatory |
| 15 | E19 | Soil condition | Compensatory |
| 18 | E22 | Sediment delivery | Compensatory |
| 20 | E24 | Stormwater pretreatment &detention | Compensatory |
| 21 | E25 | Subwatershed wetland density | Compensatory |
| 16 | F20 | Emergent vegetation % cover | Comp.—optional |
| 17 | E21 | Emergent vegetation flood resistance | Comp.—optional |
| 22 | E26 | Channels/sheet flow | Compensatory |

6.4 DOWNSTREAM WATER QUALITY PROTECTION

This rates the wetland's ability and opportunity to protect valuable downstream resources. Valuable downstream resources include recreational waters (i.e. lakes, streams, rivers, creeks, etc) and potable water supplies. The level of functioning is determined based on runoff characteristics, sedimentation processes, nutrient cycling, and the presence and location of significant downstream water resources. Runoff characteristics that are evaluated include: land use and soils in the upstream watershed, the stormwater delivery system to the wetland, and sediment delivery characteristics. The ability of the wetland to remove sediment from stormwater is determined by emergent vegetation and overland flow characteristics. A high nutrient removal rating indicates dense vegetation and sheet flow to maximize nutrient uptake and residence time within the wetland. The opportunity for a wetland to protect a valuable water resource diminishes with distance from the wetland so wetlands with valuable waters within 0.5 miles downstream have the greatest opportunity to provide protection.

Compute Functional Index for Downstream Water Quality Protection

This functional index computation was derived from a combination of Nutrient Cycling and Retention of Particulates functions in the HGM Prairie Pothole draft guidebook⁵⁴ with the downstream sensitivity concept from *The Minnesota Wetland Evaluation Methodology*. Three major processes make up equal portions of the Downstream Water Quality Protection function²⁵ with a measure of opportunity to protect downstream resources; each process is comprised of two to four observable parameters.

1. **Rate, Quantity, and Quality of Runoff to the Wetland:** this is characterized by the conditions in the upstream watershed; both land use and soils, that affect the sediment and nutrient loads to the wetland, and by the existing storm water delivery system to the wetland (Upland watershed conditions, storm water runoff, evidence of sediment delivery, and upland buffer each comprise 1/16 of the entire downstream water quality functional index based on their contribution to sediment removal).
2. **Sedimentation:** this is characterized by the presence of flow-through emergent vegetation density and by the overland flow characteristics within the wetland. A wetland with primarily sheet flow through the wetland and dense emergent vegetation density will allow sediment to drop out more effectively than a wetland with channel flow and no vegetation (When all parameters are applicable; emergent vegetative density and overland flow characteristics each make up 1/8 of the total downstream water quality functional index based on their contribution to sediment removal).
3. **Nutrient Uptake:** this is characterized by the outlet configuration and vegetative characteristics. A wetland with long water retention times has more capacity to remove nutrients from the water column via physical and biological processes. Vegetation slows floodwaters by creating frictional drag in proportion to stem density which allows sediment particles to settle out, thereby improving the water quality for downstream uses (Outlet characteristics and vegetative density each make up 1/8 of the total downstream water quality functional index based on their contribution to nutrient uptake).

²⁵ Derived from a combination of Nutrient Cycling and Retention of Particulates functions in the HGM Prairie Pothole draft guidebook (Lee et al., 1997) with the downstream sensitivity concept from *The Minnesota Wetland Evaluation Methodology*.

4. **Downstream Sensitivity:** if the wetland contributes to the maintenance of water quality within one-half mile of a recreational water body or potable water supply source downstream, it operates at a higher functioning level than a similar wetland farther from or without significant downstream water resources (This factor accounts for ¼ of the total downstream water quality functional index).

Downstream Water Quality Functional Index Computations:

1. If $12=0$, then: $(14+20_{\text{reversed}} + 18 + (23+24+26)/3 + (16+17)/2 + 27)/6$
 2. If $12>0$, then: $(14+20_{\text{reversed}} + 18 + (23+24+26)/3 + (16+17)/2 + 27 + 12)/7$

No changes to the formula are necessary if $16=0$.

Entire Formula:

(Dominant upland land use{ 14} + Stormwater runoff pretreatment & detention{ 20_{reversed} } + Sediment delivery { 18} + (Upland buffer width{ 23}WQ + Upland buffer vegetative cover{ 24} + Upland buffer slope { 26})/3 + (Flow-through %emergent vegetative cover{ 16} + Flow-through emergent vegetative roughness{ 17})/2 + Downstream sensitivity{ 27}+ Outlet for flood{ 12})/7

Downstream Water Quality Variables

| MnRAM # | Excel # | Variable Description | Type of Interaction |
|---------|---------|---|-----------------------|
| 14 | E18 | Dominant upland land use | Controlling |
| 20 | E24 | Stormwater runoff pretreatment &detention | Controlling |
| 18 | E22 | Sediment delivery | Controlling |
| 23 | G27 | Upland buffer width | Comp. |
| 24 | G28 | Upland area management | Comp. |
| 26 | G34 | Upland area slope | Comp. |
| 16 | F20 | Emergent vegetation (% cover) | Comp.—optional |
| 17 | E21 | Emergent vegetation (roughness coefficient) | Comp.—optional |
| 27 | E39 | Downstream sensitivity | Comp. |
| 12 | E16 | Outlet for flood | Controlling--optional |

6.5 MAINTENANCE OF WETLAND WATER QUALITY

The sustainability of a wetland is partially driven by the quality and quantity of stormwater runoff entering the wetland. The ability of the wetland to sustain its characteristics is evaluated based on characteristics of the contributing subwatershed and indicators within the wetland. Subwatershed conditions which affect the wetland's sustainability in relation to water quality impacts include: upland land use; sediment delivery characteristics to the wetland; stormwater runoff volumes and rates; and the extent, condition, and width of upland buffer. Indicators of nutrient loading to the wetland indicate that a diverse wetland may not be sustainable. Indicators that a wetland has been affected by nutrient loading include the presence of monotypic vegetation and/or algal blooms.

This functional index was derived from a combination of sources including MNRAM, HGM, WEM, WET, and experiences of the project team. The sustainability of a wetland

is partially driven by the quality and quantity of stormwater runoff entering the wetland. The ability of the wetland to sustain its characteristics is evaluated based on characteristics of the contributing subwatershed and indicators within the wetland. Subwatershed conditions which affect the wetland's sustainability in relation to water quality impacts include: upland land use; sediment delivery characteristics to the wetland; stormwater runoff volumes and rates; and the extent, condition, and width of upland buffer. Indicators of nutrient loading to the wetland indicate that a diverse wetland may not be sustainable. Indicators that a wetland has been affected by nutrient loading include the presence of monotypic vegetation and/or algal blooms.

Wetland Water Quality Functional Index Computation:

$$(3e*2+14+20_{\text{reversed}} + (23+24+26)/3+18+28)/7$$

Entire Formula:

(Vegetative Diversity/Integrity{3e*2} + Dominant upland land use{14} + Stormwater runoff pretreatment & detention{20_{reversed}} + (Upland buffer width{23}WQ + Upland buffer vegetative cover {24} + Upland buffer slope {26})/3 + Sediment delivery {18})/2 + Nutrient loading {28})/7

Wetland Water Quality Variables

| MnRAM # | Excel # | Variable Description | Type of Interaction |
|---------|---------|---|---------------------|
| 3e | D6*2 | Vegetative Diversity/Integrity | Contributing |
| 14 | E18 | Dominant upland land use | Contributing |
| 20 | F24 | Stormwater runoff pretreatment and detention—RR | Contributing |
| 23 | G27 | Upland buffer width | Contributing |
| 24 | G28 | Upland area management | Contributing |
| 26 | G34 | Upland area slope | Contributing |
| 18 | E22 | Sediment delivery | Contributing |
| 28 | E40 | Nutrient loading | Contributing |

This functional index was derived from a combination of sources including MNRAM, HGM, WEM, WET, and experiences of the project team. The sustainability of a wetland is partially driven by the quality and quantity of stormwater runoff entering the wetland. The ability of the wetland to sustain its characteristics is evaluated based on characteristics of the contributing subwatershed and indicators within the wetland. Subwatershed conditions which affect the wetland's sustainability in relation to water quality impacts include: upland land use; sediment delivery characteristics to the wetland; stormwater runoff volumes and rates; and the extent, condition, and width of upland buffer. Indicators of nutrient loading to the wetland indicate that a diverse wetland may not be sustainable. Indicators that a wetland has been affected by nutrient loading include the presence of monotypic vegetation and/or algal blooms.

6.6 SHORELINE PROTECTION

Shoreline protection is evaluated only for those wetlands adjacent to lakes, streams, or deepwater habitats. The function is rated based on the wetlands opportunity to protect the shoreline; i.e. wetlands located in areas frequently experiencing large waves and high

currents have the best opportunity to protect the shore. In addition, shore areas composed of sands and loams with little vegetation or shallow-rooted vegetation will benefit the most from shoreline wetlands. The wetland width, vegetative cover, and resistance of the vegetation to erosive forces determine the wetland's ability to protect the shoreline.

Each of the five parameters contributes equally²⁶: based primarily on the characteristics presented in WEM with a simple, straightforward computation of the index assuming all characteristics contribute equally.

| MnRAM # | Excel # | Variable Description | Type of Interaction |
|---------|---------|--|---------------------|
| 29 | E41 | Shoreline? | Controlling |
| 30 | E42 | Rooted shoreline vegetation (% cover) | Contributing |
| 31 | E43 | Wetland width (average) | Contributing |
| 32 | E44 | Emergent vegetation erosion resistance | Contributing |
| 33 | E45 | Shoreline erosion potential | Contributing |
| 34 | E46 | Bank protection ability | Contributing |

Shoreline Protection Functional Index Computation:

If 29=1, then:

$$\text{Shoreline Protection Index} = (30+31+32+33+34)/5$$

Entire Formula:

(Rooted shoreline vegetation {30} + Average shoreline wetland width {31} + Emergent vegetation erosion resistance {32} + (Shoreline erosion potential {33} + Bank protection ability {34})/5

6.7 MAINTENANCE OF CHARACTERISTIC WILDLIFE HABITAT STRUCTURE

The ability of a wetland to support various wildlife species is difficult to determine due to the specific requirements of the many wildlife species that utilize wetlands. This function determines the value of a wetland for wildlife in a more general sense, and not based on any specific species. The characteristics evaluated to determine the wildlife habitat function include: vegetative quality, outlet characteristics (which control hydrologic regime), upland land use, wetland soil type and conditions, water quality of storm water runoff entering the wetland, upland buffer extent, condition, and diversity; the interspersions of wetlands in the area; barriers to wildlife movement; wetland size; vegetative and community interspersions within the wetland; and amphibian breeding potential and overwintering habitat.

Thirteen parameters are weighed equally as described below; vegetative quality weighted double the other factors. The questions are borrowed or modified from MNRAM, WET, WEM, and HGM methodologies, combined to provide a measure of wildlife habitat in general, not focusing on any particular species.

If Rare Wildlife (35) or Rare Natural Community (36) are true, then this Index is Exceptional.

²⁶ Based primarily on the characteristics presented in WEM.

If Special Features d, g, or j are checked, then this Index is Exceptional, otherwise, follow conditions below:

If 37=0 and 38=0 and 39=0, then:

$$(3e*2+40+41+(23+24+25)/3+13+20)/7$$

If 38=0 and 39=0, then:

$$(3e*2+37+40+41+(23+24+25)/3+13+20)/8$$

If 37=0 and 39=0, then:

$$(3e*2+38+40+41+(23+24+25)/3+13+20)/8$$

If 37=0 and 38=0, then:

$$(3e*2+39+40+41+(23+24+25)/3+13+20)/8$$

If 39=0, then:

$$(3e*2+37+38+40+41+(23+24+25)/3+13+20)/9$$

If 38=0, then:

$$(3e*2+39+37+40+41+(23+24+25)/3+13+20)/9$$

If 37=0, then:

$$(3e*2+39+38+40+41+(23+24+25)/3+13+20)/9$$

If 37>0 and 38>0 and 39>0, then:

$$(3e*2+39+37+38+40+41+(23+24+25)/3+13+20)/10$$

Entire Equation:

(Vegetative Diversity/Integrity {3e*2} + Wetland Detritus {39} + Vegetation Interspersion {37} + Community Interspersion {38} + Wetland Interspersion {40} + Wildlife Barriers {41} + (Upland buffer width {23}WQ + Upland Area Management{24} + Upland area diversity {25})/3 + Outlet natural hydrologic regime {13}+ Stormwater runoff pretreatment and detention 20)/11

| MnRAM # | Excel # | Variable Description | Type of Interaction |
|---------|---------|--|---------------------|
| 41 | E53 | Wildlife barriers | Controlling |
| 3e | D6 | Vegetative Ranking (communities' weighted average) | Compensatory |
| 39 | E51 | Wetland detritus (n/a) | |
| 23 | I27 | Upland buffer average width | |
| 24 | G28 | Upland area management | |
| 25 | G31 | Upland area diversity | |
| 13 | E17 | Outlet natural hydrologic regime | |
| 20 | F24 | Stormwater runoff pretreatment & detention—RR | |
| 37 | F49 | Vegetation interspersion (n/a) | |
| 38 | F50 | Community interspersion (n/a) | |
| 40 | E52 | Wetland interspersion | |

6.8 MAINTENANCE OF CHARACTERISTIC FISH HABITAT

The ability of the wetland to support native fish populations is determined by structural factors within the wetland as well as water quality contributions from upland factors. Wetlands rated High are lacustrine or riverine and provide spawning/nursery habitat, or

refuge for native species (included but not limited to game fish). Wetlands rated Low for fish habitat do not have a direct hydrologic connection to a waterbody with a native fishery or have poor water quality.

| MnRAM # | Excel # | Variable Description | Type of Interaction |
|---------|---------|-----------------------------|---------------------|
| 46 | E58*2 | Fish habitat quality | Controlling |
| 29 | D41 | Fringe wetland? | Contributing |
| 24 | G28 | Adjacent area management | Compensatory |
| 18 | E22 | Sediment delivery | Compensatory |
| 20 (R) | F24 | Storm water runoff | Compensatory |
| 28 | E40 | Nutrient load | Compensatory |
| 30 | E42 | Percent cover | Compensatory |
| 31 | E43 | Wetland shoreline width | Compensatory |
| 33 (R) | F45 | Shoreline erosion potential | Compensatory |

Fish Habitat Functional Index Computation:

If Special Features a or g are checked, then Fishery Habitat Index = Exceptional.

If 46=0, then Fishery Habitat = N/A

If 29=0, Fishery Habitat Index = $[(46*2)+24+18+20(R) +28]/6$

If 29>0, Fishery Habitat Index = $[(46*2)+24+18+20(R) +28+30+31+33(R)]/9$

6.9 MAINTENANCE OF CHARACT. AMPHIBIAN HABITAT FOR BREEDING/OVERWINTERING

The ability of a wetland to support various amphibian species is difficult to determine due to the specific requirements of the many amphibian species that depend on wetlands. This function determines the value of a wetland for amphibians in general, not based on specific species. An adequate wetland hydroperiod and the presence or absence of predatory fish are considered to be limiting variables for this function. In general, wetlands must remain inundated until early to mid-June to allow the larval stages to metamorphose into adults. Because many amphibians are partly terrestrial, the characteristics evaluated to determine the amphibian habitat function include numerous hydrology and terrestrial measures. The characteristics evaluated include: upland land use, upland buffer width, water quality of storm water runoff entering the wetland, barriers to wildlife movement, and amphibian breeding potential and overwintering habitat.

An adequate wetland hydroperiod (Question 42) is considered to be the primary limiting variable for this functional index. If the hydroperiod is insufficient for breeding, the wetland rating for amphibian use will be Not Sufficient. The status of predatory fish in the wetland (Q.43) is a secondary limiting factor to the final rating; the lowest rating for this variable, however, is 0.1 (Low), rather than zero (Not Sufficient).

Amphibians' ability to use a particular wetland for over wintering is a contributing factor in rating the wetland's functional index (Q.44). Because most amphibians are partly terrestrial, the extent of upland buffer habitat surrounding the wetland (Q.23) is an

important habitat component²⁷ and is weighted by a factor of two. Question 14 (Upland Land Use) is also included as an indicator of the quality of the surrounding upland habitat⁵⁶. Unnatural fluctuations in water depth in wetlands from conducted storm water runoff can impair reproductive success in amphibians, which often attach their eggs to stems of wetland vegetation, e.g., salamanders, tree frogs, green frogs, and wood frogs²⁸. Extreme water level fluctuations during winter may also cause mortality in overwintering reptiles and amphibians²⁹. Thus, Question 20 is included in the formula, with a reverse rating. Question 41 (Barriers) is included because access to and from the wetland by amphibians is an important factor in habitat quality³⁰.

Amphibian Habitat Functional Index Computation:

If 42=0, then N/A

Otherwise: Amphibian Habitat Index = $(43) * [(44 + 2 * 23_{\text{wildlife}} + 14 + 41 + 20_{\text{reversed}}) / 6]$

Entire Formula:

If Amphibian Breeding Potential-Hydroperiod {42} is applicable, then: $(\text{Amphibian Breeding Potential-Predator Fish } \{43\}) * \{[(\text{Amphibian Overwintering Habitat } \{44\} + 2 * \text{Upland Buffer Width } (23)_{\text{Wildlife}} + \text{Dominant Upland Land Use } \{14\} + \text{Barriers } \{41\} + \text{Stormwater Input } \{20_{\text{reverse}}\}) / 6\}$

Amphibian Habitat Variables

| MnRAM # | Excel # | Variable Description | Type of Interaction |
|---------|---------|---|---------------------|
| 42 | D54 | Amphibian breeding potential—hydroperiod | Controlling |
| 43 | D55 | Amphibian breeding potential—fish presence | Controlling |
| 44 | E56 | Amphibian overwintering habitat | Compensatory |
| 23 | I27 | Upland buffer width | Compensatory |
| 41 | E53 | Wildlife barriers | Compensatory |
| 14 | E18 | Dominant upland land use | Compensatory |
| 20 | F24 | Stormwater runoff pretreatment & detention—RR | Compensatory |

6.10 AESTHETICS/RECREATION/EDUCATION/CULTURAL/SCIENCE

The aesthetics/recreation/education/cultural and science function and value of each wetland is evaluated based on the wetland's visibility, accessibility, evidence of recreational uses, evidence of human influences (e.g. noise and air pollution) and any known educational or cultural purposes. Accessibility of the wetland is key to its aesthetic or educational appreciation. While dependent on accessibility, a wetland's functional level could be evaluated by the view it provides observers. Distinct contrast

²⁷ Knutson et al., 2000

²⁸ Richter and Azous, 1995

²⁹ Hall and Cuthbert, 2000

³⁰ Knutson, et al., 1999; Findlay and Bourdages, 2000; Semlitsch, 2000.

between the wetland and surrounding upland may increase its perceived importance. Also, diversity of wetland types or vegetation communities may increase its functional level as compared to monotypic open water or vegetation. Excess negative human influence on the wetland is counted double in the formula.

All questions contribute equally to the overall index.

| MnRAM # | Excel # | Variable Description | Type of Interaction |
|---------|---------|------------------------------------|---------------------|
| 48 | E60 | Rare educational opportunity | Controlling |
| 49 | E61 | Wetland visibility | Compensatory |
| 50 | E62 | Proximity to population | Compensatory |
| 51 | E63 | Public ownership | Compensatory |
| 52 | E64 | Public access | Compensatory |
| 53 | E65 | Human influence—wetland | Compensatory |
| 54 | E66 | Human influence—viewshed | Compensatory |
| 55 | E67 | Spatial buffer | Compensatory |
| 56 | E68 | Recreational activities in wetland | Compensatory |

Aesthetics/Recreation/Education/Cultural/Science Functional Index Computations:

If Special Features c, h, or u is checked³¹, or

If 48=1, then Index = Exceptional;

If 53=0.1 (Low), then = $(50+51+52+2*53+54+55+56)/8$

If 53>0.1, then = $(49+50+51+52+53+54+55+56)/8$

Entire Formula

$(\text{Wetland Visibility } \{49\} + \text{Proximity to Population } \{50\} + \text{Public Ownership } \{51\} + \text{Public Access } \{52\} + \text{Human Influence - Wetland } \{53\} + \text{Human Influence - Viewshed } \{54\} + \text{Spatial Buffer } \{55\} + \text{Recreational Activities in Wetland } \{56\})/8$

6.11 COMMERCIAL USES

This question considers the nature of any commercially-valuable use of the wetland and requires the assessor to consider how such use may be a detriment to the sustainability of the wetland. Some row crops can be planted in Type 1 wetlands after spring flooding has ceased and still have adequate time to grow to maturity. This non-wetland-dependent agricultural use of wetlands may include hay, pasture/grazing, or row crops such as soybeans or corn. Wetland-dependent crops include wild rice and cranberries, which rely on the wetland hydrology for part of their life cycle.

³¹ c = Designated scientific and natural area; h = Archeologic or historic site designated by the State Historic Preservation Office; u = State or Federal designated wilderness area.

Sustainable uses of the wetland would not require modifying a natural wetland. Products in this category would include collection of botanical products, wet native grass seed, floral decorations, wild rice, black spruce, white cedar, and tamarack. Sustainable uses may require modification of the natural hydrology, such as for wetland-dependent crops (rice, cranberries). Haying and grazing can be less intrusive agricultural activities utilized more or less casually when hydrologic conditions permit; light pasture and occasional haying would be considered more or less sustainable. Like peat-mining, cropping is an unsustainable use of the wetland as it results in severe alterations of wetland characteristics (soil, vegetation, hydrology).

| MnRAM # | Excel # | Variable Description | Type of Interaction |
|---------|---------|-----------------------------------|---------------------|
| 57 | E69 | Commercial crop—hydrologic impact | Controlling |

Commercial Uses Functional Index = 57

6.12 GROUND-WATER INTERACTION

The ground water interaction function is the most difficult to assess. Here the most likely type of ground water interaction is determined, i.e. recharge or discharge, or a combination. In many cases, a wetland will exhibit both recharge and discharge characteristics, however one is usually more dominant. Several wetland and watershed characteristics are evaluated to determine the likely interaction including: wetland soil type, upland land use, upland soil types and wetland size, wetland hydroperiod, wetland outlet characteristics, and topographic relief.

The purpose of this function is strictly to determine the likelihood of the appropriate ground-water interaction based on observable characteristics of the wetland and watershed. The significance of ground water as a component of the wetland water budget is the most difficult functional characteristic to determine without large quantities of detailed hydrologic and geologic information. The following methodology takes the most easily observable and distinct measures of recharge/discharge relationships from the *Wetland Evaluation Technique*³² and the *Hydrogeomorphic Assessment Methodology*³³. In many wetlands, surface water and ground water both make significant contributions to the water budget, but occasionally recharge or discharge is dominant. The goal here is to identify the dominant ground-water interaction (if there is one) to help guide future management and provide an indication when additional information may be warranted.

³² Adamus, et al., 1987

³³ Magee and Hollands, 1998

- If 5 or 6 of questions 58-63 are answered the same, this indicates a strong likelihood that the most frequently stated interaction exerts the primary influence on the wetland.
- If 3-4 questions are answered the same, then the wetland is likely influenced by a combination of both recharge and discharge interactions (i.e. both types of ground water interaction are likely to be present at some point during most years).

58. Wetland Soils – from HGM system functional assessments and Novitzki

59. Subwatershed Land Use/Imperviousness – taken from WET Volume I

60. Wetland Size and Upland Soils – taken from WET Volume I and HGM

61. Wetland Hydrologic Regime– taken from WET Volume I and HGM

62. Inlet/Outlet Configuration – taken from WET Volume I and HGM

63. Upland Topographic Relief – taken from WET Volume I

Special Concerns for Recharge Wetlands

Wherever ground water recharge is indicated as the **primary** interaction and the wetland lies within a sensitive ground water area (**Special Feature Question q**), a contribution area to a public water supply, or a wellhead protection area (**Special Feature Question r**), it should be recorded as Exceptional for the ground water/wetland function.

6.13 WETLAND RESTORATION POTENTIAL

The potential for wetland restoration is determined based on the ease with which the wetland could be restored, the number of landowners within the historic wetland basin, the size of the potential restoration area, the potential for establishing buffer areas or water quality ponding, and the extent and type of hydrologic alteration. Each variable uses the High, Medium, Low rating rather than raw numbers—see MnRAM for individual ranges.

| MnRAM # | Excel # | Variable Description | Type of Interaction |
|---------|---------|---|---------------------|
| 64 | D79 | Wetland Restoration Potential | Controlling |
| 65 | F80 | Number of Landowners Affected | Contributing |
| 21 | E25 | Subwatershed Wetland Density | Contributing |
| 66b | F82 | Total Wetland Restored Size (Potential) | Contributing |
| 66c | F83 | Calculated potential new wetland area | Contributing |
| 67 | F84 | Potential Buffer Width | Contributing |
| 68 | F85 | Likelihood of Restoration Success | Contributing |

If 64="Yes", then Wetland Restoration Potential = (65+21+66b+66c+67+68)/6,

Otherwise, if 64="No" then "N/A"

Entire Formula

(Landowners Affected by Restoration (65)+Subwatershed Wetland Density (21)+Wetland Restoration Size (66b)+Proportion of Wetland Drained (66c)+Potential Buffer Width (67)+Likelihood of Restoration Success (68))/6

6.14 WETLAND SENSITIVITY TO STORMWATER INPUT AND URBAN DEVELOPMENT

The sensitivity of the wetland to stormwater and urban development is determined based on guidance within the *Storm-Water and Wetlands: Planning and Evaluation Guidelines for Addressing Potential Impacts of Urban Storm-Water and Snow-Melt Runoff on Wetlands*, State of Minnesota Storm-Water Advisory Group, June, 1997.

Use habitat proportions from Vegetative Integrity section and enter into a formula to compute answer according to the following criteria³⁴.

- Exceptional = Sedge meadows, open and coniferous bogs, calcareous fens, low prairies, wet to wet-mesic prairies, coniferous swamps, lowland hardwood swamps, or seasonally flooded basins.
- A = Shrub-carrs, alder thickets, diverse fresh wet meadows dominated by native species, diverse shallow and deep marshes, and diverse shallow, open water communities.
- B = Floodplain forests, fresh wet meadows dominated by reed canary grass, shallow and deep marshes dominated by cattail, reed canary grass, giant reed or purple loosestrife, and shallow, open water communities with low to moderate vegetative diversity.
- C = Gravel pits, cultivated hydric soils, or dredge/fill disposal sites.

6.15 ADDITIONAL STORMWATER TREATMENT NEEDS

This rates the sustainability of the wetland with regard to stormwater discharges to the wetland. The need for additional stormwater treatment prior to discharge to the wetland is rated based on the overall rating for Maintenance of Wetland Water Quality. If a wetland is severely degraded by stormwater inputs, the rating will be low, since a diverse, high quality wetland will not be sustainable.

Use functional rating for Maintenance of Wetland Water Quality (MWWQ) as follows (this index is rated strictly from the measure of the water quality in the wetland and the sustainability, i.e. if the water quality in the wetland is low, additional stormwater treatment is needed to protect the wetland and the rating is low):

Use Value for Maintenance of Wetland Water Quality Index (D76, Excel spreadsheet) and apply to criteria below.

- A = Maintenance of Wetland Water Quality Index > 0.66 (no additional treatment needed)
- B = $0.33 < \text{Maintenance of Wetland Water Quality Index} \leq 0.66$ (sediment removal needed)

³⁴ Taken directly from State of Minnesota Storm-Water Advisory Group, 1997.

C = Maintenance of Wetland Water Quality Index < 0.33 (sediment and nutrient removal needed)

- Knutson, M.G., J.R. Sauer, D.A. Olsen, M.J. Mossman, L.M. Hemesath and M.J. Lannoo. 1999. *Effects of landscape composition and wetland fragmentation on frog and toad abundance and species richness in Iowa and Wisconsin, U.S.A.* Conservation Biology 13:1437-1446.
- Knutson, M.G., J.R. Sauer, D.A. Olsen, M.J. Mossman, L.M. Hemesath and M.J. Lannoo. 2000. *Landscape associations of frog and toad species in Iowa and Wisconsin, U.S.A.* Jour. Iowa Acad. Sci. 107(3):134-145.
- Knutson, P. L., J. C. Ford, and M. R. Inskeep. 1981. *National Survey of Planted Salt Marshes (vegetative stabilization and wave stress).* Wetlands 3:129-153.
- Kuchler, A.W. 1967. Vegetation Mapping. The Ronald Press, New York, New York.
- Lannoo, M.J. 1998. *Amphibian conservation and wetland management in the upper midwest: a catch 22 for the cricket frog.* In Status and Conservation of Midwestern amphibians. M.J. Lannoo, ed. University of Iowa Press, Iowa City, Iowa. Pages 331-339.
- Lee, L.C. and Mark M. Brinson, William J. Kleindl, P. Michael Whited, Michael Gilbert, Wade L. Nutter, Dennis F. Whigham, Dave DeWald. 1997. *Revised Operational Draft Guidebook for the Hydrogeomorphic Assessment of Temporary and Seasonal Prairie Pothole Wetlands.* Seattle, WA. pp. 116+app.
- Lehtinen, R.M., S.M. Galatowitsch and J.R. Tester. 1999. *Consequences of habitat loss and fragmentation for wetland amphibian assemblages.* Wetlands 19:1-12.
- Magee, D.W. and Garrett G. Hollands. 1998. *A Rapid Procedure for Assessing Wetland Functional Capacity.* Association of State Wetland Managers. 190 pp.
- Michigan DNR. 1981. *Manual for Wetland Evaluation Techniques.* Michigan Department of Natural Resources, Lansing, MI.
- Mitch & Gosselink 2000. Wetlands 3rd Edition.
- Ogawa, H. and J. W. Male. 1983. *The Flood Mitigation Potential of Inland Wetlands.* Water Resources Research Center. University of Massachusetts.
- Oldfield, B. and J.J. Moriarty. 1994. *Amphibians and reptiles native to Minnesota.* University of Minnesota Press, Minneapolis, Minnesota.
- Reed, P.B. 1988 (and 1993 supplement). National List of Plant Species that Occur in Wetlands, U. S. Fish and Wildlife Service National Wetland Inventory, St. Petersburg, FL.
- Rheinhardt, R., Mark M. Brinson, N. Eric Fleming, J. Glenn Sandifer, Jr., 1997. *Deciduous Wetland Flats Interim HGM Model.* 50 pp.
- Richter, Klaus O. and Amanda L. Azous. 1995. Amphibian occurrence and wetland characteristics in the Puget Sound Basin. Wetlands 15:305-312.
- Seaberg, J.K. and Douglas D. Hansen. 2000. *Metropolitan Area Groundwater Model Project Summary: Northwest Province, Layers 1, 2, and 3 Model, Version 1.00.* Minnesota Pollution Control Agency.
- Semlitsch, R.D. 2000. Principles for management of aquatic-breeding amphibians. J. Wildlife Mgmt. 64:615-631.
- Shaw, S. and C.G. Fredine. 1956. *Wetlands of the United States Circular 39.* U.S. Fish and Wildlife

Service. U.S. Government Printing Office, Washington, D.C.

Smith, S.G. 1986. *The cattails (Typha): Interspecific ecological differences and problems of identification*. Proceedings of the 5th Annual Conference of the NALMS. Geneva, Illinois, 357-362.

Snodgrass, J.W., M.J. Komoroski, A.L. Bryan, Jr., and J. Burger, Jr. 2000. *Relationships among isolated wetland size, hydroperiod, and amphibian species richness: implications for wetland regulations*. Conservation Biology 14:414-419.

State of Minnesota Storm-Water Advisory Group. 1997. *Storm-Water and Wetlands: Planning and Evaluation Guidelines for Addressing Potential Impacts of Urban Storm-Water and Snow-Melt Runoff on Wetlands*.

Thill, D., and M. Jacobson. 2000. *The Minnehaha Creek Routine Assessment Method*, in conjunction with the Minnehaha Creek Watershed District.

Environmental Laboratory. 1987. Corps of Engineers Wetlands Delineation Manual. Technical Report Y-87-1, Final Report.

Verry, E.S. 1988. *The Hydrology of Wetlands and Man's Influence on it*. International Symposium on the Hydrology of Wetlands in Temperate and Cold Regions. Joensuu, Finland.

Wells, J. Mike Mueller, John Parker, Bruce Gerbig, S. Jatnieks-Straumanis, Bruce Wilson, John Kittelson, Paul Richert, Teri Sardinas, Scott LaChance, and Larry Smith, 1988. *The Minnesota Wetland Evaluation Methodology for the North Central United States*. Corps of Engineers in conjunction with the Minnesota Environmental Quality Board Wetland Evaluation Methodology Task Force, John R. Wells, Chairman.

Whited, D., Galatowitsch, S., Tester, J.R., Schik, K., Lehtinen, R., Husveth, J. 2000. Landscape and Urban Planning, 49 (1-2): 49-65.

Williams, R.E. 1968. *Flow of groundwater adjacent to a small, closed basin in glacial till*. Water Resources Research 4:777-783.

BMP References:

Board of Water and Soil Resources, Association of Metropolitan Soil and Water Conservation Districts. 1989. Minnesota Construction Site Erosion and Sediment Control Planning Handbook. St. Paul.

Minnesota Pollution Control Agency. 2000. Protecting Water Quality in Urban Areas: Best Management Practices for Dealing with Storm Water Runoff from Urban, Suburban and Developing Areas of Minnesota. Minneapolis, March, 2000.

Barr Engineering Company. 2001. Minnesota Urban Small Sites BMP Manual: Stormwater Best Management Practices for Cold Climates. Metropolitan Council Environmental Services, St. Paul, MN.

Appendix G

Vegetation Shoreline Buffer Brochure Examples



Sullivan Shoreline Planting



Project: A 375 square foot shoreline planting along Crystal Lake, covering approximately 50 linear feet of shoreline. Erosion control blanket, native shrubs, and deep-rooted native plant plugs were used to stabilize the existing slope.

Costs: Project material costs were estimated at \$935.

Funding: Dakota County SWCD provided technical assistance and Blue Thumb Grant in the amount of \$100. The City of Burnsville provided Neighborhood Water Resources Enhancement Grant.



Location:

Burnsville
Minnesota



Practice:

Shoreline Planting

Shoreline Benefits:

Reduced erosion and sediment into the receiving waterbody

Improved aesthetics

Improved water quality

Slope stabilization

Partners:

Black Dog Watershed Management Organization

City of Burnsville

Watershed:

Minnesota River

Construction:

July
2009

Dakota County Soil and Water Conservation District

4100 220th St. W., Suite 102, Farmington, MN 55024 651-480-7777 www.dakotaswcd.org

Revised 8/4/09

Fay Shoreline



Project: A 600 square foot shoreline planting.

Costs: Project material costs were estimated at \$1,847.

Funding: Dakota County SWCD provided technical assistance and Blue Thumb Grant in the amount of \$250.



Practice:

Shoreline planting and Native garden

Benefits:

Runoff volume reduction

Improved aesthetics

Improved water quality

Opportunity for public education and outreach

Wildlife habitat

Slope stabilization

Partner:

Black Dog Watershed Management Organization

Watershed:

Minnesota River


Construction:

2013



Location:

Burnsville
Minnesota



Dakota County Soil and Water Conservation District

4100 220th St. W., Suite 102, Farmington, MN 55024 651-480-7777 www.dakotaswcd.org

Revised 9/18/2013

COADY SHORELINE PLANTING



BEFORE

Shoreline planting is the use of native vegetation to protect a shoreline from existing or potential erosion



AFTER

PROJECT: Installation of a 1000 square foot shoreline planting


COST: Project materials cost estimated at \$3,192

FUNDING: Landowners receive a \$250 Blue Thumb grant as well as technical assistance provided by the Dakota County SWCD



LOCATION:

Burnsville MN
Bluebill Bay Road



PRACTICE:

- Shoreline Planting

BENEFITS:

- Shoreline stabilization and erosion reduction
- Improved water quality
- Improved wildlife habitat
- Opportunity for public education and outreach
- Improved aesthetics

PARTNERS:

- Black Dog Watershed Management Organization

WATERBODY:

- Crystal Lake

WATERSHED:

- Minnesota River

INSTALLATION:

- Summer 2014

Appendix H

Buckthorn Management Guidelines

Buckthorn Management Guidelines

Goal: Restore native plant communities in designated natural areas and other park locations by controlling and removing non-native invasive species.

Buckthorn belongs to the *Rhamnaceae* family. It is native to Europe and Asia, first appearing in the U.S. in the late 1700s. Buckthorn quickly naturalized in the woodlands of the northeastern states. Today buckthorn flourishes in the understory of Minnesota woodlands and in brushy thickets along roadsides and fields. It has become a major plant pest in natural woodlands and wetlands.

Buckthorn can grow to 15-20 feet and has dark green elliptical or oval leaves. In the fall its leaves hang on late into the season and without much color change. It starts easily from seed and will tolerate almost any soil condition or location. In partial shade it will outstretch its neighbors toward the light.

Buckthorn removal is recommended for those areas where the native plant community has been displaced by buckthorn species and where there is a high likelihood that the native plant community can be enhanced and restored.

Restoration of the native communities is the overall intent of non-native eradication efforts.

Volunteer Considerations

Volunteers must be trained in species identification, removal techniques and other aspects related to the eradication/restoration efforts.

Identification of buckthorn by volunteers is best performed during the month of October.

Process

Buckthorn removal is a long-term process requiring several steps over a three- to four-year period. Pulling seedlings, cutting and removing mature plants, chemically treating stumps and replanting the site with native species are critical to the long-term success of restoration efforts.

Staff are responsible for cutting mature plants and chemically treating the stumps in areas designated for restoration. A 20%-25% solution of glyphosate (Roundup) with a dye is used to paint, chemically treat, and mark the stumps.

Volunteer procedures

1. Hand pulling allowed by volunteers with training or under the supervision of a "trained" volunteer supervisor.
2. Use of loppers allowed by volunteers.
3. No use of power tools or chemicals by volunteers; chemicals and power tool use only by staff or contractor.
4. Volunteers must sign waiver form.

Recommended chronology of restoration activities with volunteers

Year one

- Seedlings cut or pulled (September-November)
- Mature trees cut by staff and/or volunteers in late fall (October-December)
- Stumps or stems chemically treated by staff immediately after cutting
- Removal of brush to a chipping location (or pile on site for burning)

Year two

- Remove seedlings by hand pulling or cutting and treating (June-November)
- Follow-up cutting by staff and/or volunteers in late fall (October-December) and chemically treat stump and stems.

Year three

- Seedling removal by hand pulling or cutting and treating as necessary
- Plant native understory shrubs, trees, ferns, wildflowers and grasses to approximate prior native plant community.

Year four

- Continued monitoring and buckthorn seedling removal

Other removal techniques

Mechanical

- Prescribed fire for seedlings; prescribed burns in early spring and fall annually or biannually to control buckthorn may have to be continued for several years

Chemical

- Cut-stump and stem treatment with glyphosate; 20%-25% active ingredient cut-stump; or basal bark spray treatment around the stem with 25-50% a.i. triclopyr (Garlon) – consideration of worker safety issues will dictate chemical selection. Glyphosate products registered for wetland/aquatic use should be used on water bodies and wetlands. Sponge applicators can help prevent chemical spill or spread to workers.
- Fosamine, a non-selective bud inhibitor for woody species, can be applied as a basal bark treatment in the fall at 3% a.i. concentration in winter

Another technique is goat rental.

The method of buckthorn control should be selected based on the site, safety concerns, and opportunities for continued vegetation management.

Other Sources for Guidance

University of Minnesota:

<http://www.extension.umn.edu/environment/agroforestry/woody-vegetation-control.html>

University of Wisconsin:

http://mipncontroldatabase.wisc.edu/search?name=common_buckthorn&habitat=7&season=7

Minnesota Department of Natural Resources:

<http://www.dnr.state.mn.us/invasives/terrestrialplants/woody/buckthorn/control.html>

U.S. Department of Agriculture Natural Resources Conservation Service:

<https://efotg.sc.egov.usda.gov/references/public/MN/797Buckthorn.pdf>

See Buckthorn Control Quick Guide for a summary of control techniques.

Appendix I


Lac Lavon Prairie Restoration Area

Provided by the City of Burnsville



**Lac Lavon
Prairie Planting**

Project:



Date:
Notes: