

Appendix C

2003-2020 Habitat Assessment Monitoring Results

**Appendix C: 2003-2009 Habitat Assessment Monitoring Results
Black Dog Watershed Management Organization**

Water Body	Monitoring Year	Approximate Proportion of the Water Body Which is Deep Water Habitat (~ > 20 ft. depth)	Vegetation Quality - Wet Areas											Vegetation Quality - Upland							Erosion/Sedimentation			
			Submergent Zone Sampling					Vegetated Emergent Zone Sampling						Upland Buffer Sampling										
			Overall Submergent Vegetative Quality ¹	Approximate Proportion of Water Body Typically Dominated By Submergent Vegetation (~ 2 - 20 ft. depth)	Average Native Plant Occurrence or Density Rating ^{2,3}	Total Number of Native Species ⁵	Exotic Species		Emergent Zone Vegetative Quality ⁶	Approximate Proportion of Emergent Zone (0 - 2 ft. depth) Within The Water Body	Approximate Total Percent Vegetative Cover Within The Entire Emergent Zone ⁷	Total Number of Native Wetland Plant Species ⁸	Exotic Species		Overall Upland Buffer Quality ¹⁰	Unmanicured Buffer Width ¹¹	Estimated Total Vegetative Cover (Percent Range) ¹²	Total Number of Native Plant Species ¹³	Buffer Continuity (Percent Surrounding Water Body) ¹⁴	Exotic Species		Shoreline Erosion (Percent of Shoreline) ¹⁶	Sediment Deltas (Yes/No)	
Crystal	2003	15%	Moderate	80%	1.5	15	2	1.1	1.1	Moderate	5%	26-50%	18	4	26-50%	Moderate	<10 ft.	>95%	16	26-50%	2	15-40%	0-10%	No
	2004		Excellent		1.2	14	2	1.1	2.9	Excellent		26-50%	16	6	26-50%	Moderate	<10 ft.	>95%	16	26-50%	4	15-40%	0-10%	No
	2005		Moderate		1.2	13	2	1.1	2.7	Excellent		26-50%	16	6	26-50%	Moderate	<10 ft.	>95%	17	26-50%	3	15-40%	0-10%	No
	2006		Excellent		1.0	17	2	1.5	3.2	Excellent		26-50%	18	8	26-50%	Moderate	<10 ft.	>95%	17	26-50%	3	15-40%	0-10%	No
	2007		Excellent		1.5	16	2	1.6	3.4	Excellent		26-50%	22	10	26-50%	Moderate	<10 ft.	>95%	15	26-50%	5	15-40%	0-10%	No
	2008		Moderate		1.3	15	2	1.6	2.5	Excellent		26-50%	21	12	26-50%	Moderate	<10 ft.	>95%	15	26-50%	5	15-40%	0-10%	No
	2009		Moderate		1.3	14	2	1.6	2.8	Excellent		26-50%	20	11	26-50%	Moderate	<10 ft.	>95%	15	26-50%	7	15-40%	0-10%	No
Keller	2003	0%	Moderate	90%	1.9	4	1	3.2	3.2	Poor	10%	51-75%	5	2	51-75%	Moderate	<10 ft.	>95%	7	76-100%	6	>40%	0-10%	No
	2004		Moderate		1.7	5	1	1.8	2.5	Moderate		51-75%	6	2	51-75%	Moderate	<10 ft.	>95%	7	76-100%	6	>40%	0-10%	No
	2005		Moderate		1.3	5	2	1.0	1.1	Moderate		51-75%	7	2	26-50%	Moderate	<10 ft.	>95%	8	76-100%	7	>40%	0-10%	No
	2006		Moderate		2.0	5	2	1.8	2.5	Moderate		51-75%	8	2	26-50%	Moderate	<10 ft.	>95%	8	76-100%	8	>40%	0-10%	No
	2007		Moderate		2.1	3	2	2.4	3.8	Moderate		51-75%	9	3	26-50%	Moderate	<10 ft.	>95%	5	76-100%	9	>40%	0-10%	No
	2008		Moderate		2.2	3	2	2.2	2.9	Moderate		51-75%	9	3	26-50%	Moderate	<10 ft.	>95%	6	76-100%	12	>40%	0-10%	No
	2009		Poor		3.0	2	2	2.7	3.3	Moderate		51-75%	9	4	26-50%	Moderate	<10 ft.	>95%	4	76-100%	11	>40%	0-10%	No
Kingsley	2003	0%	Moderate	95%	2.7	7	0	0.0	0.0	Excellent	5%	51-75%	11	2	0-25%	Excellent	25-50 ft.	>95%	15	51-75%	4	15-40%	0-10%	No
	2004		Moderate		2.7	7	0	0.0	0.0	Excellent		51-75%	11	2	0-25%	Excellent	25-50 ft.	>95%	15	51-75%	4	15-40%	0-10%	No
	2005		Moderate		2.6	7	1	1.0	1.0	Excellent		51-75%	15	6	0-25%	Excellent	25-50 ft.	>95%	19	76-100%	2	15-40%	0-10%	No
	2006		Excellent		1.8 ¹⁷	13 ¹⁸	1	1.0	1.0	Excellent		51-75%	15	6	0-25%	Excellent	25-50 ft.	>95%	19	76-100%	3	15-40%	0-10%	No
	2007		Excellent		1.6	13	1	1.0	1.0	Excellent		51-75%	19	6	0-25%	Excellent	25-50 ft.	>95%	21	76-100%	4	15-40%	0-10%	No
	2008		Moderate		2.9	5	0	0.0	0.0	Excellent		51-75%	18	5	0-25%	Excellent	25-50 ft.	>95%	25	76-100%	4	15-40%	0-10%	No
	2009		Excellent		2.0	11	1	1.0	1.0	Excellent		51-75%	16	5	0-25%	Excellent	25-50 ft.	>95%	23	76-100%	5	15-40%	0-10%	No
Lac Lavon	2003	25%	Poor	70%	2.0	7	1	1.0	1.0	Poor	5%	0-25%	14	5	0-25%	Poor	<10 ft.	<75%	12	0-25%	17	>40%	0-10%	No
	2004		Moderate		0.9	9	2	1.6	1.9	Moderate		0-25%	15	5	0-25%	Poor	<10 ft.	<75%	12	0-25%	17	>40%	0-10%	No
	2005		Moderate		2.3	5	1	2.0	2.0	Excellent		0-25%	20	10	0-25%	Poor	<10 ft.	<75%	12	0-25%	16	>40%	0-10%	No
	2006		Moderate		1.6	10 ¹⁹	2	2.5	4.0	Excellent		0-25%	16	13	0-25%	Poor	<10 ft.	<75%	11	0-25%	19	>40%	0-10%	No
	2007		Excellent		1.8	10 ²⁰	3	1.8	4.0	Excellent		0-25%	16	12	0-25%	Poor	<10 ft.	<75%	12	0-25%	18	>40%	0-10%	No
	2008		Poor		1.0	5	2	1.0	1.0	Moderate		0-25%	14	9	0-25%	Poor	<10 ft.	<75%	9	0-25%	13	>40%	0-10%	No
	2009		Moderate		1.6	10	2	2.5	4.0	Moderate		0-25%	13	8	0-25%	Poor	<10 ft.	<75%	9	0-25%	11	>40%	0-10%	No
Orchard	2003	20%	Poor	75%	1.2	13	1	2.3	3.4	Moderate	5%	26-50%	16	5	26-50%	Moderate	<10 ft.	>95%	5	26-50%	5	>40%	0-10%	No
	2004		Moderate		1.2	13	1	2.3	2.3	Excellent		26-50%	17	5	26-50%	Moderate	<10 ft.	>95%	5	26-50%	5	>40%	0-10%	No
	2005		Moderate		1.3	14	1	1.8	2.6	Moderate		26-50%	14	6	26-50%	Moderate	<10 ft.	>95%	5	26-50%	5	>40%	0-10%	No
	2006		Moderate		1.2	13	1	1.7	3.4	Excellent		26-50%	18	9	26-50%	Moderate	<10 ft.	>95%	5	26-50%	5	>40%	0-10%	No
	2007		Moderate		1.3	11	1	1.9	3.3	Excellent		26-50%	18	9	26-50%	Moderate	<10 ft.	>95%	3	26-50%	5	>40%	0-10%	No
	2008		Moderate		1.3	14	1	1.6	2.8	Excellent		26-50%	16	8	26-50%	Moderate	<10 ft.	>95%	3	26-50%	7	>40%	0-10%	No
	2009		Moderate		1.6	11	1	1.7	2.5	Excellent		26-50%	16	8	26-50%	Moderate	<10 ft.	>95%	3	26-50%	6	>40%	0-10%	No
Sunset Pond	2003	0%	Moderate	75%	3.0	11	1	1.0	1.0	Poor	25%	76 - 100%	5	5	76-100%	Moderate	10-25 ft.	75-95%	10	51-75%	15	15-40%	0-10%	Yes
	2004		Excellent		2.2	11	0	0.0	0.0	Poor		76 - 100%	4	3	76-100%	Moderate	10-25 ft.	75-95%	10	51-75%	18	15-40%	0-10%	Yes
	2005		Excellent		2.1	10	1	1.0	1.0	Poor		76 - 100%	6	4	76-100%	Moderate	10-25 ft.	75-95%	9	76-100%	20	>40%	0-10%	Yes
	2006		Moderate		2.6	11	1	1.0	1.0	Poor		76 - 100%	7	4	76-100%	Moderate	10-25 ft.	75-95%	9	76-100%	19	>40%	0-10%	No
	2007		Excellent		1.9	12	1	1.0	1.0	Moderate		76-100%	11	6	76-100%	Moderate	10-25 ft.	75-95%	8	76-100%	19	>40%	0-10%	No
	2008		Excellent		1.8	10	1	2.0	3.0	Poor		76-100%	10	5	76-100%	Moderate	10-25 ft.	75-95%	5	76-100%	15	>40%	0-10%	No
	2009		Moderate		2.2	11	1	3.0	3.0	Poor		76-100%	10	5	76-100%	Moderate	10-25 ft.	75-95%	6	76-100%	17	>40%	0-10%	Yes

**Appendix C: 2003-2009 Habitat Assessment Monitoring Results
Black Dog Watershed Management Organization**

The following footnotes pertain to 2003-2009 data.

¹Overall Submergent Vegetative Quality rating is the average of the exotic species density, macrophyte density, and total number of native: >0.66 = Excellent, 0.33-0.66 = Moderate, <0.33 = Poor.

Overall Submergent Vegetative Quality	Avg. Exotic Species Density	Exotic Species Density/ Occurrence Rating Score	Avg. Macrophyte Density	Avg. Macrophyte Density Rating Score	Total Number of Native Species In Submergent Zone	Species Richness Rating	Total Overall Diversity Score
Poor	>2.0	0.1	0.0 - 1.0 and >3.0	0.1	<9	0.1	< 0.33
Moderate	>0 - 2.0	0.5	1.0 - 1.5 and > 2.5 to 3.0	0.5	9-14	.25-.75	0.33 - 0.66
Excellent	0	1.0	1.5 to 2.5	1.0	>14	1.0	> 0.66

²Plant occurrence ratings are a relative measure of the amount of native submergent vegetation with a scale from 1 to 5; 1 = lowest density (present on only 1 of 4 casts), 5 = highest density (hook full of vegetation on 4 of 4 casts).

³Density data for Crystal, Keller, and Orchard Lakes were collected by Blue Water Science. Numerous sample plots were conducted over the entire water body. A density scale of 1 to 4 was utilized (max = 4) by estimating the amount of vegetation obtained by rake casts and also transforming visual observations.

⁴Maximum exotic plant occurrence ratings represent the worst case scenario of curlyleaf pondweed density early in the growing season and/or Eurasian watermilfoil when it is most prolific later in the growing season.

⁵The Total Number of Native Species within the submergent zone for Crystal, Keller and Orchard Lakes is based on a detailed survey conducted by Blue Water Science; and for Kingsley Lake, Lac Lavon, and Sunset Pond, based on a survey by Barr Engineering and volunteers. The survey of the 3 water bodies conducted by Blue Water Science involved

the sampling of numerous sample plots or stations. The survey for Lac Lavon, Kingsley, and Sunset Pond is based on 3 sampling locations and a visual survey during travels on the water body: <7 = Poor, 7-14 = Moderate, >14 = Excellent.

⁶Emergent Zone Vegetative Quality is the average of the following parameters within the emergent zone: the approximate total percent coverage, the total number of native wetland species, and the percent coverage of exotic species: >0.66 = Excellent, 0.33-0.66 = Moderate, <0.33 = Poor.

Emergent Zone Vegetative Quality	Percent Cover	Percent Cover Rating Score	Total Number of Native Wetland Plants	Number of Native Wetland Plants Rating Score	Percent Cover of Exotics	Percent Cover of Exotics Rating Score	Overall Emergent Zone Quality Score
Poor	0-25%	0.1	<0= 5	0.1	76-100%	0.1	< 0.33
Moderate	76-100% or 26-50%	0.5	>5 - 15	0.66 - 0.33	26-75%	.33-.66	0.33 - 0.66
Excellent	51-75%	1.0	> 15	1.0	0-25%	1.0	> 0.66

⁷Approximate Total Percent Vegetative Cover Within the Entire Emergent Zone (0-2 ft. depth) is estimated based on the three sampling locations and a visual survey during travels around the water body. Estimates are broken into four categories: 0-25%=Poor, 26-50%=Moderate, 51-75%=Excellent, 76-100%=Moderate.

⁸The Total Number of Native Wetland Plant Species within the emergent zone is based on 3 sampling locations and a visual survey during travels on the water body: 0-5 = Poor, 6-15 = Moderate, >15 = Excellent.

⁹Total Exotic Emergent Percent Coverage, out of the entire emergent zone area, is estimated based on the three sampling locations and a visual survey during travels around the water body. Estimates are broken into four categories: 0-25%=Excellen(1.0), 26-50%=Moderate(0.5), 51-75%=Poor(0.0), 76-100%=Poor(0.1)

¹⁰Overall Upland Buffer Quality is determined based on the average of the four upland buffer quality parameters, with the exception of the number of exotic species present and the number of native plant species: >0.66 = Excellent, 0.33-0.66 = Moderate, <0.66 = Poor.

Overall Upland Buffer Quality	Percent Cover	Percent Cover Rating Score	Exotics Percent Cover Range	Exotics Percent Cover Rating Score	Buffer Width Range	Buffer Width Rating Score	Buffer Continuity Percent Range	Buffer Continuity Rating Score	Overall Upland Buffer QualityScore
Poor	<75%	0.1	>40%	0.1	<10 ft.	0.1	0-25%	0.1	< 0.33
Moderate	75-95%	0.5	15-40%	0.5	10-50 ft.	0.4 - 0.7	25-75%	.4-.7	0.33 - 0.66
Excellent	>95%	1.0	<15%	1.0	>50 ft.	1.0	76-100%	1.0	> 0.66

¹¹Unmanicured (upland) Buffer Width is divided into four categories: Excellent(1.0) = >50 ft, High(0.7) = 25-50 ft, Moderate(0.4) = 10-25 ft, and Low(0.1) = <10 ft.

¹²Estimated Total Vegetative Cover (Percent Range) for upland buffer is the proportion of the ground covered by vegetation within 50 feet of the wetland/upland transition zone. The percent cover is divided into three categories: Excellent(1.0) = >95%, Moderate(0.5) = 75 - 95%, and Poor(0.1) = <75%.

¹³The Total Number of Native Plant Species within the unmanicured upland buffer zone is based on 3 sampling locations and a visual survey.

¹⁴(Upland) Buffer Continuity is a measure of the proportion of the water body surrounded by the unmanicured, native upland buffer. This measure is divided into four categories: Excellent(1.0) = 76 - 100%, High(0.7) = 51 - 75%, Medium(0.4) = 26 - 50%, and Low(0.1) = 0 - 25%.

¹⁵Upland buffer exotic species "Percent of Total Coverage" is the percent cover of exotic species within the unmanicured upland buffer, which is divided into three categories: Excellent(1.0) = <15%, Moderate(0.5) = 15 - 40%, and Poor(0.1) = >40%.

¹⁶The presence of shoreline erosion is determined by the approximate percentage of the shoreline affected and is divided into the following three categories: 0 - 10%, 11 - 25%, 26 - 100%.

¹⁷The 2006 plant occurrence rating is lower (has improved), when compared to past assessment years primarily due to the low occurrence of additional plants found during a more detailed survey of the lake. The more detailed plant survey was conducted to better understand the extent of curlyleaf pondweed.

¹⁸The number of plant species documented in 2006, when compared to past assessment years, increased primarily due to additional plants found during a more detailed survey of the lake. The more detailed plant survey was conducted to better understand the extent of curlyleaf pondweed.

¹⁹The number of native submergent plant species documented in 2006, was incorrectly represented as 11 in the 2006 annual report. The actual number of native submergent plant species documented in 2006 was 10.

²⁰Native plant species were noted by the Minnesota Department of Natural Resouces during an October 25, 2007 macropyte survey and used in the 2007 annual report.

Rating Code: Poor Moderate Excellent

Table 1: Orchard Lake 2012 and 2017 Habitat Assessment Monitoring Results Black Dog Watershed Management Organization

Monitoring Year	Submergent Zone								
	Approximate Proportion of the Water Body Which is Deep Water Habitat (~ > 20 ft. depth)	Overall Submergent Zone Quality ¹	Approximate Proportion of Water Body Typically Dominated By Submergent Vegetation (~ 2 - 20 ft. depth)	Native Species		Mean Coefficient of Conservatism Value	Exotic Species		
				Average Native Plant Density Rating ^{2,3}	Total Number of Native Species ⁵		Total Number of Species	Average Exotic Plant Density Rating ^{2,3}	Maximum Exotic Plant Density Rating ⁴
2012	20%	Moderate	75%	2.0 (Moderate)	13 (High)	5.4 (Moderate)	1	1.7 (Moderate)	3.0 (Poor)
2017	20%	High	75%	1.2 (Excellent)	16 (Excellent)	5.2 (Moderate)	2	1.1 (Moderate)	1.5 (Moderate)

Monitoring Year	Emergent Zone						
	Overall Emergent Zone Quality ⁶	Approximate Proportion of Emergent Zone (0 - 2 ft. depth) Within The Water Body	Approximate Total Percent Vegetative Cover Within The Entire Emergent Zone ⁷	Total Number of Native Wetland Plant Species ⁸	Mean Coefficient of Conservatism Value	Exotic Species	
						Number of Species	Total Exotic Emergent Percent Coverage ⁹
2012	Moderate	5%	26-50% (Moderate)	43 (Excellent)	3.1 (Moderate)	12	51-75% (Moderate)
2017	Moderate	15%	51-75% (High)	50 (Excellent)	2.7 (Poor)	13	51-75% (Moderate)

Monitoring Year	Upland Buffer								Erosion/Sedimentation	
	Overall Upland Buffer Quality ¹⁰	Unmanicured Buffer Width ¹¹	Estimated Total Vegetative Cover (Percent Range) ¹²	Total Number of Native Plant Species ¹³	Mean Coefficient of Conservatism Value	Buffer Continuity (Percent Surrounding Water Body) ¹⁴	Exotic Species		Shoreline Erosion (Percent of Shoreline) ¹⁶	Sediment Deltas (Yes/No)
							Number of Species	Percent of Total Coverage ¹⁵		
2012	Poor	<10 ft. (Poor)	>95% (High)	19 (Moderate)	1.6 (Poor)	0-25% (Poor)	20	>40% (Poor)	0-10%	No
2017	Moderate	<10 ft. (Poor)	>95% (High)	25 (High)	1.9 (Poor)	0-25% (Poor)	21	>40% (Poor)	0-10%	No

Table 1: Orchard Lake 2017 Habitat Assessment Monitoring Results Black Dog Watershed Management Organization

The following changes were made to the 2011 - 2017 monitoring and analysis:

- Monitor one or two water bodies per year. Kingsley Lake in 2011 - Conduct a meandering survey of submergent, emergent, and upland buffer zones rather than monitoring of plot locations. Orchard Lake in 2012, Crystal Lake in 2013, Lac Lavon in 2014, Keller Lake in 2015, Kingsley Lake in 2016, Orchard Lake in 2017 - Conduct a meandering survey of submergent, emergent, and upland buffer zones. In addition, the emergent and upland buffer plot locations were evaluated.
- Changes were made in 2011 through 2017 to the calculations to include floristic quality as part of the assessment. These changes include adding a rating of "High" to the categories to accommodate MPCA ratings for floristic quality. These changes included adding a Rating Code:

Poor	Moderate	High or Excellent
------	----------	-------------------

The following footnotes pertain to 2011 through 2017 data:

¹**Overall Submergent Zone Quality** rating is the average of the rating scores for the following parameters: average exotic plant density, average native plant density, total number of native species, and C-value rating: >0.80 = Excellent, 0.67-0.80 = High, 0.33-0.66 = Moderate, <0.33 = Poor.

Overall Submergent Zone Quality	Avg. Exotic Plant Density	Exotic Plant Density Rating Score	Avg. Native Plant Density	Avg. Native Plant Density Rating Score	Total Number of Native Species In Submergent Zone	Species Richness Rating Score	Mean Coefficient of Conservatism Value (C-Value)	C-Value Rating (using MPCA values, 2007)	Total Overall Submergent Zone Quality Score
Poor	>2.0	0.1	> 1.75	0.1	<7	0.1	0 - <3	0.10	< 0.33
Moderate	>1.0 - 2.0	0.5	1.25 - 1.75	0.5	>7 - <9	0.5	>3 - <6	0.50	0.33 - 0.66
High	>0 - 1.0	0.75			>9 - <14	0.75	>6 - <9	0.75	0.67 - 0.80
Excellent	0	1.0	1.0 to 1.25	1.0	>14	1.0	>9 - 10	1.00	> 0.80

²Plant density ratings are a relative measure of the total amount of submergent vegetation covering the submergent zone, with a scale from 1 to 4 according to MN DNR methodology. The rating system is based on a 1 to 3 scale. Therefore the density results were converted to match the rating system.

³Density data for Orchard Lake were collected by Blue Water Science using a stratified line transect survey throughout the lake.

⁴Maximum exotic plant density ratings represent the worst case scenario of curlyleaf pondweed density early in the growing season and/or Eurasian watermilfoil when it is most prolific later in the growing season.

⁵The Total Number of Native Species within the submergent zone for Orchard Lake was collected by Blue Water Science using a stratified line transect survey. The additional category of "High" was added in 2011 through 2017 and values were adjusted to: <7 = Poor, 7-9 = Moderate, 9-14 = High, >14 = Excellent.

⁶**Overall Emergent Zone Quality** is the average of the rating scores for the following parameters within the emergent zone: the total percent coverage, the total number of native wetland plant species, the percent coverage of exotic species, and the C-Value Rating: >0.80 = Excellent, 0.67-0.80 = High, 0.33-0.66 = Moderate, <0.33 = Poor.

Overall Emergent Zone Quality	Percent Cover	Percent Cover Rating Score	Total Number of Native Wetland Plant Species	Number of Native Wetland Plant Species Rating Score	Percent Cover of Exotics	Percent Cover of Exotics Rating Score	Mean Coefficient of Conservatism Value (C-Value)	C-Value Rating (using MPCA values, 2007)	Overall Emergent Zone Quality Score
Poor	0-25%	0.1	< or= 5	0.1	76-100%	0.1	0 - <3	0.10	< 0.33
Moderate	76-100% or 26-50%	0.5	6 - 10	0.33	51-75%	0.33	>3 - <6	0.50	0.33 - 0.66
High	51-75%	1.0	11 - 15	0.66	26-50%	0.66	>6 - <9	0.75	0.67 - 0.80
Excellent	51-75%	1.0	> 15	1.0	0-25%	1.0	>9 - 10	1.00	> 0.80

Table 1: Orchard Lake 2017 Habitat Assessment Monitoring Results Black Dog Watershed Management Organization

⁷Approximate Total Percent Vegetative Cover Within the Entire Emergent Zone (0-2 ft. depth) is estimated based on the three sampling locations and a visual survey during travels around the water body. Estimates are broken into the following categories: 0-25%=Poor, 26-50%=Moderate, 51-75%=High and Excellent, 76-100%=Moderate.

⁸The Total Number of Native Wetland Plant Species within the emergent zone is based on 3 sampling locations, a meandering visual survey during travels on the water body, and walking along the shoreline: 0-5 = Poor, 6-10 = Moderate, 11-15 = High, and >15 = Excellent.

⁹Total Exotic Emergent Percent Coverage, out of the entire emergent zone area, is estimated based on two plot locations, a meandering visual survey during travels on the water body, and walking along the shoreline. Estimates are broken into four categories: 0-25%=Excellent (1.0), 26-50%=High (0.66), 51-75%=Moderate (0.33), 76-100%=Poor (0.1)

¹⁰**Overall Upland Buffer Quality** is determined based on the average of the six upland buffer quality parameter rating scores: >0.80 = Excellent, 0.67-0.80 = High, 0.33-0.66 = Moderate, <0.33 = Poor.

Overall Upland Buffer Quality	Percent Cover	Percent Cover Rating Score	Exotics Percent Cover Range	Exotics Percent Cover Rating Score	Buffer Width Range	Buffer Width Rating Score	Buffer Continuity Percent Range	Buffer Continuity Rating Score	Mean Coefficient of Conservatism Value (C-Value)	C-Value Rating (using MPCA values, 2007)	Number of Native Species	Number of Native Species Rating Score	Overall Upland Buffer Quality Score
Poor	<75%	0.1	>40%	0.1	<10 ft.	0.1	0-25%	0.1	0 - <3	0.10	<5	0.1	< 0.33
Moderate	75-95%	0.5	15-40%	0.5	10-25 ft.	0.4	25-50%	0.4	>3 - <6	0.50	5-20	0.33	0.33 - 0.66
High	>95%	1.0	<15%	1.0	25-50 ft.	0.7	51-75%	0.7	>6 - <9	0.75	20-30	0.66	0.67 - 0.80
Excellent	>95%	1.0	<15%	1.0	>50 ft.	1.0	76-100%	1.0	>9 - 10	1.00	>30	1.0	> 0.80

¹¹Unmanicured (upland) Buffer Width is divided into four categories: Excellent (1.0) = >50 ft, High (0.7) = 25-50 ft, Moderate (0.4) = 10-25 ft, and Low (0.1) = <10 ft.

¹²Estimated Total Vegetative Cover (Percent Range) for upland buffer is the proportion of the ground covered by vegetation within 50 feet of the wetland/upland transition zone. The percent cover is divided into three categories: High and Excellent (1.0) = >95%, Moderate (0.5) = 75 - 95%, and Poor (0.1) = <75%.

¹³The Total Number of Native Plant Species within the unmanicured upland buffer zone is based on two plot locations and a meandering visual survey along the shoreline.

¹⁴(Upland) Buffer Continuity is a measure of the proportion of the water body surrounded by the unmanicured, native upland buffer. This measure is divided into four categories: Excellent (1.0) = 76 - 100%, High (0.7) = 51 - 75%, Medium (0.4) = 26 - 50%, and Low (0.1) = 0 - 25%.

¹⁵Upland buffer exotic species "Percent of Total Coverage" is the percent cover of exotic species within the unmanicured upland buffer, which is divided into three categories: High and Excellent (1.0) = <15%, Moderate (0.5) = 15 - 40%, and Poor (0.1) = >40%.

¹⁶The presence of shoreline erosion is determined by the approximate percentage of the shoreline affected and is divided into the following three categories: 0 - 10%, 11 - 25%, 26 - 100%.

Table 1: Crystal Lake 2013 and 2018 Habitat Assessment Monitoring Results Black Dog Watershed Management Organization

Monitoring Year	Submergent Zone Sampling								
	Approximate Proportion of the Water Body Which is Deep Water Habitat (~ > 20 ft. depth)	Overall Submergent Zone Quality ¹	Approximate Proportion of Water Body Typically Dominated By Submergent Vegetation (~ 2 - 20 ft. depth)	Native Species		Mean Coefficient of Conservatism Value	Exotic Species		
				Average Native Plant Density Rating ^{2,3}	Total Number of Native Species ⁵		Total Number of Species	Average Exotic Plant Density Rating ^{2,3}	Maximum Exotic Plant Density Rating ⁴
2013	15%	High	80%	1.2 (Excellent)	18 (Excellent)	4.9 (Moderate)	2	1.8 (Moderate)	2.2 (Poor)
2018	15%	High	80%	1.2 (Excellent)	15 (Excellent)	5.0 (Moderate)	2	1.2 (Moderate)	1.4 (Moderate)

Monitoring Year	Vegetated Emergent Zone Sampling						
	Overall Emergent Zone Quality ⁶	Approximate Proportion of Emergent Zone (0 - 2 ft. depth) Within The Water Body	Approximate Total Percent Vegetative Cover Within The Entire Emergent Zone ⁷	Total Number of Native Wetland Plant Species ⁸	Mean Coefficient of Conservatism Value	Exotic Species	
						Number of Species	Total Exotic Emergent Percent Coverage ⁹
2013	High	5%	26-50% (Moderate)	36 (Excellent)	3.0 (Moderate)	10	26-50% (High)
2018	High	5%	26-50% (Moderate)	50 (Excellent)	3.3 (Moderate)	9	26-50% (High)

Monitoring Year	Upland Buffer Sampling								Erosion/Sedimentation	
	Overall Upland Buffer Quality ¹⁰	Unmanicured Buffer Width ¹¹	Estimated Total Vegetative Cover (Percent Range) ¹²	Total Number of Native Plant Species ¹³	Mean Coefficient of Conservatism Value	Buffer Continuity (Percent Surrounding Water Body) ¹⁴	Exotic Species		Shoreline Erosion (Percent of Shoreline) ¹⁶	Sediment Deltas (Yes/No)
							Number of Species	Percent of Total Coverage ¹⁵		
2013	Moderate	<10 ft. (Poor)	>95% (High)	39 (Excellent)	2.6 (Poor)	26-50% (Moderate)	16	15-40% (Moderate)	0-10%	No
2018	Moderate	<10 ft. (Poor)	>95% (High)	54 (Excellent)	2.7 (Poor)	26-50% (Moderate)	20	15-40% (Moderate)	0-10%	No

Table 1: Crystal Lake 2018 Habitat Assessment Monitoring Results Black Dog Watershed Management Organization

The following changes were made to the 2011 - 2018 monitoring and analysis:

- Monitor one or two water bodies per year. Kingsley Lake in 2011 and 2016, Orchard Lake in 2012 and 2017, Crystal Lake in 2013 and 2018, Lac Lavon in 2014, Keller Lake in 2015 - Conduct a meandering survey of submergent, emergent, and upland buffer zones. In addition, the emergent and upland buffer plot locations were evaluated.
- Changes were made in 2011 through 2018 to the calculations to include floristic quality as part of the assessment. These changes include adding a rating of "High" to the categories to accommodate MPCA ratings for floristic quality. These changes included adding a Rating Code:

Poor	Moderate	High or Excellent
------	----------	-------------------

The following footnotes pertain to 2011 through 2018 data:

¹**Overall Submergent Zone Quality** rating is the average of the rating scores for the following parameters: average exotic plant density, average native plant density, total number of native species, and C-value rating: >0.80 = Excellent, 0.67-0.80 = High, 0.33-0.66 = Moderate, <0.33 = Poor.

Overall Submergent Zone Quality	Avg. Exotic Plant Density	Exotic Plant Density Rating Score	Avg. Native Plant Density	Avg. Native Plant Density Rating Score	Total Number of Native Species In Submergent Zone	Species Richness Rating Score	Mean Coefficient of Conservatism Value (C-Value)	C-Value Rating (using MPCA values, 2007)	Total Overall Submergent Zone Quality Score
Poor	>2.0	0.1	> 1.75	0.1	<7	0.1	0 - <3	0.10	< 0.33
Moderate	>1.0 - 2.0	0.5	1.25 - 1.75	0.5	>7 - <9	0.5	>3 - <6	0.50	0.33 - 0.66
High	>0 - 1.0	0.75			>9 - <14	0.75	>6 - <9	0.75	0.67 - 0.80
Excellent	0	1.0	1.0 to 1.25	1.0	>14	1.0	>9 - 10	1.00	> 0.80

²Plant density ratings are a relative measure of the total amount of submergent vegetation covering the submergent zone, with a scale from 1 to 3.

³Density data for Orchard Lake were collected by Blue Water Science using a stratified line transect survey throughout the lake.

⁴Maximum exotic plant density ratings represent the worst case scenario of curlyleaf pondweed density early in the growing season and/or Eurasian watermilfoil when it is most prolific later in the growing season.

⁵The Total Number of Native Species within the submergent zone for Orchard Lake was collected by Blue Water Science using a stratified line transect survey. The additional category of "High" was added in 2011 through 2018 and values were adjusted to: <7 = Poor, 7-9 = Moderate, 9-14 = High, >14 = Excellent.

⁶**Overall Emergent Zone Quality** is the average of the rating scores for the following parameters within the emergent zone: the total percent coverage, the total number of native wetland plant species, the percent coverage of exotic species, and the C-Value Rating: >0.80 = Excellent, 0.67-0.80 = High, 0.33-0.66 = Moderate, <0.33 = Poor.

Overall Emergent Zone Quality	Percent Cover	Percent Cover Rating Score	Total Number of Native Wetland Plant Species	Number of Native Wetland Plant Species Rating Score	Percent Cover of Exotics	Percent Cover of Exotics Rating Score	Mean Coefficient of Conservatism Value (C-Value)	C-Value Rating (using MPCA values, 2007)	Overall Emergent Zone Quality Score
Poor	0-25%	0.1	< or= 5	0.1	76-100%	0.1	0 - <3	0.10	< 0.33
Moderate	76-100% or 26-50%	0.5	6 - 10	0.33	51-75%	0.33	>3 - <6	0.50	0.33 - 0.66
High	51-75%	1.0	11 - 15	0.66	26-50%	0.66	>6 - <9	0.75	0.67 - 0.80
Excellent	51-75%	1.0	> 15	1.0	0-25%	1.0	>9 - 10	1.00	> 0.80

Table 1: Crystal Lake 2018 Habitat Assessment Monitoring Results Black Dog Watershed Management Organization

⁷Approximate Total Percent Vegetative Cover Within the Entire Emergent Zone (0-2 ft. depth) is estimated based on the three sampling locations and a visual survey during travels around the water body. Estimates are broken into the following categories: 0-25%=Poor, 26-50%=Moderate, 51-75%=High and Excellent, 76-100%=Moderate.

⁸The Total Number of Native Wetland Plant Species within the emergent zone is based on 3 sampling locations, a meandering visual survey during travels on the water body, and walking along the shoreline: 0-5 = Poor, 6-10 = Moderate, 11-15 = High, and >15 = Excellent.

⁹Total Exotic Emergent Percent Coverage, out of the entire emergent zone area, is estimated based on two plot locations, a meandering visual survey during travels on the water body, and walking along the shoreline. Estimates are broken into four categories: 0-25%=Excellent (1.0), 26-50%=High (0.66), 51-75%=Moderate (0.33), 76-100%=Poor (0.1)

¹⁰**Overall Upland Buffer Quality** is determined based on the average of the six upland buffer quality parameter rating scores: >0.80 = Excellent, 0.67-0.80 = High, 0.33-0.66 = Moderate, <0.33 = Poor.

Overall Upland Buffer Quality	Percent Cover	Percent Cover Rating Score	Exotics Percent Cover Range	Exotics Percent Cover Rating Score	Buffer Width Range	Buffer Width Rating Score	Buffer Continuity Percent Range	Buffer Continuity Rating Score	Mean Coefficient of Conservatism Value (C-Value)	C-Value Rating (using MPCA values, 2007)	Number of Native Species	Number of Native Species Rating Score	Overall Upland Buffer Quality Score
Poor	<75%	0.1	>40%	0.1	<10 ft.	0.1	0-25%	0.1	0 - <3	0.10	<5	0.1	< 0.33
Moderate	75-95%	0.5	15-40%	0.5	10-25 ft.	0.4	25-50%	0.4	>3 - <6	0.50	5-20	0.33	0.33 - 0.66
High	>95%	1.0	<15%	1.0	25-50 ft.	0.7	51-75%	0.7	>6 - <9	0.75	20-30	0.66	0.67 - 0.80
Excellent	>95%	1.0	<15%	1.0	>50 ft.	1.0	76-100%	1.0	>9 - 10	1.00	>30	1.0	> 0.80

¹¹Unmanicured (upland) Buffer Width is divided into four categories: Excellent (1.0) = >50 ft, High (0.7) = 25-50 ft, Moderate (0.4) = 10-25 ft, and Low (0.1) = <10 ft.

¹²Estimated Total Vegetative Cover (Percent Range) for upland buffer is the proportion of the ground covered by vegetation within 50 feet of the wetland/upland transition zone. The percent cover is divided into three categories: High and Excellent (1.0) = >95%, Moderate (0.5) = 75 - 95%, and Poor (0.1) = <75%.

¹³The Total Number of Native Plant Species within the unmanicured upland buffer zone is based on two plot locations and a meandering visual survey along the shoreline.

¹⁴(Upland) Buffer Continuity is a measure of the proportion of the water body surrounded by the unmanicured, native upland buffer. This measure is divided into four categories: Excellent (1.0) = 76 - 100%, High (0.7) = 51 - 75%, Medium (0.4) = 26 - 50%, and Low (0.1) = 0 - 25%.

¹⁵Upland buffer exotic species "Percent of Total Coverage" is the percent cover of exotic species within the unmanicured upland buffer, which is divided into three categories: High and Excellent (1.0) = <15%, Moderate (0.5) = 15 - 40%, and Poor (0.1) = >40%.

¹⁶The presence of shoreline erosion is determined by the approximate percentage of the shoreline affected and is divided into the following three categories: 0 - 10%, 11 - 25%, 26 - 100%.

Table 1: Lac Lavon 2014 and 2019 Habitat Assessment Monitoring Results Black Dog Watershed Management Organization

Monitoring Year	Submergent Zone Sampling								
	Approximate Proportion of the Water Body Which is Deep Water Habitat (~ > 20 ft. depth)	Overall Submergent Zone Quality ¹	Approximate Proportion of Water Body Typically Dominated By Submergent Vegetation (~ 2 - 20 ft. depth)	Native Species		Mean Coefficient of Conservatism Value	Exotic Species		
				Average Native Plant Density Rating ^{2,3}	Total Number of Native Species ⁵		Total Number of Species	Average Exotic Plant Density Rating ^{2,3}	Maximum Exotic Plant Density Rating ⁴
2014	25%	Moderate	70%	1.4 (Moderate)	12 (High)	4.6 (Moderate)	2	2.0 (Moderate)	3.0 (Poor)
2019	25%	Moderate	70%	1.5 (Moderate)	12 (High)	4.5 (Moderate)	2	1.7 (Moderate)	3.0 (Poor)

Monitoring Year	Vegetated Emergent Zone Sampling						
	Overall Emergent Zone Quality ⁶	Approximate Proportion of Emergent Zone (0 - 2 ft. depth) Within The Water Body	Approximate Total Percent Vegetative Cover Within The Entire Emergent Zone ⁷	Total Number of Native Wetland Plant Species ⁸	Mean Coefficient of Conservatism Value	Exotic Species	
						Number of Species	Total Exotic Emergent Percent Coverage ⁹
2014	Moderate	5%	0-25% (Poor)	32 (Excellent)	2.3 (Poor)	15	26-50% (High)
2019	Moderate	5%	0-25% (Poor)	38 (Excellent)	2.4 (Poor)	17	26-50% (High)

Monitoring Year	Upland Buffer Sampling								Erosion/Sedimentation	
	Overall Upland Buffer Quality ¹⁰	Unmanicured Buffer Width ¹¹	Estimated Total Vegetative Cover (Percent Range) ¹²	Total Number of Native Plant Species ¹³	Mean Coefficient of Conservatism Value	Buffer Continuity (Percent Surrounding Water Body) ¹⁴	Exotic Species		Shoreline Erosion (Percent of Shoreline) ¹⁶	Sediment Deltas (Yes/No)
							Number of Species	Percent of Total Coverage ¹⁵		
2014	Poor	<10 ft. (Poor)	>95% (High)	32 (Excellent)	1.3 (Poor)	0-25% (Poor)	31	>40% (Poor)	0-10%	No
2019	Poor	<10 ft. (Poor)	75-95% (Moderate)	56 (Excellent)	2.0 (Poor)	0-25% (Poor)	41	>40% (Poor)	0-10%	No

Table 1: Lac Lavon 2019 Habitat Assessment Monitoring Results Black Dog Watershed Management Organization

The following changes were made to the 2011 - 2019 monitoring and analysis:

- Monitor one or two water bodies per year. Kingsley Lake in 2011 and 2016, Orchard Lake in 2012 and 2017, Crystal Lake in 2013 and 2018, Lac Lavon in 2014 and 2019, Keller Lake in 2015 - Conduct a meandering survey of submergent, emergent, and upland buffer zones. In addition, the emergent and upland buffer plot locations were evaluated.
- Changes were made in 2011 through 2019 to the calculations to include floristic quality as part of the assessment. These changes include adding a rating of "High" to the categories to accommodate MPCA ratings for floristic quality. These changes included adding a Rating Code:

Poor	Moderate	High or Excellent
------	----------	-------------------

The following footnotes pertain to 2011 through 2019 data:

¹**Overall Submergent Zone Quality** rating is the average of the rating scores for the following parameters: average exotic plant density, average native plant density, total number of native species, and C-value rating: >0.80 = Excellent, 0.67-0.80 = High, 0.33-0.66 = Moderate, <0.33 = Poor.

Overall Submergent Zone Quality	Avg. Exotic Plant Density	Exotic Plant Density Rating Score	Avg. Native Plant Density	Avg. Native Plant Density Rating Score	Total Number of Native Species In Submergent Zone	Species Richness Rating Score	Mean Coefficient of Conservatism Value (C-Value)	C-Value Rating (using MPCA values, 2007)	Total Overall Submergent Zone Quality Score
Poor	>2.0	0.1	> 1.75	0.1	<7	0.1	0 - <3	0.10	< 0.33
Moderate	>1.0 - 2.0	0.5	1.25 - 1.75	0.5	>7 - <9	0.5	>3 - <6	0.50	0.33 - 0.66
High	>0 - 1.0	0.75			>9 - <14	0.75	>6 - <9	0.75	0.67 - 0.80
Excellent	0	1.0	1.0 to 1.25	1.0	>14	1.0	>9 - 10	1.00	> 0.80

²Plant density ratings are a relative measure of the total amount of submergent vegetation covering the submergent zone, with a scale from 1 to 3.

³Density data for Lac Lavon were collected by Matt Berg using a point intercept survey throughout the lake.

⁴Maximum exotic plant density ratings represent the worst case scenario of curlyleaf pondweed density early in the growing season and/or Eurasian watermilfoil when it is most prolific later in the growing season.

⁵The Total Number of Native Species within the submergent zone for Lac Lavon was collected by Matt Berg using a point intercept survey.

The additional category of "High" was added in 2011 through 2019 and values were adjusted to: <7 = Poor, 7-9 = Moderate, 9-14 = High, >14 = Excellent.

⁶**Overall Emergent Zone Quality** is the average of the rating scores for the following parameters within the emergent zone: the total percent coverage, the total number of native wetland plant species, the percent coverage of exotic species, and the C-Value Rating: >0.80 = Excellent, 0.67-0.80 = High, 0.33-0.66 = Moderate, <0.33 = Poor.

Overall Emergent Zone Quality	Percent Cover	Percent Cover Rating Score	Total Number of Native Wetland Plant Species	Number of Native Wetland Plant Species Rating Score	Percent Cover of Exotics	Percent Cover of Exotics Rating Score	Mean Coefficient of Conservatism Value (C-Value)	C-Value Rating (using MPCA values, 2007)	Overall Emergent Zone Quality Score
Poor	0-25%	0.1	< or= 5	0.1	76-100%	0.1	0 - <3	0.10	< 0.33
Moderate	76-100% or 26-50%	0.5	6 - 10	0.33	51-75%	0.33	>3 - <6	0.50	0.33 - 0.66
High	51-75%	1.0	11 - 15	0.66	26-50%	0.66	>6 - <9	0.75	0.67 - 0.80
Excellent	51-75%	1.0	> 15	1.0	0-25%	1.0	>9 - 10	1.00	> 0.80

Table 1: Lac Lavon 2019 Habitat Assessment Monitoring Results Black Dog Watershed Management Organization

⁷Approximate Total Percent Vegetative Cover Within the Entire Emergent Zone (0-2 ft. depth) is estimated based on the three sampling locations and a visual survey during travels around the water body. Estimates are broken into the following categories: 0-25%=Poor, 26-50%=Moderate, 51-75%=High and Excellent, 76-100%=Moderate.

⁸The Total Number of Native Wetland Plant Species within the emergent zone is based on 3 sampling locations, a meandering visual survey during travels on the water body, and walking along the shoreline: 0-5 = Poor, 6-10 = Moderate, 11-15 = High, and >15 = Excellent.

⁹Total Exotic Emergent Percent Coverage, out of the entire emergent zone area, is estimated based on two plot locations, a meandering visual survey during travels on the water body, and walking along the shoreline. Estimates are broken into four categories: 0-25%=Excellent (1.0), 26-50%=High (0.66), 51-75%=Moderate (0.33), 76-100%=Poor (0.1)

¹⁰**Overall Upland Buffer Quality** is determined based on the average of the six upland buffer quality parameter rating scores: >0.80 = Excellent, 0.67-0.80 = High, 0.33-0.66 = Moderate, <0.33 = Poor.

Overall Upland Buffer Quality	Percent Cover	Percent Cover Rating Score	Exotics Percent Cover Range	Exotics Percent Cover Rating Score	Buffer Width Range	Buffer Width Rating Score	Buffer Continuity Percent Range	Buffer Continuity Rating Score	Mean Coefficient of Conservatism Value (C-Value)	C-Value Rating (using MPCA values, 2007)	Number of Native Species	Number of Native Species Rating Score	Overall Upland Buffer Quality Score
Poor	<75%	0.1	>40%	0.1	<10 ft.	0.1	0-25%	0.1	0 - <3	0.10	<5	0.1	< 0.33
Moderate	75-95%	0.5	15-40%	0.5	10-25 ft.	0.4	25-50%	0.4	>3 - <6	0.50	5-20	0.33	0.33 - 0.66
High	>95%	1.0	<15%	1.0	25-50 ft.	0.7	51-75%	0.7	>6 - <9	0.75	20-30	0.66	0.67 - 0.80
Excellent	>95%	1.0	<15%	1.0	>50 ft.	1.0	76-100%	1.0	>9 - 10	1.00	>30	1.0	> 0.80

¹¹Unmanicured (upland) Buffer Width is divided into four categories: Excellent (1.0) = >50 ft, High (0.7) = 25-50 ft, Moderate (0.4) = 10-25 ft, and Low (0.1) = <10 ft.

¹²Estimated Total Vegetative Cover (Percent Range) for upland buffer is the proportion of the ground covered by vegetation within 50 feet of the wetland/upland transition zone. The percent cover is divided into three categories: High and Excellent (1.0) = >95%, Moderate (0.5) = 75 - 95%, and Poor (0.1) = <75%.

¹³The Total Number of Native Plant Species within the unmanicured upland buffer zone is based on two plot locations and a meandering visual survey along the shoreline.

¹⁴(Upland) Buffer Continuity is a measure of the proportion of the water body surrounded by the unmanicured, native upland buffer. This measure is divided into four categories: Excellent (1.0) = 76 - 100%, High (0.7) = 51 - 75%, Medium (0.4) = 26 - 50%, and Low (0.1) = 0 - 25%.

¹⁵Upland buffer exotic species "Percent of Total Coverage" is the percent cover of exotic species within the unmanicured upland buffer, which is divided into three categories: High and Excellent (1.0) = <15%, Moderate (0.5) = 15 - 40%, and Poor (0.1) = >40%.

¹⁶The presence of shoreline erosion is determined by the approximate percentage of the shoreline affected and is divided into the following three categories: 0 - 10%, 11 - 25%, 26 - 100%.

Table 1: Keller Lake 2015 and 2020 Habitat Assessment Monitoring Results Black Dog Watershed Management Organization

Monitoring Year	Submergent Zone								
	Approximate Proportion of the Water Body Which is Deep Water Habitat (~ > 20 ft. depth)	Overall Submergent Zone Quality ¹	Approximate Proportion of Water Body Typically Dominated By Submergent Vegetation (~ 2 - 20 ft. depth)	Native Species		Mean Coefficient of Conservatism Value	Exotic Species		
				Average Native Plant Density Rating ^{2,3}	Total Number of Native Species ⁵		Total Number of Species	Average Exotic Plant Density Rating ^{2,3}	Maximum Exotic Plant Density Rating ⁴
2015	0%	Poor	90%	1.3 (Moderate)	2 (Poor)	1.5 (Poor)	2	1.8 (Moderate)	2.2 (Poor)
2020	0%	Moderate	90%	1.2 (Excellent)	2 (Poor)	1.5 (Poor)	2	1.8 (Moderate)	2.3 (Poor)

Monitoring Year	Vegetated Emergent Zone						
	Overall Emergent Zone Quality ⁶	Approximate Proportion of Emergent Zone (0 - 2 ft. depth) Within The Water Body	Approximate Total Percent Vegetative Cover Within The Entire Emergent Zone ⁷	Total Number of Native Wetland Plant Species ⁸	Mean Coefficient of Conservatism Value	Exotic Species	
						Number of Species	Total Exotic Emergent Percent Coverage ⁹
2015	High	10%	51-75% (High)	28 (Excellent)	2.3 (Poor)	8	26-50% (High)
2020	High	10%	51-75% (High)	36 (Excellent)	2.4 (Poor)	10	26-50% (High)

Monitoring Year	Upland Buffer								Erosion/Sedimentation	
	Overall Upland Buffer Quality ¹⁰	Unmanicured Buffer Width ¹¹	Estimated Total Vegetative Cover (Percent Range) ¹²	Total Number of Native Plant Species ¹³	Mean Coefficient of Conservatism Value	Buffer Continuity (Percent Surrounding Water Body) ¹⁴	Exotic Species		Shoreline Erosion (Percent of Shoreline) ¹⁶	Sediment Deltas (Yes/No)
							Number of Species	Percent of Total Coverage ¹⁵		
2015	Moderate	25-50 ft. (High)	>95% (High)	20 (Moderate)	1.6 (Poor)	76-100% (Excellent)	10	>40% (Poor)	0-10%	No
2020	Moderate	25-50 ft. (High)	>95% (High)	42 (Excellent)	1.8 (Poor)	51-75% (High)	29	>40% (Poor)	0-10%	No

Table 1: Keller Lake 2020 Habitat Assessment Monitoring Results Black Dog Watershed Management Organization

The following changes were made to the 2011 - 2020 monitoring and analysis:

- Monitor one or two water bodies per year. Kingsley Lake in 2011 and 2016, Orchard Lake in 2012 and 2017, Crystal Lake in 2013 and 2018, Lac Lavon in 2014 and 2019, Keller Lake in 2015 and 2020 - Conduct a meandering survey of submergent, emergent, and upland buffer zones. In addition, the emergent and upland buffer plot locations were evaluated.
- Changes were made in 2011 through 2020 to the calculations to include floristic quality as part of the assessment. These changes include adding a rating of "High" to the categories to accommodate MPCA ratings for floristic quality. These changes included adding a Rating Code:

Poor	Moderate	High or Excellent
------	----------	-------------------

The following footnotes pertain to 2011 through 2020 data:

¹**Overall Submergent Zone Quality** rating is the average of the rating scores for the following parameters: average exotic plant density, average native plant density, total number of native species, and C-value rating: >0.80 = Excellent, 0.67-0.80 = High, 0.33-0.66 = Moderate, <0.33 = Poor.

Overall Submergent Zone Quality	Avg. Exotic Plant Density	Exotic Plant Density Rating Score	Avg. Native Plant Density	Avg. Native Plant Density Rating Score	Total Number of Native Species In Submergent Zone	Species Richness Rating Score	Mean Coefficient of Conservatism Value (C-Value)	C-Value Rating (using MPCA values, 2007)	Total Overall Submergent Zone Quality Score
Poor	>2.0	0.1	> 1.75	0.1	<7	0.1	0 - <3	0.10	< 0.33
Moderate	>1.0 - 2.0	0.5	1.25 - 1.75	0.5	>7 - <9	0.5	>3 - <6	0.50	0.33 - 0.66
High	>0 - 1.0	0.75			>9 - <14	0.75	>6 - <9	0.75	0.67 - 0.80
Excellent	0	1.0	1.0 to 1.25	1.0	>14	1.0	>9 - 10	1.00	> 0.80

²Plant density ratings are a relative measure of the total amount of submergent vegetation covering the submergent zone, with a scale from 1 to 3.

³Density data for Keller Lake were collected by Blue Water Science using a point intercept survey throughout the lake.

⁴Maximum exotic plant density ratings represent the worst case scenario of curlyleaf pondweed density early in the growing season and/or Eurasian watermilfoil when it is most prolific later in the growing season.

⁵The Total Number of Native Species within the submergent zone for Keller Lake was collected by Blue Water Science using a point intercept survey.

The additional category of "High" was added in 2011 through 2020 and values were adjusted to: <7 = Poor, 7-9 = Moderate, 9-14 = High, >14 = Excellent.

⁶**Overall Emergent Zone Quality** is the average of the rating scores for the following parameters within the emergent zone: the total percent coverage, the total number of native wetland plant species, the percent coverage of exotic species, and the C-Value Rating: >0.80 = Excellent, 0.67-0.80 = High, 0.33-0.66 = Moderate, <0.33 = Poor.

Overall Emergent Zone Quality	Percent Cover	Percent Cover Rating Score	Total Number of Native Wetland Plant Species	Number of Native Wetland Plant Species Rating Score	Percent Cover of Exotics	Percent Cover of Exotics Rating Score	Mean Coefficient of Conservatism Value (C-Value)	C-Value Rating (using MPCA values, 2007)	Overall Emergent Zone Quality Score
Poor	0-25%	0.1	< or= 5	0.1	76-100%	0.1	0 - <3	0.10	< 0.33
Moderate	76-100% or 26-50%	0.5	6 - 10	0.33	51-75%	0.33	>3 - <6	0.50	0.33 - 0.66
High	51-75%	1.0	11 - 15	0.66	26-50%	0.66	>6 - <9	0.75	0.67 - 0.80
Excellent	51-75%	1.0	> 15	1.0	0-25%	1.0	>9 - 10	1.00	> 0.80

Table 1: Keller Lake 2020 Habitat Assessment Monitoring Results Black Dog Watershed Management Organization

⁷Approximate Total Percent Vegetative Cover Within the Entire Emergent Zone (0-2 ft. depth) is estimated based on the two plot locations and a visual survey walking along the shoreline. Estimates are broken into the following categories: 0-25%=Poor, 26-50%=Moderate, 51-75%=High and Excellent, 76-100%=Moderate.

⁸The Total Number of Native Wetland Plant Species within the emergent zone is based on 2 plot locations and a visual survey walking along the shoreline: 0-5 = Poor, 6-10 = Moderate, 11-15 = High, and >15 = Excellent.

⁹Total Exotic Emergent Percent Coverage, out of the entire emergent zone area, is estimated based on two plot locations, a visual survey walking along the shoreline. Estimates are broken into four categories: 0-25%=Excellent (1.0), 26-50%=High (0.66), 51-75%=Moderate (0.33), 76-100%=Poor (0.1)

¹⁰**Overall Upland Buffer Quality** is determined based on the average of the six upland buffer quality parameter rating scores: >0.80 = Excellent, 0.67-0.80 = High, 0.33-0.66 = Moderate, <0.33 = Poor.

Overall Upland Buffer Quality	Percent Cover	Percent Cover Rating Score	Exotics Percent Cover Range	Exotics Percent Cover Rating Score	Buffer Width Range	Buffer Width Rating Score	Buffer Continuity Percent Range	Buffer Continuity Rating Score	Mean Coefficient of Conservatism Value (C-Value)	C-Value Rating (using MPCA values, 2007)	Number of Native Species	Number of Native Species Rating Score	Overall Upland Buffer Quality Score
Poor	<75%	0.1	>40%	0.1	<10 ft.	0.1	0-25%	0.1	0 - <3	0.10	<5	0.1	< 0.33
Moderate	75-95%	0.5	15-40%	0.5	10-25 ft.	0.4	25-50%	0.4	>3 - <6	0.50	5-20	0.33	0.33 - 0.66
High	>95%	1.0	<15%	1.0	25-50 ft.	0.7	51-75%	0.7	>6 - <9	0.75	20-30	0.66	0.67 - 0.80
Excellent	>95%	1.0	<15%	1.0	>50 ft.	1.0	76-100%	1.0	>9 - 10	1.00	>30	1.0	> 0.80

¹¹Unmanicured (upland) Buffer Width is divided into four categories: Excellent (1.0) = >50 ft, High (0.7) = 25-50 ft, Moderate (0.4) = 10-25 ft, and Low (0.1) = <10 ft.

¹²Estimated Total Vegetative Cover (Percent Range) for upland buffer is the proportion of the ground covered by vegetation within 50 feet of the wetland/upland transition zone. The percent cover is divided into three categories: High and Excellent (1.0) = >95%, Moderate (0.5) = 75 - 95%, and Poor (0.1) = <75%.

¹³The Total Number of Native Plant Species within the unmanicured upland buffer zone is based on two plot locations and a meandering visual survey along the shoreline.

¹⁴(Upland) Buffer Continuity is a measure of the proportion of the water body surrounded by the unmanicured, native upland buffer. This measure is divided into four categories: Excellent (1.0) = 76 - 100%, High (0.7) = 51 - 75%, Medium (0.4) = 26 - 50%, and Low (0.1) = 0 - 25%.

¹⁵Upland buffer exotic species "Percent of Total Coverage" is the percent cover of exotic species within the unmanicured upland buffer, which is divided into three categories: High and Excellent (1.0) = <15%, Moderate (0.5) = 15 - 40%, and Poor (0.1) = >40%.

¹⁶The presence of shoreline erosion is determined by the approximate percentage of the shoreline affected and is divided into the following three categories: 0 - 10%, 11 - 25%, 26 - 100%.

Appendix D

2003–2020 Recommended and Completed Management Actions

**Table D-1: 2009 Recommended and Completed Management Actions
Black Dog Watershed Management Organization Habitat Monitoring**

Strategic Water Body	Problem Identified	Recommendation	Proposed Action	Benefits	Implementation Period	Completed 2004-2009 Actions Which May Improve Wildlife Habitat and/or Water Quality
Crystal	Unmanicured, native vegetation in adjacent upland and emergent zone is narrow and not continuous, limiting wildlife benefits.	1. Increase width and continuity of native upland buffer and emergent zone.	Conduct an educational workshop and lakescaping demonstration project. Assist lakeshore owners with funding and obtaining any needed MnDNR permits for potential upland buffer and emergent zone enhancements. Restore sustainable native communities	Inform/show lakeshore property owners how a native upland buffer and native emergent zone can improve functions and values of the lake and improve aesthetics. Increase wildlife habitat.	Spring - Fall Spring - Fall	2009: Operation of the ferric chloride treatment system halted due to low water levels. The City of Burnsville harvested curlyleaf pondweed. In late 2009, the City of Burnsville treated 14 acres of buckthorn within Crystal West Park. In 2009 and 2008, garlic mustard within the upland buffer was removed/pulled. 2004-2008: The BDWMO resumed and continued operation of the ferric chloride treatment system. The City of Burnsville: 1) excavated/enhanced four stormwater treatment ponds (including West Buck Hill Park), which reduced the phosphorus loading into the lake, and 2) conducted annual harvesting of Eurasian watermilfoil and curlyleaf pondweed. The City of Lakeville excavated/enhanced the Bluebird stormwater treatment pond.
	Purple loosestrife is present.	2. Continue to control and manage purple loosestrife.	Control and manage. For large stands of purple loosestrife, contact the MnDNR to request a release of purple loosestrife-controlling beetles. For a few small colonies of purple loosestrife, hand pull or dig the plants out before they go to seed.	Increase/maintain wildlife habitat.	Spring - Fall	
	Curlyleaf pondweed is present.	3. Control curlyleaf pondweed	Control by harvesting or chemical treatment.	Maintain wildlife habitat.	Late Spring	
	Eurasian watermilfoil is present.	4. Control Eurasian watermilfoil.	Control by chemical treatment.	Maintain wildlife habitat.	Summer	
Keller Lake	Unmanicured, native vegetation in adjacent upland is narrow and not continuous, limiting wildlife benefits.	1. Increase width and continuity of native upland buffer.	Conduct an educational workshop and lakescaping demonstration project. Assist lakeshore owners with funding of potential upland buffer enhancements. Restore sustainable native communities	Inform/show lakeshore property owners how a native upland buffer can improve functions and values of the lake and improve aesthetics. Increase wildlife habitat.	Spring - Fall Spring - Fall	In 2010 the City of Apple Valley may construct Whitney Pond for stormwater treatment within the Keller Lake watershed. 2009: Due to low water levels, operation of the ferric chloride treatment system halted and no harvesting of curlyleaf pondweed was conducted. The City of Burnsville stabilized approximately one hundred feet of shoreline on the southeast edge of the lake. Logs were interlaced and secured along the shoreline and red-osier dogwood live stakes were installed along the eroding banks. 2004 - 2008: The Cities of Apple Valley and Burnsville partnered to conduct annual harvesting of curlyleaf pondweed. 2005: The City of Apple Valley excavated and enhanced Redwood Pond, which will decrease phosphorus loading into Keller Lake. Also, in 2010 the City of Apple Valley may construct Whitney Pond for stormwater treatment within the Keller Lake watershed.
	Purple loosestrife is present.	2. Continue to control and manage purple loosestrife.	Control and manage. For large stands of purple loosestrife, contact the MnDNR to request a release of purple loosestrife-controlling beetles. For a few small colonies of purple loosestrife, hand pull or dig the plants out before they go to seed.	Increase/maintain wildlife habitat.	Spring - Fall	
	Curlyleaf pondweed dominates the lake in late spring-early summer.	3. Continue control of curlyleaf pondweed.	Control as recommended by the MnDNR. Since the MnDNR designates Keller Lake as a "Natural Environment", a special permit is needed to chemically treat the lake.	Maintain wildlife habitat.	Summer	
	Eurasian watermilfoil is present.	4. Control Eurasian watermilfoil.	Control as recommended by the MnDNR. Since the MnDNR designates Keller Lake as a "Natural Environment", a special permit is needed to chemically treat the lake.	Maintain wildlife habitat.	Summer	
Kingsley Lake	Curlyleaf pondweed is present.	1. Conduct a detailed late spring macrophyte survey to ascertain densities and extent of coverage.	Consider control measures, dependent on results of an detailed early growing season survey.	Maintain wildlife habitat.	Late Spring	2005 - 2008: Annually, the City of Lakeville and members of the Kingsley Lake Homeowner's Association removed purple loosestrife plants and common buckthorn from portions of the lake and the upland buffer surrounding the lake. On March 6, 2008, soil sediment samples were collected on Kingsley Lake by Blue Water Science (BWS) and the City of Lakeville. Based on the results of the soil analysis, the BWS report stated that "curlyleaf pondweed is not expected to produce heavy growth conditions (where plants top out in a solid canopy) in Kingsley Lake." However, since curlyleaf pondweed may typically die-off prior to the early-June habitat assessment, the peak density and percent total coverage of curlyleaf pondweed is uncertain. To date, it is unclear if curlyleaf pondweed densities and percent coverage have been relatively consistent or increasing within the lake over the last few years. In 2008, a Kingsley lakeshore resident, inspired by the Blue Thumb program, commenced shoreline stabilization utilizing native plants.
	Common buckthorn dominates portions of the upland buffer.	2. Conduct an evaluation of common buckthorn, followed by removal.	Remove buckthorn. Volunteer groups and contractors can effectively remove buckthorn by pulling, cutting, and treating stumps with herbicide.	Increase wildlife habitat.	Open	
	Purple loosestrife is present.	3. Continue to control and manage purple loosestrife.	Control and manage. For large stands of purple loosestrife, contact the MnDNR to request a release of purple loosestrife-controlling beetles. For a few small colonies of purple loosestrife, hand pull or dig the plants out before they go to seed.	Increase/maintain wildlife habitat.	Spring - Fall	
	Hybrid cattail and reed canary grass are present.	4. Control hybrid cattail and reed canary grass.	Control hybrid cattail and reed canary grass now before colonies become more abundant. The herbicide Rodeo™ can be used to effectively control both invasive emergent species.	Increase/maintain wildlife habitat.	Spring-Summer	
Lac Lavon	Eurasian watermilfoil dominates portions of the lake.	1. Continue to manage Eurasian watermilfoil.	Control by chemical treatment as recommended by MnDNR.	Increase/maintain wildlife habitat and water quality	Spring-Summer	2006: The Cities of Burnsville and Apple Valley and the lake homeowners partnered to fund a fluridone treatment for control of Eurasian watermilfoil. The treatment is expected to provide control of Eurasian watermilfoil for three years, while allowing native plant species to rebound. The cities have continued to inform the MnDNR of the ongoing treatments and the MnDNR proposes to continue aquatic plant surveys to study the effects of whole-lake fluridone treatments. However, no MnDNR macrophyte survey was conducted in 2008.
	Curlyleaf pondweed is present.	2. Monitor presence of curlyleaf pondweed.	Control if increased occurrence and subsequent midsummer die off threatens water quality)	Identify the problem before it becomes difficult to treat.	Spring	
	Unmanicured, native vegetation in adjacent upland is narrow and not continuous, limiting wildlife benefits.	3. Increase width/creation of native upland buffer.	Conduct an educational workshop and lakescaping demonstration project. Assist lakeshore owners with funding of potential upland buffer enhancements. Restore sustainable native communities	Inform/show lakeshore property owners of how a native upland buffer can improve functions and values of the lake and improve aesthetics. Increase wildlife habitat.	Spring - Fall Spring - Fall	
Orchard Lake	Curlyleaf pondweed dominates the lake in late spring-early summer.	1. Continue curlyleaf pondweed control measures.	Control and manage	Increase/maintain wildlife habitat and water quality.	Late Spring - Early summer	2009: The City of Lakeville conducted herbicide treatment for curlyleaf pondweed within the northeast bay (~20 acres). The herbicide treatment resulted in lake-wide control of curlyleaf pondweed. 2004-2008: The City of Lakeville provides lakeshore owners with shoreline restoration information. However, to date, no plans have been made for potential future shoreline restoration projects. Annually, the City of Lakeville harvested approximately 70 acres of curlyleaf pondweed. 2007: A small area of lakeshore, near the boat launch, was restored using native plants.
	Unmanicured, native vegetation in adjacent upland is narrow and not continuous, limiting wildlife benefits.	2. Increase width and continuity of native upland buffer.	Conduct an educational workshop and lakescaping demonstration project. Assist lakeshore owners with funding of potential upland buffer enhancements. Restore sustainable native communities	Inform/show lakeshore property owners how a native upland buffer can improve functions and values of the lake and improve aesthetics. Increase wildlife habitat.	Spring - Fall Spring - Fall	
	Purple loosestrife is present.	3. Conduct a detailed evaluation of purple loosestrife, followed by removal/control.	Control and manage by hand-pulling if only a few plants are present or introduce beetles if numerous plants are present.	Increase/maintain wildlife habitat.	Spring - Summer	
Sunset Pond	Extensive algal bloom	1. Reduce phosphorus loading into the pond.	Construct/install: catch basin sumps, prefabricated treatment devices (e.g. Stormceptor), infiltration facilities within the watershed, or other more conventional methods. Conduct more frequent street sweepings.	Improve wildlife habitat, fishery habitat, and aesthetics/recreation.	Open	In 2009, as in past years, the City of Burnsville actively managed the restored native buffer adjacent to the pond, the surrounding prairie restoration area, and portions of the emergent zone. Specifically, in 2007 through 2009 the City of Burnsville conducted spot spraying of invasive vegetation, such as reed canary grass, thistle, and purple loosestrife. A prescription burn, inter-seeding of prairie species, and buckthorn removal were conducted in 2008 to increase the plant diversity in the upland area.
	Maintained turf grass remains within portions of the upland buffer.	2. Enhance/maintain upland buffer.	Continue restoring sustainable native communities	Improve wildlife habitat and water quality.	Spring - Fall	
	Exotic species are dominant in emergent zone, and include narrow-leaf cattail, hybrid cattail, and reed canary grass.	3. Manage exotic species within emergent zone.	Selective herbicide treatments to reduce presence of exotic species	Allow for the establishment of more diverse native species that provide better wildlife values.	Spring - Fall	
	Presence of curlyleaf pondweed observed in 2003 and 2005 through 2008.	4. Conduct a late spring macrophyte survey to ascertain densities and extent of coverage.	consider control measures dependent on the results of an early growing season survey.	Maintain wildlife habitat. Reduce down-stream phosphorus loading.	Late Spring	
	The southern portion of the pond is shallow (1 to 3 feet deep).	5. Create a "navigation channel".	Excavate and remove sediment.	Improve wildlife habitat, fishery habitat, and aesthetics/recreation.	Winter	

**Table 2: 2011 Recommended and Completed Management Actions for Kingsley Lake
Black Dog Watershed Management Organization Habitat Monitoring**

Problem Identified	Recommendation	Proposed Action	Benefits	Implementation Period	Completed 2004-2009 Actions Which May Improve Wildlife Habitat and/or Water Quality
Curlyleaf pondweed is present in some years.	Conduct a detailed late spring macrophyte survey to ascertain densities and extent of coverage.	Consider control measures, dependent on results of a detailed early growing season survey.	Maintain wildlife habitat.	Late Spring	<p>2005 - 2008: Annually, the City of Lakeville and members of the Kingsley Lake Homeowner's Association removed purple loosestrife plants and common buckthorn from portions of the lake and the upland buffer surrounding the lake. Purple loosestrife beetles were released by the MnDNR prior to 2002. Follow up monitoring by the MnDNR indicates that beetles are present at a population that the MnDNR feels is appropriate for biological control. On March 6, 2008, soil sediment samples were collected on Kingsley Lake by Blue Water Science (BWS) and the City of Lakeville. Based on the results of the soil analysis, the BWS report stated that "curlyleaf pondweed is not expected to produce heavy growth conditions (where plants top out in a solid canopy) in Kingsley Lake." However, since curlyleaf pondweed may typically die-off prior to the early-June habitat assessment, the peak density and percent total coverage of curlyleaf pondweed is uncertain. To date, it is unclear if curlyleaf pondweed densities and percent coverage have been relatively consistent or increasing within the lake over the last few years. In 2008, a Kingsley Lake lakeshore resident, inspired by the Blue Thumb program, commenced shoreline stabilization utilizing native plants.</p>
Common buckthorn dominates portions of the upland buffer.	Conduct an evaluation of common buckthorn, followed by removal.	Remove buckthorn. Volunteer groups and contractors can effectively remove buckthorn by pulling, cutting, and treating stumps with herbicide.	Increase wildlife habitat.	Open	
Purple loosestrife is present.	Continue to control and manage purple loosestrife.	Control and manage. For a few small colonies of purple loosestrife, hand pull or dig the plants out before they go to seed. Continue to request monitoring from the MnDNR to assure beetles are present and at appropriate populations for biological control.	Increase/maintain wildlife habitat.	Spring - Fall	
Hybrid cattail and reed canary grass are present.	Control hybrid cattail and reed canary grass.	Control hybrid cattail and reed canary grass now before colonies become more abundant. The herbicide Rodeo™ can be used to effectively control both invasive emergent species.	Increase/maintain wildlife habitat.	Spring-Summer	
Stormwater drainage from impervious surfaces is directed into the lake.	Redirect stormwater for infiltration prior to discharge.	Install a rainwater garden or other suitable method for infiltration.	Improve water quality	Open	
Bare soil on steep slope could cause erosion and sedimentation into lake.	Vegetate hillslope.	Plant vegetation suited for steep slopes along hillside to prevent erosion.	Improve water quality	Open	
Upland buffer areas lacking naturalized vegetation.	Improve the shoreline with a naturalized upland buffer.	Rather than manicured turf grass, gravel, and managed plantings with bare soil, the shoreline could be vegetated with native grasses and wildflowers. A landscape architect could create inviting spaces and views for restaurant customers to enjoy.	Increase wildlife habitat and Improve water quality	Open	
Emergent and upland buffer areas contain non-native invasive vegetation.	Replace non-native invasive vegetation with native vegetation.	Treat non-native invasive vegetation and then seed with an appropriate BWSR seed mix.	Increase/maintain wildlife habitat.	Spring-Summer	

**Table 2: 2012 Recommended and Completed Management Actions for Orchard Lake
Black Dog Watershed Management Organization Habitat Monitoring**

Problem Identified	Recommendation	Proposed Action	Benefits	Implementation Period	Completed 2004-2012 Actions Which May Improve Wildlife Habitat and/or Water Quality
Curlyleaf pondweed dominates the lake in late spring-early summer.	Continue curlyleaf pondweed control measures.	Continue to control and manage. See Figure 3 for locations of curlyleaf pondweed.	Increase wildlife habitat, improve water quality, vegetative diversity, aesthetics, and recreation.	Late Spring - Early summer	<p>1999 through 2012: The City of Lakeville conducts aquatic vegetation monitoring twice/year.</p> <p>2009 through 2012: The City of Lakeville conducted annual herbicide treatment for curlyleaf pondweed.</p> <p>2004 through 2008: Annually, the City of Lakeville harvested approximately 70 acres of curlyleaf pondweed.</p> <p>2010: Adjacent to the southwest end of the lake, an aeration system was installed in Orchard Pond to precipitate out phosphorus and improve water quality flowing into Orchard Lake.</p> <p>2004 through 2012: The City of Lakeville annually provides lakeshore owners with shoreline restoration information and encourages homeowners to take advantage of the Blue Thumb restoration program.</p> <p>One shoreline resident started a restoration project in 2012.</p> <p>2007: A small area of lakeshore, near the boat launch, was restored using native plants.</p> <p>2002: Purple loosestrife beetles were released by the MNDNR. Follow up monitoring indicates that beetles are present at a population that the MNDNR feels is appropriate for biological control of purple loosestrife plants.</p>
Upland buffer areas lacking naturalized vegetation within publicly owned properties.	Increase width and continuity of native upland buffer.	<p>To expand on the shoreline restoration that was done near the boat launch in 2007, the adjacent upland buffer could also be restored to naturalized native vegetation and not mowed (Potential Restoration Area #1 as shown in Appendix A and Figure 5).</p> <p>In the Wayside Park Area, non-native invasive vegetation including common buckthorn, vetch, spotted knapweed, and cattails could be removed and replaced with native vegetation. The naturalized upland buffer could be widened (Potential Restoration Area #2 as shown in Appendix A and Figure 5).</p> <p>At the beach area, there is a timber wall which is currently being used for fishing. A shoreline restoration could be done in this area (Potential Restoration Area #3 as shown in Appendix A and Figure 5).</p> <p>On the northwest side of the lake, one property owned by the City of Lakeville (adjacent to residential shoreline properties) could be restored to naturalized vegetation and provide an example for adjacent residential landowners for shoreline and upland buffer restoration (Potential Restoration Area #4 as shown in Appendix A and Figure 5).</p>	Increase wildlife habitat. Improve water quality. Improve vegetative diversity and aesthetics.	Spring - Fall	
Upland buffer areas lacking naturalized vegetation. Most of the residential properties have turf grass up the the lakeshore edge.	Increase width and continuity of native upland buffer.	<p>Restore sustainable native communities. Rather than manicured turf grass, sand, and bare soil, the shoreline could be vegetated with native grasses and wildflowers. A native upland buffer can improve functions and values of the lake and improve aesthetics (Potential Restoration Area #5 as shown in Appendix A and Figure 5).</p>	Increase wildlife habitat. Improve water quality. Improve vegetative diversity and aesthetics.	Spring - Fall	
Purple loosestrife is present.	Continue to control and manage purple loosestrife.	Control and manage. For a few small colonies of purple loosestrife, hand pull or dig the plants out before they go to seed. Continue to request monitoring from the MNDNR to assure beetles are present and at appropriate populations for biological control (See Figures 3 and 5 for location of purple loosestrife).	Increase wildlife habitat. Improve vegetative diversity.	Spring - Fall	

**Table 2: 2013 Recommended and Completed Management Actions for Crystal Lake
Black Dog Watershed Management Organization Habitat Monitoring**

Problem Identified	Recommendation	Proposed Action	Benefits	Implementation Period	Completed Actions Which May Improve Wildlife Habitat and/or Water Quality
Curlyleaf pondweed dominates the lake in late spring-early summer.	Continue curlyleaf pondweed control measures.	Continue to control and manage. See Blue Water Science report for locations of curlyleaf pondweed.	Increase wildlife habitat, improve water quality, vegetative diversity, aesthetics, and recreation.	Late Spring - Early summer	
Upland buffer areas lacking naturalized vegetation within publicly owned properties.	Increase width and continuity of native upland buffer.	The width and density of naturalized shoreline buffer at the location of Emergent Plot #1 near the swimming area has improved significantly since 2009. The adjacent upland buffer could also be restored to naturalized native vegetation and not mowed (Potential Restoration Areas #1 through 4 as shown in Figure 4 and photos).	Increase wildlife habitat. Improve water quality. Improve vegetative diversity and aesthetics.	Spring - Fall	<p>1999 through 2013: The City of Burnsville conducts aquatic vegetation monitoring twice/year.</p> <p>2003 through 2013: The City of Burnsville conducted annual harvesting of curlyleaf pondweed.</p> <p>2004-2008:</p> <ul style="list-style-type: none"> -The BDWMO operated the ferric chloride treatment system.
Upland buffer areas lacking naturalized vegetation. Most of the residential properties have turf grass up the the lakeshore edge.	Increase width and continuity of native upland buffer.	Restore sustainable native communities. Rather than manicured turf grass, sand, and bare soil, the shoreline could be vegetated with native grasses and wildflowers. A native upland buffer can improve functions and values of the lake and improve aesthetics (Potential Restoration Area #5 as shown in Figure 4 and photos).	Increase wildlife habitat. Improve water quality. Improve vegetative diversity and aesthetics.	Spring - Fall	<ul style="list-style-type: none"> -The City of Burnsville: 1) excavated/enhanced four stormwater treatment ponds (including West Buck Hill Park), which reduced the phosphorus loading into the lake, and 2) conducted annual harvesting of Eurasian watermilfoil and curlyleaf pondweed.
Purple loosestrife is present.	Continue to control and manage purple loosestrife.	Continue to control. For a few small colonies of purple loosestrife, hand pull or dig the plants out before they go to seed.	Increase wildlife habitat. Improve vegetative diversity.	Spring - Fall	<ul style="list-style-type: none"> -The City of Lakeville excavated/enhanced the Bluebill stormwater treatment pond. <p>In 2009 and 2008, garlic mustard within the upland buffer was removed/pulled.</p> <p>In late 2009, the City of Burnsville treated 14 acres of buckthorn within Crystal West Park.</p>
Eurasian watermilfoil is present.	Control Eurasian watermilfoil.	Control by chemical treatment.	Maintain wildlife habitat.	Summer	

**Table 2: 2014 Recommended and Completed Management Actions for Lac Lavon
Black Dog Watershed Management Organization Habitat Monitoring**

Problem Identified	Recommendation	Proposed Action	Benefits	Implementation Period	Completed Actions Which May Improve Wildlife Habitat and/or Water Quality
Curlyleaf pondweed dominates the lake in late spring-early summer.	Continue curlyleaf pondweed control measures.	Continue to control and manage. See Macrophyte Survey Results for locations of curlyleaf pondweed.	Increase wildlife habitat, improve water quality, vegetative diversity, aesthetics, and recreation.	Late Spring - Early summer	Aquatic plant surveys were conducted by Barr in 2013 and 2014.
Eurasian watermilfoil is present.	Control Eurasian watermilfoil.	Control by chemical treatment. See Macrophyte Survey Results for locations of Eurasian watermilfoil	Maintain wildlife habitat.	Summer	In 2006, the cities of Burnsville and Apple Valley and the lake homeowners partnered to fund a fluridone treatment for control of Eurasian watermilfoil. Aquatic plant surveys were conducted by Barr in 2013 and 2014.
Purple loosestrife is present.	Continue to control and manage purple loosestrife.	Continue to control. For a few small colonies of purple loosestrife, hand pull or dig the plants out before they go to seed. See Macrophyte Survey Results for locations of purple loosestrife	Increase wildlife habitat. Improve vegetative diversity.	Spring - Fall	Purple loosestrife removal on shallow island areas was completed by the cities of Apple Valley and Burnsville in 2011.
Upland buffer areas lacking naturalized vegetation within publicly owned properties.	Increase width and continuity of native upland buffer.	Expand native prairie planting to include area to the east, which is dominated by knapweed. This could become a tall grass prairie. Potential Restoration Area #1	Increase wildlife habitat. Improve water quality. Improve vegetative diversity and aesthetics.	Spring - Fall	In 2013, the city of Burnsville installed a native prairie planting converting a sand beach and turf grass to prairie and wetland vegetation.
Upland buffer areas in city parks contain non-native invasive vegetation such as buckthorn, Siberian elm, leafy spurge, and spotted knapweed.	Continue to control and manage non-native invasive vegetation	Continue to control and manage non-native invasive vegetation Potential Restoration Area #2	Increase wildlife habitat. Improve vegetative diversity and aesthetics	Spring - Fall	Some invasive species control for Canada thistle and knapweed was conducted on the new native planting area in 2014. In 2010, the city of Apple Valley released about 150 spotted knapweed seedhead boring weevils in Lac Lavon Park in Apple Valley. Continued management of the vegetation communities and shoreline restoration activities will help to maintain and improve wildlife habitat, vegetation diversity, aesthetics, and recreation
Impervious surfaces and turf grass in the Apple Valley park near the fishing pier can collect pollutants in stormwater and flow directly into the lake, decreasing water quality.	Increase areas of naturalized vegetation to slow down and pretreat stormwater prior to entering the lake.	Strategically create buffer strips with naturalized vegetation adjacent to impervious surfaces to slow down and pretreat stormwater prior to entering the lake. Potential Restoration Area #3	Improve water quality	Spring - Fall	
Upland buffer areas lacking naturalized vegetation. Most of the residential properties have turf grass or sand up to the lakeshore edge.	Increase width and continuity of native upland buffer.	Restore sustainable native communities. Rather than manicured turf grass, sand, and bare soil, the shoreline could be vegetated with native grasses and wildflowers. A native upland buffer can improve functions and values of the lake and improve aesthetics. Potential Restoration Area #4	Increase wildlife habitat. Improve water quality. Improve vegetative diversity and aesthetics.	Spring - Fall	One raingarden was installed in the backyard of a shoreline property owner on Highview Drive in Apple Valley through the Blue Thumb program. The establishment of shoreline restoration projects (especially contiguous) on residential properties in the future will help balance out the differences in upland buffer habitat between city owned property and residential property.

**Table 2: 2015 Recommended and Completed Management Actions for Keller Lake
Black Dog Watershed Management Organization Habitat Monitoring**

Problem Identified	Recommendation	Proposed Action	Benefits	Implementation Period	Completed Actions Which May Improve Wildlife Habitat and/or Water Quality
Curlyleaf pondweed dominates the lake in late spring-early summer.	Continue curlyleaf pondweed control measures.	Continue to control and manage. See Appendix A Aquatic Plant Survey for locations of curlyleaf pondweed.	Increase wildlife habitat, improve water quality, vegetative diversity, aesthetics, and recreation.	Late Spring - Early summer	Aquatic plant surveys have been conducted by Blue Water Science 1998-2015. Iron dosing occurred from 1999 through 2008. Mechanical harvesting is conducted each year since 2004.
Eurasian watermilfoil is present.	Control Eurasian watermilfoil.	Continue to monitor. Control as recommended by the MnDNR. Since the MnDNR designates Keller Lake as a "Natural Environment Lake", chemical treatment is not allowed.	Maintain wildlife habitat.	Summer	Aquatic plant surveys have been conducted by Blue Water Science 1998-2015.
The inlet coming from the stormwater pond at the south end of Keller Lake is surrounded by bare soil or sparse vegetation.	Re-vegetated bare areas to prevent soil erosion into Keller Lake.	Seed or plant bare areas with native vegetation. Potential Restoration Area #1	Improve water quality and vegetative diversity.	Spring or Fall	
Shoreline fishing traffic is causing bare soil areas along the shoreline.	Re-vegetated bare areas to prevent soil erosion into Keller Lake.	Create designated stone walkways for fishing access. Potential Restoration Area #2	Improve water quality, vegetative diversity, and aesthetics.	Spring - Fall	
The southern public park is littered with trash and other dumped items especially near the shoreline.	Clean up the litter.	Organize a neighborhood clean-up project to pick up trash and other dumped items along the south shoreline of the lake. Potential Restoration Area #3	Improve aesthetics. Potentially prevent harm to wildlife. Prevent migration of trash into lake.	Spring - Fall	
Upland buffer areas in city parks contain non-native invasive vegetation such as buckthorn and garlic mustard.	Continue to control and manage non-native invasive vegetation	Continue to control and manage non-native invasive vegetation. Pull garlic mustard within the City of Burnsville property at the north end of the lake. Continue to remove and treat new growth of buckthorn in city parks. Potential Restoration Area #4	Increase wildlife habitat. Improve vegetative diversity and aesthetics	Spring - Fall	Buckthorn appears to have been previously removed in the park along the southern shoreline.
Upland buffer areas lacking naturalized vegetation. Some of the residential properties have narrow buffers with turf grass close to the lakeshore edge.	Increase width and continuity of native upland buffer.	Restore sustainable native communities. Manicured turf grass near the shoreline could be vegetated with native grasses and wildflowers. A native upland buffer can improve functions and values of the lake and improve aesthetics. Potential Restoration Area #5	Increase wildlife habitat. Improve water quality. Improve vegetative diversity and aesthetics.	Spring - Fall	Most residential properties allow a narrow width of naturalized vegetation to prevent soil erosion, however a wider buffer of native vegetation could help improve wildlife habitat, vegetative diversity, and aesthetics.

**Table 2: 2016 Recommended and Completed Management Actions for Kingsley Lake
Black Dog Watershed Management Organization Habitat Monitoring**

Problem Identified	Recommendation	Proposed Action	Benefits	Implementation Period	Completed Actions Which May Improve Wildlife Habitat and/or Water Quality
Curlyleaf pondweed is present in some years.	Continue to monitor	Consider control measures, if densities and locations increase to an extent of concern. See Appendix A Aquatic Plant Survey for locations of curlyleaf pondweed.	Increase wildlife habitat, improve water quality, vegetative diversity, aesthetics, and recreation.	Late Spring - Early summer	On March 6, 2008, soil sediment samples were collected on Kingsley Lake by Blue Water Science (BWS) and the City of Lakeville. Based on the results of the soil analysis, the BWS report stated that "curlyleaf pondweed is not expected to produce heavy growth conditions (where plants top out in a solid canopy) in Kingsley Lake."
Common buckthorn dominates portions of the upland buffer.	Conduct an evaluation of common buckthorn, followed by removal.	Remove buckthorn. Volunteer groups and contractors can effectively remove buckthorn by pulling, cutting, and treating stumps with herbicide. See Figure 4, Potential Restoration Area #1	Increase wildlife habitat. Improve vegetative diversity and aesthetics	Spring - Fall	From 2005-2008, the City of Lakeville and members of the Kingsley Lake Association removed common buckthorn from portions of the lake and the upland buffer surrounding the lake.
Purple loosestrife is present.	Continue to control and manage purple loosestrife.	Control and manage. For a few small colonies of purple loosestrife, hand pull or dig the plants out before they go to seed. See Figure 4 for purple loosestrife locations.	Increase/maintain wildlife habitat.	Spring - Fall	From 2005-2008, the City of Lakeville and members of the Kingsley Lake Association removed purple loosestrife plants from portions of the lake and the upland buffer surrounding the lake. Purple loosestrife beetles were released by the MnDNR prior to 2002. Follow up monitoring by the MnDNR indicates that beetles are present at a population that the MnDNR feels is appropriate for biological control.
Stormwater drainage from impervious surfaces is directed into the lake.	Redirect stormwater for infiltration prior to discharge.	Install a rainwater garden, pervious pavement, or other suitable method for infiltration. See Figure 4, Potential Restoration Area #2.	Improve water quality	Open	
Bare soil on steep slope could cause erosion and sedimentation into lake.	Re-vegetate bare areas to prevent soil erosion into Kingsley Lake.	Plant vegetation suited for steep slopes along hillside to prevent erosion. See Figure 4, Potential Restoration Area #3	Improve water quality	Spring - Fall	
Upland buffer areas lacking naturalized vegetation.	Increase width and continuity of native upland buffer.	Rather than manicured turf grass, gravel, and managed plantings with bare soil, the shoreline could be vegetated with native grasses and wildflowers. See Figure 4, Potential Restoration Areas #4 through 7. See Appendix G for examples of improvements. See also island shoreline areas becoming bare from YMCA camper overuse (Figure 4, Potential Restoration Areas 10 and 11).	Improve water quality, increase wildlife habitat. Improve vegetative diversity and aesthetics.	Spring - Fall	In 2008, a Kingsley Lake lakeshore resident, inspired by the Blue Thumb program, commenced shoreline stabilization utilizing native plants.
Emergent zone and upland buffer areas contain non-native invasive vegetation.	Continue to control and manage non-native invasive vegetation, including, but not limited to reed canary grass, hybrid cattail, and yellow iris.	Treat non-native invasive vegetation and then seed with an appropriate BWSR seed mix. See Figure 4, Potential Restoration Areas #8 and 9. Remove yellow iris (See Appendix A for locations of yellow iris). The MN DNR may require a permit for cattail treatment and yellow iris removal if below the OHW. Dense reed canary grass is located at Plot 2b as shown of Figure 3 . Dense hybrid cattail is located at Plot 1b as shown on Figure 3 .	Increase/maintain wildlife habitat.	Spring-Summer	

**Table 2: 2017 Recommended and Completed Management Actions for Orchard Lake
Black Dog Watershed Management Organization Habitat Monitoring**

Problem Identified	Recommendation	Proposed Action	Benefits	Implementation Period	Completed Actions Which May Improve Wildlife Habitat and/or Water Quality
Curlyleaf pondweed is common in early spring	Continue to monitor, control, and manage.	Continue to treat curlyleaf pondweed where growth is predicted to be heavy. See Appendix A Aquatic Plant Survey for more details.	Increase wildlife habitat, improve water quality, vegetative diversity, aesthetics, and recreation.	Late Spring - Early summer	From 1999-2017, the City of Lakeville contracts Blue Water Science to conduct aquatic plant surveys twice per year. Curlyleaf pondweed was harvested annually from 2004-2009. Herbicide treatments were conducted annually from 2009-2012 and 2015-2017.
Purple loosestrife is present.	Continue to control and manage purple loosestrife.	Control and manage. For a few small colonies of purple loosestrife, hand pull or dig the plants out before they go to seed. See Figure 4 for purple loosestrife locations.	Increase/maintain wildlife habitat.	Spring - Fall	Purple loosestrife beetles were released by the MnDNR prior to 2002. Follow up monitoring by the MnDNR indicates that beetles are present at a population that the MnDNR feels is appropriate for biological control.
Stormwater drainage from impervious surfaces is directed into the lake.	Redirect stormwater for infiltration prior to discharge.	Install a rainwater garden, pervious pavement, or other suitable method for infiltration and establish a naturalized upland buffer. See Figure 4 and Site Photos, Potential Restoration Area #6.	Improve water quality	Open	Two raingardens were completed on 175th St W. In 2010, adjacent to the southwest end of the lake, an aeration system was installed in Orchard Pond to precipitate out phosphorus and improve water quality flowing into Orchard Lake.
Bare soil along shoreline could cause erosion and sedimentation into lake.	Re-vegetate bare areas to prevent soil erosion into Orchard Lake.	Improve soil and plant vegetation along shoreline to prevent erosion. Establish a canoe and kayak access at Wayside Park. See Figure 4 and Site Photos, Potential Restoration Area #4 and #5.	Improve water quality	Spring - Fall	The City of Lakeville removed a dilapidated timber wall and attempted a shoreline restoration south of the beach, however, the soil was too poor for the plantings to become established. North of the beach, a concrete wall was built to prevent shoreline erosion.
Upland buffer areas lacking naturalized vegetation.	Increase width and continuity of native upland buffer.	Rather than manicured turf grass the shoreline could be vegetated with native grasses and wildflowers. See Figure 4 and Site Photos, Potential Restoration Areas #1-3, 7 and 8. See Appendix G for examples of improvements.	Improve water quality, increase wildlife habitat. Improve vegetative diversity and aesthetics.	Spring - Fall	2004 through 2012: The City of Lakeville annually provides lakeshore owners with shoreline restoration information and encourages homeowners to take advantage of the Blue Thumb restoration program. Two residential shoreline restoration projects have been completed. One is located north of the beach area and one is on 175th St. W. 2007: A small area of lakeshore, near the boat launch, was restored using native plants.

**Table 2: 2018 Recommended and Completed Management Actions for Crystal Lake
Black Dog Watershed Management Organization Habitat Monitoring**

Problem Identified	Recommendation	Proposed Action	Benefits	Implementation Period	Completed Actions Which May Improve Wildlife Habitat and/or Water Quality
Curlyleaf pondweed dominates the lake in late spring-early summer.	Continue curlyleaf pondweed control measures.	Continue to control and manage. See Appendix A Blue Water Science report for locations of curlyleaf pondweed.	Increase wildlife habitat, improve water quality, vegetative diversity, aesthetics, and recreation.	Late Spring - Early summer	1999 through 2018: The City of Burnsville conducts aquatic vegetation monitoring twice/year. 2003 through 2018: The City of Burnsville conducted annual harvesting of curlyleaf pondweed.
Eurasian watermilfoil is present.	Control Eurasian watermilfoil.	Control by chemical treatment. See Appendix A Blue Water Science report for locations of Eurasian watermilfoil.	Increase wildlife habitat, improve water quality, vegetative diversity, aesthetics, and recreation.	Summer	
Common and glossy buckthorn are present	Control common and glossy buckthorn	Remove buckthorn. Volunteer groups and contractors can effectively remove buckthorn by pulling, cutting, and treating stumps with herbicide. See Appendix H for buckthorn management guidelines. See Appendix I for locations of buckthorn.	Increase wildlife habitat. Improve vegetative diversity and aesthetics	Fall	In 2009, the City of Burnsville treated 14 acres of buckthorn within Crystal Lake West Park (Appendix I).
Garlic mustard is present	Control garlic mustard	Organize a volunteer neighborhood group to pull garlic mustard. See Appendix I for locations of garlic mustard.	Increase wildlife habitat. Improve vegetative diversity and aesthetics	Spring	In 2008 and 2009, the City of Burnsville removed garlic mustard within the upland buffer (Appendix I)
Purple loosestrife is present.	Continue to control and manage purple loosestrife.	Control and manage. For a few small colonies of purple loosestrife, hand pull or dig the plants out before they go to seed. See Appendix I for locations of purple loosestrife.	Increase wildlife habitat. Improve vegetative diversity.	Spring - Fall	Purple loosestrife beetles were released by the MnDNR prior to 2002. Follow up monitoring by the MnDNR indicates that beetles are present at a population that the MnDNR feels is appropriate for biological control.
Bare soil areas are present along shoreline in Crystal Lake West Park area.	Re-vegetate bare soil areas to prevent soil erosion into Crystal Lake and create designated stone walkways for fishing access.	Exposed soil along the shoreline of Crystal Lake West Park Area could be re-vegetated to prevent shoreline erosion. Strategically located stones could provide fishing access to prevent disturbance of vegetation after it is established. (Potential Restoration Area #1 as shown in Figure 4 and photos)	Improve water quality and prevent erosion.	Spring - Fall	
Timber retaining wall in Tyecke Park area is in poor condition.	Repair timber retaining wall to prevent soil erosion into Crystal Lake.	Steep slopes in the Tyecke Park area are well protected with mature naturalized vegetation, however a timber retaining wall along the shoreline may need to be repaired or replaced to prevent slope destabilization and erosion. (Potential Restoration Area #2 as shown in Figure 4 and photos)	Prevent erosion	Winter	
Shoreline areas lacking naturalized vegetation within publicly owned beach area. Some areas have mowed turf grass close to the lakeshore edge.	Increase width and continuity of native upland buffer.	The upland buffer near the location of Plot #1C and shoreline to the south, and north of the beach area could be restored to naturalized native vegetation and not mowed (Potential Restoration Areas #3 and 4 as shown in Figure 4 and photos).	Increase wildlife habitat. Improve water quality. Improve vegetative diversity and aesthetics.	Spring - Fall	The width and density of naturalized shoreline buffer at the location of Emergent Plot #1B near the beach area has improved significantly since 2009.
Shoreline areas lacking naturalized vegetation within residential properties. Most of the residential properties have turf grass up the the lakeshore edge.	Increase width and continuity of native upland buffer.	Rather than manicured turf grass, the shoreline could be vegetated with native grasses and wildflowers. (Potential Restoration Area #5 as shown in Figure 4 and photos).	Increase wildlife habitat. Improve water quality. Improve vegetative diversity and aesthetics.	Spring - Fall	Six residential property owners have completed shoreline restoration projects using either City of Burnsville or Dakota Soil and Water Conservation District grants.

Table 2 2019 Recommended and Completed Management Actions for Lac Lavon – Black Dog Watershed Management Organization Habitat Monitoring

Problem Identified	Recommendation	Proposed Action	Benefits	Implementation Period	Completed Actions Which May Improve Wildlife Habitat and/or Water Quality
Curly-leaf pondweed dominates the lake in late spring-early summer.	Continue curly-leaf pondweed control measures.	Continue to control and manage. Detailed results are available upon request.	Increase wildlife habitat, improve water quality, vegetative diversity, aesthetics, and recreation.	Late Spring - Early summer	Aquatic plant surveys were conducted in 2013, 2014, and 2019.
Eurasian watermilfoil is present.	Control Eurasian watermilfoil.	Control by chemical treatment. Detailed results are available upon request.	Increase wildlife habitat, improve water quality, vegetative diversity, aesthetics, and recreation.	Summer	In 2006, the cities of Burnsville and Apple Valley and the lake homeowners partnered to fund a one-time fluridone treatment for control of Eurasian watermilfoil. Aquatic plant surveys were conducted in 2013, 2014, and 2019.
Purple loosestrife is present.	Continue to control and manage purple loosestrife.	Continue to control. For a few small colonies of purple loosestrife, hand pull or dig the plants out before they go to seed.	Increase wildlife habitat. Improve vegetative diversity.	Spring - Fall	Purple loosestrife removal on shallow island areas was completed by the cities of Apple Valley and Burnsville in 2011. Purple loosestrife beetles were released by the MnDNR prior to 2002. Follow up monitoring by the MnDNR indicates that beetles are present at a population that the MnDNR feels is appropriate for biological control.
Shoreline areas lacking naturalized vegetation within publicly owned properties.	Increase width and continuity of native upland buffer.	Expand native prairie planting to include area to the east, which is dominated by knapweed. This could become a tall grass prairie. Potential Restoration Area #1	Increase wildlife habitat. Improve water quality. Improve vegetative diversity and aesthetics.	Spring - Fall	In 2013, the city of Burnsville installed a native prairie planting converting a sand beach and turf grass to prairie and wetland vegetation.
Shoreline areas in city parks contain non-native invasive vegetation such as buckthorn, Siberian elm, leafy spurge, and spotted knapweed.	Continue to control and manage non-native invasive vegetation	Continue to control and manage non-native invasive vegetation Potential Restoration Area #2	Increase wildlife habitat. Improve vegetative diversity and aesthetics	Spring - Fall	Some invasive species control for Canada thistle and knapweed was conducted on the new native planting area in 2014. In 2010, the city of Apple Valley released about 150 spotted knapweed seedhead boring weevils in Lac Lavon Park in Apple Valley. Continued management of the vegetation communities and shoreline restoration activities will help to maintain and improve wildlife habitat, vegetation diversity, aesthetics, and recreation
Impervious surfaces and turf grass in the Apple Valley park near the fishing pier can collect pollutants in stormwater and flow directly into the lake, decreasing water quality.	Increase areas of naturalized vegetation adjacent to impervious surfaces to slow down and pretreat stormwater prior to entering the lake.	Strategically create buffer strips of naturalized vegetation adjacent to the bituminous lake access pathway to slow down and pretreat stormwater prior to entering the lake. Potential Restoration Area #3	Improve water quality	Spring - Fall	
Upland buffer areas lacking naturalized vegetation. Most of the residential properties have turf grass or sand up to the lakeshore edge.	Increase width and continuity of native upland buffer.	Restore sustainable native communities. Rather than manicured turf grass, sand, and bare soil, the shoreline could be vegetated with native grasses and wildflowers. A native upland buffer can improve functions and values of the lake and improve aesthetics. Potential Restoration Area #4	Increase wildlife habitat. Improve water quality. Improve vegetative diversity and aesthetics.	Spring - Fall	One native prairie restoration project was installed in the backyard of a shoreline property owner on Highview Drive in Apple Valley through the Dakota Soil and Water Conservation District program. The establishment of shoreline restoration projects (especially contiguous) on residential properties in the future will help balance out the differences in upland buffer habitat between city owned property and residential property.

Table 2 2020 Recommended and Completed Management Actions for Keller Lake – Black Dog Watershed Management Organization Habitat Monitoring

Problem Identified	Recommendation	Proposed Action	Benefits	Implementation Period	Completed Actions Which May Improve Wildlife Habitat and/or Water Quality
Curly-leaf pondweed dominates the lake in late spring-early summer.	Continue curly-leaf pondweed control measures.	Continue to control and manage. See Appendix A for details.	Increase wildlife habitat, improve water quality, vegetative diversity, aesthetics, and recreation.	Late Spring - Early summer	Aquatic plant surveys have been conducted by Blue Water Science since 1998. A curly-leaf pondweed turion survey was also conducted in 2020. Iron dosing occurred during 1996 - 2008. Mechanical harvesting was conducted from 2004 - 2015. Herbicide treatments were conducted from 2017 – 2020.
Eurasian watermilfoil is present.	Control Eurasian watermilfoil.	Continue to control and manage. See Appendix A for details.	Increase wildlife habitat, improve water quality, vegetative diversity, aesthetics, and recreation.	Summer	Aquatic plant surveys have been conducted by Blue Water Science since 1998.
Low native aquatic vegetative diversity in the submergent zone.	Continue to increase native aquatic plant diversity.	Continue to monitor and assess. See Appendix A for details.	Increase wildlife habitat, improve water quality, vegetative diversity, aesthetics, and recreation.	Summer	A seedbank assessment was conducted in 2020. Native aquatic plant re-introduction began in 2020.
The inlet coming from the stormwater pond at the south end of Keller Lake is surrounded by bare soil or sparse vegetation.	Re-vegetate bare areas to prevent soil erosion into Keller Lake.	Seed or plant bare areas with native vegetation. Potential Restoration Area #1	Improve water quality and vegetative diversity.	Spring or Fall	
Shoreline pedestrian observation and fishing traffic is causing bare soil areas along the shoreline.	Re-vegetate bare areas to prevent soil erosion into Keller Lake.	Create designated stone walkways for observation and fishing access. Potential Restoration Area #2	Improve water quality, vegetative diversity, and aesthetics.	Spring-Fall	
The southern public park is littered with trash and other dumped items, especially near the shoreline.	Clean up the litter.	Organize a neighborhood clean-up project to pick up trash and other dumped items along the south shoreline of the lake. Potential Restoration Area #3	Improve aesthetics. Potentially prevent harm to wildlife. Prevent migration of trash into lake.	Spring - Fall	
Shoreline areas in city parks contain non-native invasive vegetation such as buckthorn, spotted knapweed, and garlic mustard.	Continue to control and manage non-native invasive vegetation	Continue to control and manage non-native invasive vegetation. Pull garlic mustard within the City of Burnsville property at the north end of the lake. Continue to remove and treat new growth of buckthorn in city parks. Potential Restoration Area #4	Increase wildlife habitat. Improve vegetative diversity and aesthetics	Spring - Fall	Buckthorn appears to have been previously removed in the park along the southern shoreline, however new seedlings are emerging. Continued management of the vegetation communities and shoreline restoration activities will help to maintain and improve wildlife habitat, vegetation diversity, aesthetics, and recreation
Upland buffer areas lacking naturalized vegetation. Many of the residential properties have narrow buffers with lawns mowed to the lakeshore edge.	Increase width and continuity of native upland buffer.	Restore sustainable native communities. Rather than manicured turf grass, the shoreline could be vegetated with native grasses and wildflowers. A native upland buffer can improve functions and values of the lake and improve aesthetics. Potential Restoration Area #5	Increase wildlife habitat. Improve water quality. Improve vegetative diversity and aesthetics.	Spring - Fall	The Dakota Soil and Water Conservation District program assists homeowners with establishment of shoreline restoration projects. Additional restoration projects (especially contiguous) on residential properties in the future will help balance out the differences in upland buffer habitat between city owned property and residential property.
Shoreline areas lacking naturalized vegetation within publicly owned properties.	Increase width and continuity of native upland buffer.	Adjust mowing distance away from shoreline in Apple Valley's Keller Park. Potential Restoration Area #6	Increase wildlife habitat. Improve water quality. Improve vegetative diversity and aesthetics.	Spring - Fall	
A portion of Lac Lavon Park in Burnsville south of Keller Lake includes a large area of mowed turf grass with no apparent use.	Consider using this area for a native prairie restoration with meandering trails and educational signs.	Recruit volunteers through neighborhood or organizations to transform the lawn into a prairie. Potential Restoration Area #7	Improve vegetative diversity and aesthetics. Increase wildlife habitat and provide pollinator habitat. Provide recreational and educational opportunities.	Spring - Fall	
Sediment from the Lac Lavon Park parking lot is directed into the floodplain forest along the south shoreline of Keller Lake.	Prevent sediment from entering the floodplain forest area.	Install a pre-treatment system such as a rain garden or sediment trap to collect sediment from the parking lot. Follow up with routine maintenance of sediment clean-out. Potential Restoration Area #8	Protect floodplain forest from further degradation and prevent sediment from reaching Keller Lake, thereby improving water quality.	Winter	
Japanese hedge parsley is located along the edge of the path near the storm pond north of Keller Lake. This is a new non-native invasive species in Minnesota. Early detection and control will prevent infestations.	Eradicate- Pull, cut or mow before flowering. Monitor area for new seedlings. Spray rosettes or bolting plants in spring with 1-2% glyphosate or 1-2% triclopyr. In fall, use herbicides on rosettes.	A report of this species at this location was submitted through the Great Lakes Early Detection Network online application. Follow up with the Dakota County Cooperative Weed Management contact to verify action will be taken for eradication. Potential Restoration Area #9	Prevent spread and infestation.	Spring - Fall	

Appendix E

2016 Kingsley Lake MNRAM 3.4 Wetland Functional Assessment Results

Wetland Functional Assessment Summary

Wetland Name	Hydrogeomorphology	Maintenance of Hydrologic Regime	Flood/Stormwater/Attenuation	Downstream Water Quality	Maintenance of Wetland Water Quality	Shoreline Protection
Kingsley Lake	Depressional/Flow-through (apparent inlet and outlet), Depressional/Flow-through (apparent inlet and outlet)	0.75	0.58	0.63	0.66	0.64
		High	Moderate	Moderate	High	Moderate

Additional Information

Wetland Name	Maintenance of Characteristic Wildlife Habitat Structure	Maintenance of Characteristic Fish Habitat	Maintenance of Characteristic Amphibian Habitat	Aesthetics/Recreation/Education/Cultural	Commercial Uses	Ground-Water Interaction	Wetland Restoration Potential	Wetland Sensitivity to Stormwater and Urban Development	Additional Stormwater Treatment Needs
Kingsley Lake	0.70	0.80	0.06	0.59	0.00	Discharge	0.00	1.00	0.66
	High	High	Low	Moderate	Not Applicable		Not Applicable	High	High

Wetland Community Summary

Wetland Name	Location	Vegetative Diversity/Integrity							
		Community			Wetland Proportion	Individual Community Rating	Highest Wetland Rating	Average Wetland Rating	Weighted Average Wetland Rating
		Cowardin Classification	Circular 39	Plant Community					
Kingsley Lake	19-114-21-11-001	L2AB2G	Type 5	Shallow, Open Water Communities	70	1	1.00	1.00	1.00
							High	High	High
		PEM1F	Type 4	Deep Marsh	30	1	1.00	1.00	1.00
							High	High	High
					100		1.00	1.00	1.00

Denotes incomplete calculation data.

Appendix F

Descriptions of MNRAM Wetland Functions

6.0 Functional Rating Formulas

GENERAL NOTE: Some questions are not applicable to particular wetlands and will be scored N/A. In these cases, rather than count N/A as zero, an alternate equation is provided that eliminates the question from the formula altogether. Because not every question has N/A as an option, formulas that do not include N/A-option questions have only one configuration.

Formulas with a “reverse rating” (marked as “R”) take the actual response and “flip” its value for the calculation, so that a question response of “A” high (value of 1.0) will be calculated as low (value of 0.1). In such a formula, medium ratings stay medium.

6.1 VEGETATIVE DIVERSITY/INTEGRITY

Table 3: Vegetative Diversity/Integrity Summary

The functional rating is based primarily on the diversity of vegetation within the wetland in comparison to an undisturbed condition for that wetland type. An exceptional rating results from one of the following conditions: 1) highly diverse wetlands with virtually no non-native species, 2) rare or critically impaired wetland communities in the watershed, or 3) the presence or previous siting of rare, threatened, or endangered plant species. A high rating indicates the presence of diverse, native wetland species and a lack of non-native or invasive species. Wetlands that rate low are primarily dominated by non-native and/or invasive species.

This table may be used when calculating Vegetative Diversity/Integrity Functional Index manually. It shows four options for calculating and presenting floristic data. If you are entering data directly into the MnRAM 3.0 database, this table does not apply.

	3A Proportion of Wetland	3B Individual Community Scores	3C Highest Quality	3D Non-Weighted Average	3E Weighted Average
Community #1	T	A		A	A
Community #2	U	B		B	B
Community #3	V	C		C	C
Community #4	W	D		D	D
Community #5	X	E		E	E
Community #6	Y	F		F	F
Community #7	Z	G		G	G
Wetland Rating Value	1.0		Highest Value	$(A+B+C+D+E+F+G)/7 =$ Ave.	$(A*T)+(B*U$ $)+(C*V)+(D$ $*W)+(E*X)+$ $(F*Y)+(G*Z$ $) =$ Wt. Ave.

If any questions #4-6 are answered yes and/or if any of the Special Features b, d, or i have been selected, enter Exceptional for the functional index. If not, compute the contribution to vegetative diversity and integrity by each plant community by doing the following: multiply the ranking for each community (Question #3b) by its total proportion in Question 3a (percent of total). Then, the functional index for the entire wetland can be calculated four ways (as follows) and should be utilized according to the scope of the project:

- 3b) Individual Community Scores:** maintain raw data as recorded.
- 3c) Highest Quality Community:** report the highest-functioning community.
- 3d) Non-Weighted Average Quality of all Communities:** straight average
- 3e) Weighted Average Quality Based on Percentage of Each Community:** multiply each community rating by its percentage, then add all together.

Vegetative Diversity/ Integrity					
	3a. Proportion of Wetland	3b. Individual Community Scores	3c. Highest Rated Community Quality	3d. Non- Weighted Average	3e. Weighted Average
Community #1	T	A	If Spec. Features b, d or i are checked then rate Exceptional (2); if either question 4, 5, or 6 are Yes, then rate Exceptional (2); else:		
Community #2	U	B			
Community #3	V	C			
Community #4	W	D			
Community #5	X	E			
Community #6	Y	F			
Community #7	Z	G			
Overall Wetland Value Rating	1.0		: Highest Value of A-G	: (A+B+C+ D+E+F+G)/7 = Ave.	:(A*T)+(B* U)+(C*V)+ (D*W)+(E* X)+(F*Y)+(G*Z) = Wt. Ave.

6.2 MAINTENANCE OF CHARACTERISTIC HYDROLOGIC REGIME

A wetland's hydrologic regime or hydroperiod is the seasonal pattern of the wetland water level that is like a hydrologic signature of each wetland type. It defines the rise and fall of a wetland's surface and subsurface water. The constancy of the seasonal patterns from year to year ensures a reasonable stability for the wetland²³. The ability of the wetland to maintain a hydrologic regime characteristic of the wetland type is evaluated based upon wetland soil and vegetation characteristics, land use within the wetland, land use within the upland watershed contributing to the wetland, and wetland outlet configuration. Maintenance of the hydrologic regime is important for maintaining a characteristic vegetative community, and is closely associated with other functions including flood attenuation, water quality and groundwater interaction.

Measures the degree of human alteration of the wetland hydrology, either by outlet control or by altering immediate watershed conditions. Each parameter is weighted equally.

MnRAM #	Excel #	Variable Description	Type of Interaction
13	E17	Outlet—natural hydrologic regime	Controlling
14	E18	Dominant upland land use	Compensatory
15	E19	Soil condition/wetland	Compensatory
20	F24	Stormwater runoff/pretreatment-Reverse	Compensatory

Hydrologic Regime Index = (13+14+15+20)/4

6.3 FLOOD AND STORMWATER STORAGE/ATTENUATION

A wetland's ability to provide flood storage and/or flood wave attenuation is dependent on many characteristics of the wetland and contributing watershed. Characteristics of the subwatershed that affect the wetlands ability to provide flood storage and attenuation include: soil types, land use and resulting stormwater runoff volume, sediment delivery from the subwatershed, and the abundance of wetlands and waterbodies in the subwatershed. Wetland characteristics which affect the wetland's ability to store and or attenuate stormwater include: condition of wetland soils; presence, extent, and type of wetland vegetation; presence and connectivity of channels; and most importantly outlet configuration. Higher rated wetlands will have an unaltered or restricted outlet, undisturbed wetland soils, dense emergent vegetation without channels, a high proportion of impervious surfaces in the subwatershed, large runoff volumes, clayey upland soils, and few wetlands present within the subwatershed.

This formula is based on the Surface Water Storage Functional Capacity Index scoring concept and equation²⁴. The formula was altered with the addition of three surface flow characteristics and two stormwater runoff parameters (Stormwater Runoff Quality/Quantity and Subwatershed Wetland Density) along with the removal of two parameters (Soil Porosity and Subsurface Outlet,

²³ Mitsch and Gosselink, 2000

²⁴ Lee et al., 1997

which is already characterized in another parameter). This index is comprised of 5 primary processes, which are weighted equally; included in each major process are one to three characteristics that equally contribute to that process.

1. **Outlet Characteristics:** Outlet characteristics
2. **Upland Watershed:** Upland land use, Upland soils,
3. **Wetland Condition/Land Use:** Wetland land use, sediment delivery
4. **Runoff Characteristics:** Stormwater runoff quality/quantity, subwatershed wetland density
5. **Surface Flow Characteristics:** Flow-through emergent vegetation density, surface flow characteristics

Flood and Stormwater Storage Index Computation:

Entire Formula: Outlet for flood retention{12} + (Dominant upland use-RR{14}+ Upland soils{19})/2 + (Soil condition{15} + Sediment delivery{18})/2 + Stormwater runoff pretreat&det{20} + Subwatershed wetland density{21})/2 + (Percent emergent vegetative cover{16} + Flow-through emergent vegetative roughness{17} + Channels/sheet flow{22})/3)/5.

1. If 12=0, then: $((14+19)/2+(15+18)/2+(20+21)/2+(16+17+22)/3)/4$
2. If 12>0, then: $(12+(14+19)/2+(15+18)/2+(20+21)/2+(16+17+22)/3)/5$

No changes to the formula are necessary if 16=0.

Flood and Stormwater Storage/Attenuation Variables

MnRAM #	Excel #	Variable Description	Type of Interaction
12	E16	Outlet—flood attenuation	Controlling—optional
14	F18	Dominant upland land use-RR	Compensatory
19	E23	Upland soils	Compensatory
15	E19	Soil condition	Compensatory
18	E22	Sediment delivery	Compensatory
20	E24	Stormwater pretreatment &detention	Compensatory
21	E25	Subwatershed wetland density	Compensatory
16	F20	Emergent vegetation % cover	Comp.—optional
17	E21	Emergent vegetation flood resistance	Comp.—optional
22	E26	Channels/sheet flow	Compensatory

6.4 DOWNSTREAM WATER QUALITY PROTECTION

This rates the wetland's ability and opportunity to protect valuable downstream resources. Valuable downstream resources include recreational waters (i.e. lakes, streams, rivers, creeks, etc) and potable water supplies. The level of functioning is determined based on runoff characteristics, sedimentation processes, nutrient cycling, and the presence and location of significant downstream water resources. Runoff characteristics that are evaluated include: land use and soils in the upstream watershed, the stormwater delivery system to the wetland, and sediment delivery characteristics. The ability of the wetland to remove sediment from stormwater is determined by emergent vegetation and overland flow characteristics. A high nutrient removal rating indicates dense vegetation and sheet flow to maximize nutrient uptake and residence time within the wetland. The opportunity for a wetland to protect a valuable water resource diminishes with distance from the wetland so wetlands with valuable waters within 0.5 miles downstream have the greatest opportunity to provide protection.

Compute Functional Index for Downstream Water Quality Protection

This functional index computation was derived from a combination of Nutrient Cycling and Retention of Particulates functions in the HGM Prairie Pothole draft guidebook⁵⁴ with the downstream sensitivity concept from *The Minnesota Wetland Evaluation Methodology*. Three major processes make up equal portions of the Downstream Water Quality Protection function²⁵ with a measure of opportunity to protect downstream resources; each process is comprised of two to four observable parameters.

1. **Rate, Quantity, and Quality of Runoff to the Wetland:** this is characterized by the conditions in the upstream watershed; both land use and soils, that affect the sediment and nutrient loads to the wetland, and by the existing storm water delivery system to the wetland (Upland watershed conditions, storm water runoff, evidence of sediment delivery, and upland buffer each comprise 1/16 of the entire downstream water quality functional index based on their contribution to sediment removal).
2. **Sedimentation:** this is characterized by the presence of flow-through emergent vegetation density and by the overland flow characteristics within the wetland. A wetland with primarily sheet flow through the wetland and dense emergent vegetation density will allow sediment to drop out more effectively than a wetland with channel flow and no vegetation (When all parameters are applicable; emergent vegetative density and overland flow characteristics each make up 1/8 of the total downstream water quality functional index based on their contribution to sediment removal).
3. **Nutrient Uptake:** this is characterized by the outlet configuration and vegetative characteristics. A wetland with long water retention times has more capacity to remove nutrients from the water column via physical and biological processes. Vegetation slows floodwaters by creating frictional drag in proportion to stem density which allows sediment particles to settle out, thereby improving the water quality for downstream uses (Outlet characteristics and vegetative density each make up 1/8 of the total downstream water quality functional index based on their contribution to nutrient uptake).

²⁵ Derived from a combination of Nutrient Cycling and Retention of Particulates functions in the HGM Prairie Pothole draft guidebook (Lee et al., 1997) with the downstream sensitivity concept from *The Minnesota Wetland Evaluation Methodology*.

4. **Downstream Sensitivity:** if the wetland contributes to the maintenance of water quality within one-half mile of a recreational water body or potable water supply source downstream, it operates at a higher functioning level than a similar wetland farther from or without significant downstream water resources (This factor accounts for ¼ of the total downstream water quality functional index).

Downstream Water Quality Functional Index Computations:

1. If 12=0, then: $(14+20_{\text{reversed}} + 18 + (23+24+26)/3 + (16+17)/2 + 27)/6$
 2. If 12>0, then: $(14+20_{\text{reversed}} + 18 + (23+24+26)/3 + (16+17)/2 + 27 + 12)/7$

No changes to the formula are necessary if 16=0.

Entire Formula:

(Dominant upland land use{ 14 } + Stormwater runoff pretreatment & detention{ 20_{reversed} } + Sediment delivery { 18 } + (Upland buffer width{ 23 }WQ + Upland buffer vegetative cover{ 24 } + Upland buffer slope { 26 })/3 + (Flow-through %emergent vegetative cover{ 16 } + Flow-through emergent vegetative roughness{ 17 })/2 + Downstream sensitivity{ 27 }+ Outlet for flood{ 12 })/7

Downstream Water Quality Variables

MnRAM #	Excel #	Variable Description	Type of Interaction
14	E18	Dominant upland land use	Controlling
20	E24	Stormwater runoff pretreatment &detention	Controlling
18	E22	Sediment delivery	Controlling
23	G27	Upland buffer width	Comp.
24	G28	Upland area management	Comp.
26	G34	Upland area slope	Comp.
16	F20	Emergent vegetation (% cover)	Comp.—optional
17	E21	Emergent vegetation (roughness coefficient)	Comp.—optional
27	E39	Downstream sensitivity	Comp.
12	E16	Outlet for flood	Controlling--optional

6.5 MAINTENANCE OF WETLAND WATER QUALITY

The sustainability of a wetland is partially driven by the quality and quantity of stormwater runoff entering the wetland. The ability of the wetland to sustain its characteristics is evaluated based on characteristics of the contributing subwatershed and indicators within the wetland. Subwatershed conditions which affect the wetland’s sustainability in relation to water quality impacts include: upland land use; sediment delivery characteristics to the wetland; stormwater runoff volumes and rates; and the extent, condition, and width of upland buffer. Indicators of nutrient loading to the wetland indicate that a diverse wetland may not be sustainable. Indicators that a wetland has been affected by nutrient loading include the presence of monotypic vegetation and/or algal blooms.

This functional index was derived from a combination of sources including MNRAM, HGM, WEM, WET, and experiences of the project team. The sustainability of a wetland

is partially driven by the quality and quantity of stormwater runoff entering the wetland. The ability of the wetland to sustain its characteristics is evaluated based on characteristics of the contributing subwatershed and indicators within the wetland. Subwatershed conditions which affect the wetland's sustainability in relation to water quality impacts include: upland land use; sediment delivery characteristics to the wetland; stormwater runoff volumes and rates; and the extent, condition, and width of upland buffer. Indicators of nutrient loading to the wetland indicate that a diverse wetland may not be sustainable. Indicators that a wetland has been affected by nutrient loading include the presence of monotypic vegetation and/or algal blooms.

Wetland Water Quality Functional Index Computation:

$$(3e*2+14+20_{\text{reversed}} +(23+24+26)/3+18+28)/7$$

Entire Formula:

$$(Vegetative\ Diversity/Integrity\{3e*2\} + Dominant\ upland\ land\ use\{14\} + Stormwater\ runoff\ pretreatment\ \&\ detention\{20_{\text{reversed}}\} + (Upland\ buffer\ width\{23\}WQ + Upland\ buffer\ vegetative\ cover\{24\} + Upland\ buffer\ slope\{26\})/3 + Sediment\ delivery\{18\})/2 + Nutrient\ loading\{28\})/7$$

Wetland Water Quality Variables

MnRAM #	Excel #	Variable Description	Type of Interaction
3e	D6*2	Vegetative Diversity/Integrity	Contributing
14	E18	Dominant upland land use	Contributing
20	F24	Stormwater runoff pretreatment and detention—RR	Contributing
23	G27	Upland buffer width	Contributing
24	G28	Upland area management	Contributing
26	G34	Upland area slope	Contributing
18	E22	Sediment delivery	Contributing
28	E40	Nutrient loading	Contributing

This functional index was derived from a combination of sources including MNRAM, HGM, WEM, WET, and experiences of the project team. The sustainability of a wetland is partially driven by the quality and quantity of stormwater runoff entering the wetland. The ability of the wetland to sustain its characteristics is evaluated based on characteristics of the contributing subwatershed and indicators within the wetland. Subwatershed conditions which affect the wetland's sustainability in relation to water quality impacts include: upland land use; sediment delivery characteristics to the wetland; stormwater runoff volumes and rates; and the extent, condition, and width of upland buffer. Indicators of nutrient loading to the wetland indicate that a diverse wetland may not be sustainable. Indicators that a wetland has been affected by nutrient loading include the presence of monotypic vegetation and/or algal blooms.

6.6 SHORELINE PROTECTION

Shoreline protection is evaluated only for those wetlands adjacent to lakes, streams, or deepwater habitats. The function is rated based on the wetlands opportunity to protect the shoreline; i.e. wetlands located in areas frequently experiencing large waves and high

currents have the best opportunity to protect the shore. In addition, shore areas composed of sands and loams with little vegetation or shallow-rooted vegetation will benefit the most from shoreline wetlands. The wetland width, vegetative cover, and resistance of the vegetation to erosive forces determine the wetland's ability to protect the shoreline.

Each of the five parameters contributes equally²⁶: based primarily on the characteristics presented in WEM with a simple, straightforward computation of the index assuming all characteristics contribute equally.

MnRAM #	Excel #	Variable Description	Type of Interaction
29	E41	Shoreline?	Controlling
30	E42	Rooted shoreline vegetation (% cover)	Contributing
31	E43	Wetland width (average)	Contributing
32	E44	Emergent vegetation erosion resistance	Contributing
33	E45	Shoreline erosion potential	Contributing
34	E46	Bank protection ability	Contributing

Shoreline Protection Functional Index Computation:

If 29=1, then:

$$\text{Shoreline Protection Index} = (30+31+32+33+34)/5$$

Entire Formula:

(Rooted shoreline vegetation {30} + Average shoreline wetland width {31} + Emergent vegetation erosion resistance {32} + (Shoreline erosion potential {33} + Bank protection ability {34})/5

6.7 MAINTENANCE OF CHARACTERISTIC WILDLIFE HABITAT STRUCTURE

The ability of a wetland to support various wildlife species is difficult to determine due to the specific requirements of the many wildlife species that utilize wetlands. This function determines the value of a wetland for wildlife in a more general sense, and not based on any specific species. The characteristics evaluated to determine the wildlife habitat function include: vegetative quality, outlet characteristics (which control hydrologic regime), upland land use, wetland soil type and conditions, water quality of storm water runoff entering the wetland, upland buffer extent, condition, and diversity; the interspersions of wetlands in the area; barriers to wildlife movement; wetland size; vegetative and community interspersions within the wetland; and amphibian breeding potential and overwintering habitat.

Thirteen parameters are weighed equally as described below; vegetative quality weighted double the other factors. The questions are borrowed or modified from MNRAM, WET, WEM, and HGM methodologies, combined to provide a measure of wildlife habitat in general, not focusing on any particular species.

If Rare Wildlife (35) or Rare Natural Community (36) are true, then this Index is Exceptional.

²⁶ Based primarily on the characteristics presented in WEM.

If Special Features d, g, or j are checked, then this Index is Exceptional, otherwise, follow conditions below:

If 37=0 and 38=0 and 39=0, then:

$$(3e^2+40+41+(23+24+25)/3+13+20)/7$$

If 38=0 and 39=0, then:

$$(3e^2+37+40+41+(23+24+25)/3+13+20)/8$$

If 37=0 and 39=0, then:

$$(3e^2+38+40+41+(23+24+25)/3+13+20)/8$$

If 37=0 and 38=0, then:

$$(3e^2+39+40+41+(23+24+25)/3+13+20)/8$$

If 39=0, then:

$$(3e^2+37+38+40+41+(23+24+25)/3+13+20)/9$$

If 38=0, then:

$$(3e^2+39+37+40+41+(23+24+25)/3+13+20)/9$$

If 37=0, then:

$$(3e^2+39+38+40+41+(23+24+25)/3+13+20)/9$$

If 37>0 and 38>0 and 39>0, then:

$$(3e^2+39+37+38+40+41+(23+24+25)/3+13+20)/10$$

Entire Equation:

(Vegetative Diversity/Integrity {3e*2} + Wetland Detritus {39} + Vegetation Interspersion {37} + Community Interspersion {38} + Wetland Interspersion {40} + Wildlife Barriers {41} + (Upland buffer width {23}WQ + Upland Area Management{24} + Upland area diversity {25})/3 + Outlet natural hydrologic regime {13}+ Stormwater runoff pretreatment and detention 20)/11

MnRAM #	Excel #	Variable Description	Type of Interaction
41	E53	Wildlife barriers	Controlling
3e	D6	Vegetative Ranking (communities' weighted average)	Compensatory
39	E51	Wetland detritus (n/a)	
23	I27	Upland buffer average width	
24	G28	Upland area management	
25	G31	Upland area diversity	
13	E17	Outlet natural hydrologic regime	
20	F24	Stormwater runoff pretreatment & detention—RR	
37	F49	Vegetation interspersion (n/a)	
38	F50	Community interspersion (n/a)	
40	E52	Wetland interspersion	

6.8 MAINTENANCE OF CHARACTERISTIC FISH HABITAT

The ability of the wetland to support native fish populations is determined by structural factors within the wetland as well as water quality contributions from upland factors. Wetlands rated High are lacustrine or riverine and provide spawning/nursery habitat, or

refuge for native species (included but not limited to game fish). Wetlands rated Low for fish habitat do not have a direct hydrologic connection to a waterbody with a native fishery or have poor water quality.

MnRAM #	Excel #	Variable Description	Type of Interaction
46	E58*2	Fish habitat quality	Controlling
29	D41	Fringe wetland?	Contributing
24	G28	Adjacent area management	Compensatory
18	E22	Sediment delivery	Compensatory
20 (R)	F24	Storm water runoff	Compensatory
28	E40	Nutrient load	Compensatory
30	E42	Percent cover	Compensatory
31	E43	Wetland shoreline width	Compensatory
33 (R)	F45	Shoreline erosion potential	Compensatory

Fish Habitat Functional Index Computation:

If Special Features a or g are checked, then Fishery Habitat Index = Exceptional.

If 46=0, then Fishery Habitat = N/A

If 29=0, Fishery Habitat Index = $[(46*2)+24+18+20(R) +28]/6$

If 29>0, Fishery Habitat Index = $[(46*2)+24+18+20(R) +28+30+31+33(R)]/9$

6.9 MAINTENANCE OF CHARACT. AMPHIBIAN HABITAT FOR BREEDING/OVERWINTERING

The ability of a wetland to support various amphibian species is difficult to determine due to the specific requirements of the many amphibian species that depend on wetlands. This function determines the value of a wetland for amphibians in general, not based on specific species. An adequate wetland hydroperiod and the presence or absence of predatory fish are considered to be limiting variables for this function. In general, wetlands must remain inundated until early to mid-June to allow the larval stages to metamorphose into adults. Because many amphibians are partly terrestrial, the characteristics evaluated to determine the amphibian habitat function include numerous hydrology and terrestrial measures. The characteristics evaluated include: upland land use, upland buffer width, water quality of storm water runoff entering the wetland, barriers to wildlife movement, and amphibian breeding potential and overwintering habitat.

An adequate wetland hydroperiod (Question 42) is considered to be the primary limiting variable for this functional index. If the hydroperiod is insufficient for breeding, the wetland rating for amphibian use will be Not Sufficient. The status of predatory fish in the wetland (Q.43) is a secondary limiting factor to the final rating; the lowest rating for this variable, however, is 0.1 (Low), rather than zero (Not Sufficient).

Amphibians' ability to use a particular wetland for over wintering is a contributing factor in rating the wetland's functional index (Q.44). Because most amphibians are partly terrestrial, the extent of upland buffer habitat surrounding the wetland (Q.23) is an

important habitat component²⁷ and is weighted by a factor of two. Question 14 (Upland Land Use) is also included as an indicator of the quality of the surrounding upland habitat⁵⁶. Unnatural fluctuations in water depth in wetlands from conducted storm water runoff can impair reproductive success in amphibians, which often attach their eggs to stems of wetland vegetation, e.g., salamanders, tree frogs, green frogs, and wood frogs²⁸. Extreme water level fluctuations during winter may also cause mortality in overwintering reptiles and amphibians²⁹. Thus, Question 20 is included in the formula, with a reverse rating. Question 41 (Barriers) is included because access to and from the wetland by amphibians is an important factor in habitat quality³⁰.

Amphibian Habitat Functional Index Computation:

If 42=0, then N/A

Otherwise: Amphibian Habitat Index = (43) * [(44 + 2*23_{wildlife} + 14 + 41 + 20_{reversed})/6]

Entire Formula:

If Amphibian Breeding Potential-Hydroperiod {42} is applicable, then: (Amphibian Breeding Potential-Predator Fish {43}) * {(Amphibian Overwintering Habitat {44}+ 2*Upland Buffer Width (23)_{wildlife} + Dominant Upland Land Use {14} + Barriers {41} + Stormwater Input {20_{reverse}}}/6}

Amphibian Habitat Variables

MnRAM #	Excel #	Variable Description	Type of Interaction
42	D54	Amphibian breeding potential—hydroperiod	Controlling
43	D55	Amphibian breeding potential—fish presence	Controlling
44	E56	Amphibian overwintering habitat	Compensatory
23	I27	Upland buffer width	Compensatory
41	E53	Wildlife barriers	Compensatory
14	E18	Dominant upland land use	Compensatory
20	F24	Stormwater runoff pretreatment & detention—RR	Compensatory

6.10 AESTHETICS/RECREATION/EDUCATION/CULTURAL/SCIENCE

The aesthetics/recreation/education/cultural and science function and value of each wetland is evaluated based on the wetland’s visibility, accessibility, evidence of recreational uses, evidence of human influences (e.g. noise and air pollution) and any known educational or cultural purposes. Accessibility of the wetland is key to its aesthetic or educational appreciation. While dependent on accessibility, a wetland's functional level could be evaluated by the view it provides observers. Distinct contrast

²⁷ Knutson et al., 2000

²⁸ Richter and Azous, 1995

²⁹ Hall and Cuthbert, 2000

³⁰ Knutson, et al., 1999; Findlay and Bourdages, 2000; Semlitsch, 2000.

between the wetland and surrounding upland may increase its perceived importance. Also, diversity of wetland types or vegetation communities may increase its functional level as compared to monotypic open water or vegetation. Excess negative human influence on the wetland is counted double in the formula.

All questions contribute equally to the overall index.

MnRAM #	Excel #	Variable Description	Type of Interaction
48	E60	Rare educational opportunity	Controlling
49	E61	Wetland visibility	Compensatory
50	E62	Proximity to population	Compensatory
51	E63	Public ownership	Compensatory
52	E64	Public access	Compensatory
53	E65	Human influence—wetland	Compensatory
54	E66	Human influence—viewshed	Compensatory
55	E67	Spatial buffer	Compensatory
56	E68	Recreational activities in wetland	Compensatory

Aesthetics/Recreation/Education/Cultural/Science Functional Index Computations:

If Special Features c, h, or u is checked³¹, or

If 48=1, then Index = Exceptional;

If 53=0.1 (Low), then = (50+51+52+2*53+54+55+56)/8

If 53>0.1, then = (49+50+51+52+53+54+55+56)/8

Entire Formula

(Wetland Visibility {49} + Proximity to Population {50} + Public Ownership {51} + Public Access {52} + Human Influence - Wetland {53} + Human Influence - Viewshed {54} + Spatial Buffer {55} + Recreational Activities in Wetland {56})/8

6.11 COMMERCIAL USES

This question considers the nature of any commercially-valuable use of the wetland and requires the assessor to consider how such use may be a detriment to the sustainability of the wetland. Some row crops can be planted in Type 1 wetlands after spring flooding has ceased and still have adequate time to grow to maturity. This non-wetland-dependent agricultural use of wetlands may include hay, pasture/grazing, or row crops such as soybeans or corn. Wetland-dependent crops include wild rice and cranberries, which rely on the wetland hydrology for part of their life cycle.

³¹ c = Designated scientific and natural area; h = Archeologic or historic site designated by the State Historic Preservation Office; u = State or Federal designated wilderness area.

Sustainable uses of the wetland would not require modifying a natural wetland. Products in this category would include collection of botanical products, wet native grass seed, floral decorations, wild rice, black spruce, white cedar, and tamarack. Sustainable uses may require modification of the natural hydrology, such as for wetland-dependent crops (rice, cranberries). Haying and grazing can be less intrusive agricultural activities utilized more or less casually when hydrologic conditions permit; light pasture and occasional haying would be considered more or less sustainable. Like peat-mining, cropping is an unsustainable use of the wetland as it results in severe alterations of wetland characteristics (soil, vegetation, hydrology).

MnRAM #	Excel #	Variable Description	Type of Interaction
57	E69	Commercial crop—hydrologic impact	Controlling

Commercial Uses Functional Index = 57

6.12 GROUND-WATER INTERACTION

The ground water interaction function is the most difficult to assess. Here the most likely type of ground water interaction is determined, i.e. recharge or discharge, or a combination. In many cases, a wetland will exhibit both recharge and discharge characteristics, however one is usually more dominant. Several wetland and watershed characteristics are evaluated to determine the likely interaction including: wetland soil type, upland land use, upland soil types and wetland size, wetland hydroperiod, wetland outlet characteristics, and topographic relief.

The purpose of this function is strictly to determine the likelihood of the appropriate ground-water interaction based on observable characteristics of the wetland and watershed. The significance of ground water as a component of the wetland water budget is the most difficult functional characteristic to determine without large quantities of detailed hydrologic and geologic information. The following methodology takes the most easily observable and distinct measures of recharge/discharge relationships from the *Wetland Evaluation Technique*³² and the *Hydrogeomorphic Assessment Methodology*³³. In many wetlands, surface water and ground water both make significant contributions to the water budget, but occasionally recharge or discharge is dominant. The goal here is to identify the dominant ground-water interaction (if there is one) to help guide future management and provide an indication when additional information may be warranted.

³² Adamus, et al., 1987

³³ Magee and Hollands, 1998

- If 5 or 6 of questions 58-63 are answered the same, this indicates a strong likelihood that the most frequently stated interaction exerts the primary influence on the wetland.
- If 3-4 questions are answered the same, then the wetland is likely influenced by a combination of both recharge and discharge interactions (i.e. both types of ground water interaction are likely to be present at some point during most years).

- 58. Wetland Soils – from HGM system functional assessments and Novitzki
- 59. Subwatershed Land Use/Imperviousness – taken from WET Volume I
- 60. Wetland Size and Upland Soils – taken from WET Volume I and HGM
- 61. Wetland Hydrologic Regime– taken from WET Volume I and HGM
- 62. Inlet/Outlet Configuration – taken from WET Volume I and HGM
- 63. Upland Topographic Relief – taken from WET Volume I

Special Concerns for Recharge Wetlands

Wherever ground water recharge is indicated as the **primary** interaction and the wetland lies within a sensitive ground water area (**Special Feature Question q**), a contribution area to a public water supply, or a wellhead protection area (**Special Feature Question r**), it should be recorded as Exceptional for the ground water/wetland function.

6.13 WETLAND RESTORATION POTENTIAL

The potential for wetland restoration is determined based on the ease with which the wetland could be restored, the number of landowners within the historic wetland basin, the size of the potential restoration area, the potential for establishing buffer areas or water quality ponding, and the extent and type of hydrologic alteration. Each variable uses the High, Medium, Low rating rather than raw numbers—see MnRAM for individual ranges.

MnRAM #	Excel #	Variable Description	Type of Interaction
64	D79	Wetland Restoration Potential	Controlling
65	F80	Number of Landowners Affected	Contributing
21	E25	Subwatershed Wetland Density	Contributing
66b	F82	Total Wetland Restored Size (Potential)	Contributing
66c	F83	Calculated potential new wetland area	Contributing
67	F84	Potential Buffer Width	Contributing
68	F85	Likelihood of Restoration Success	Contributing

If 64="Yes", then Wetland Restoration Potential = (65+21+66b+66c+67+68)/6,

Otherwise, if 64="No" then "N/A"

Entire Formula

(Landowners Affected by Restoration (65)+Subwatershed Wetland Density (21)+Wetland Restoration Size (66b)+Proportion of Wetland Drained (66c)+Potential Buffer Width (67)+Likelihood of Restoration Success (68))/6

6.14 WETLAND SENSITIVITY TO STORMWATER INPUT AND URBAN DEVELOPMENT

The sensitivity of the wetland to stormwater and urban development is determined based on guidance within the *Storm-Water and Wetlands: Planning and Evaluation Guidelines for Addressing Potential Impacts of Urban Storm-Water and Snow-Melt Runoff on Wetlands*, State of Minnesota Storm-Water Advisory Group, June, 1997.

Use habitat proportions from Vegetative Integrity section and enter into a formula to compute answer according to the following criteria³⁴.

- Exceptional = Sedge meadows, open and coniferous bogs, calcareous fens, low prairies, wet to wet-mesic prairies, coniferous swamps, lowland hardwood swamps, or seasonally flooded basins.
- A = Shrub-carrs, alder thickets, diverse fresh wet meadows dominated by native species, diverse shallow and deep marshes, and diverse shallow, open water communities.
- B = Floodplain forests, fresh wet meadows dominated by reed canary grass, shallow and deep marshes dominated by cattail, reed canary grass, giant reed or purple loosestrife, and shallow, open water communities with low to moderate vegetative diversity.
- C = Gravel pits, cultivated hydric soils, or dredge/fill disposal sites.

6.15 ADDITIONAL STORMWATER TREATMENT NEEDS

This rates the sustainability of the wetland with regard to stormwater discharges to the wetland. The need for additional stormwater treatment prior to discharge to the wetland is rated based on the overall rating for Maintenance of Wetland Water Quality. If a wetland is severely degraded by stormwater inputs, the rating will be low, since a diverse, high quality wetland will not be sustainable.

Use functional rating for Maintenance of Wetland Water Quality (MWWQ) as follows (this index is rated strictly from the measure of the water quality in the wetland and the sustainability, i.e. if the water quality in the wetland is low, additional stormwater treatment is needed to protect the wetland and the rating is low):

Use Value for Maintenance of Wetland Water Quality Index (D76, Excel spreadsheet) and apply to criteria below.

- A = Maintenance of Wetland Water Quality Index >0.66 (no additional treatment needed)
- B = $0.33 < \text{Maintenance of Wetland Water Quality Index} \leq 0.66$ (sediment removal needed)

³⁴ Taken directly from State of Minnesota Storm-Water Advisory Group, 1997.

C = Maintenance of Wetland Water Quality Index < 0.33 (sediment and nutrient removal needed)

- Knutson, M.G., J.R. Sauer, D.A. Olsen, M.J. Mossman, L.M. Hemesath and M.J. Lannoo. 1999. *Effects of landscape composition and wetland fragmentation on frog and toad abundance and species richness in Iowa and Wisconsin, U.S.A.* Conservation Biology 13:1437-1446.
- Knutson, M.G., J.R. Sauer, D.A. Olsen, M.J. Mossman, L.M. Hemesath and M.J. Lannoo. 2000. *Landscape associations of frog and toad species in Iowa and Wisconsin, U.S.A.* Jour. Iowa Acad. Sci. 107(3):134-145.
- Knutson, P. L., J. C. Ford, and M. R. Inskeep. 1981. *National Survey of Planted Salt Marshes (vegetative stabilization and wave stress).* Wetlands 3:129-153.
- Kuchler, A.W. 1967. *Vegetation Mapping.* The Ronald Press, New York, New York.
- Lannoo, M.J. 1998. *Amphibian conservation and wetland management in the upper midwest: a catch 22 for the cricket frog.* In Status and Conservation of Midwestern amphibians. M.J. Lannoo, ed. University of Iowa Press, Iowa City, Iowa. Pages 331-339.
- Lee, L.C. and Mark M. Brinson, William J. Kleindl, P. Michael Whited, Michael Gilbert, Wade L. Nutter, Dennis F. Whigham, Dave DeWald. 1997. *Revised Operational Draft Guidebook for the Hydrogeomorphic Assessment of Temporary and Seasonal Prairie Pothole Wetlands.* Seattle, WA. pp. 116+app.
- Lehtinen, R.M., S.M. Galatowitsch and J.R. Tester. 1999. *Consequences of habitat loss and fragmentation for wetland amphibian assemblages.* Wetlands 19:1-12.
- Magee, D.W. and Garrett G. Hollands. 1998. *A Rapid Procedure for Assessing Wetland Functional Capacity.* Association of State Wetland Managers. 190 pp.
- Michigan DNR. 1981. *Manual for Wetland Evaluation Techniques.* Michigan Department of Natural Resources, Lansing, MI.
- Mitch & Gosselink 2000. *Wetlands 3rd Edition.*
- Ogawa, H. and J. W. Male. 1983. *The Flood Mitigation Potential of Inland Wetlands.* Water Resources Research Center. University of Massachusetts.
- Oldfield, B. and J.J. Moriarty. 1994. *Amphibians and reptiles native to Minnesota.* University of Minnesota Press, Minneapolis, Minnesota.
- Reed, P.B. 1988 (and 1993 supplement). *National List of Plant Species that Occur in Wetlands, U. S. Fish and Wildlife Service National Wetland Inventory, St. Petersburg, FL.*
- Rheinhardt, R., Mark M. Brinson, N. Eric Fleming, J. Glenn Sandifer, Jr., 1997. *Deciduous Wetland Flats Interim HGM Model.* 50 pp.
- Richter, Klaus O. and Amanda L. Azous. 1995. *Amphibian occurrence and wetland characteristics in the Puget Sound Basin.* Wetlands 15:305-312.
- Seaberg, J.K. and Douglas D. Hansen. 2000. *Metropolitan Area Groundwater Model Project Summary: Northwest Province, Layers 1, 2, and 3 Model, Version 1.00.* Minnesota Pollution Control Agency.
- Semlitsch, R.D. 2000. *Principles for management of aquatic-breeding amphibians.* J. Wildlife Mgmt. 64:615-631.
- Shaw, S. and C.G. Fredine. 1956. *Wetlands of the United States Circular 39.* U.S. Fish and Wildlife

Service. U.S. Government Printing Office, Washington, D.C.

Smith, S.G. 1986. *The cattails (Typha): Interspecific ecological differences and problems of identification*. Proceedings of the 5th Annual Conference of the NALMS. Geneva, Illinois, 357-362.

Snodgrass, J.W., M.J. Komoroski, A.L. Bryan, Jr., and J. Burger, Jr. 2000. *Relationships among isolated wetland size, hydroperiod, and amphibian species richness: implications for wetland regulations*. Conservation Biology 14:414-419.

State of Minnesota Storm-Water Advisory Group. 1997. *Storm-Water and Wetlands: Planning and Evaluation Guidelines for Addressing Potential Impacts of Urban Storm-Water and Snow-Melt Runoff on Wetlands*.

Thill, D., and M. Jacobson. 2000. *The Minnehaha Creek Routine Assessment Method*, in conjunction with the Minnehaha Creek Watershed District.

Environmental Laboratory. 1987. Corps of Engineers Wetlands Delineation Manual. Technical Report Y-87-1, Final Report.

Verry, E.S. 1988. *The Hydrology of Wetlands and Man's Influence on it*. International Symposium on the Hydrology of Wetlands in Temperate and Cold Regions. Joensuu, Finland.

Wells, J. Mike Mueller, John Parker, Bruce Gerbig, S. Jatnieks-Straumanis, Bruce Wilson, John Kittelson, Paul Richert, Teri Sardinas, Scott LaChance, and Larry Smith, 1988. *The Minnesota Wetland Evaluation Methodology for the North Central United States*. Corps of Engineers in conjunction with the Minnesota Environmental Quality Board Wetland Evaluation Methodology Task Force, John R. Wells, Chairman.

Whited, D., Galatowitsch, S., Tester, J.R., Schik, K., Lehtinen, R., Husveth, J. 2000. Landscape and Urban Planning, 49 (1-2): 49-65.

Williams, R.E. 1968. *Flow of groundwater adjacent to a small, closed basin in glacial till*. Water Resources Research 4:777-783.

BMP References:

Board of Water and Soil Resources, Association of Metropolitan Soil and Water Conservation Districts. 1989. Minnesota Construction Site Erosion and Sediment Control Planning Handbook. St. Paul.

Minnesota Pollution Control Agency. 2000. Protecting Water Quality in Urban Areas: Best Management Practices for Dealing with Storm Water Runoff from Urban, Suburban and Developing Areas of Minnesota. Minneapolis, March, 2000.

Barr Engineering Company. 2001. Minnesota Urban Small Sites BMP Manual: Stormwater Best Management Practices for Cold Climates. Metropolitan Council Environmental Services, St. Paul, MN.

Appendix G

Vegetation Shoreline Buffer Brochure Examples


HENNINGSON

RESIDENTIAL NATIVE PLANTING



BEFORE

A native planting, also called natural landscaping, uses plants, trees and grasses that are suitable for growth in a particular geographical area



AFTER

PRACTICE:

- Native Planting

BENEFITS:

- Runoff volume reduction
- Slope stabilization
- Improved wildlife habitat
- Opportunity for public education and outreach
- Improved aesthetics

PARTNERS:

- Black Dog Watershed Management Organization

WATERSHED:

- Black Dog

INSTALLATION:

- Summer 2021

PROJECT: Installation of a 500 sq. ft. residential native garden.


COST: Project materials cost estimated at \$2,360

FUNDING: Landowners receive a \$250 Landscaping for Clean Water grant as well as technical assistance provided by the Dakota County Soil and Water Conservation District

Black Dog
Watershed Management Organization


LOCATION:

Kenosha Ave
Burnsville



DILLMAN

NATIVE SHORELINE PLANTING



A native shoreline planting is the use of native vegetation to protect a shoreline from existing or potential erosion.



PRACTICE:

- Native Shoreline Planting

BENEFITS:

- Shoreline stabilization and erosion reduction
- Improved water quality
- Improved wildlife habitat
- Opportunity for public education and outreach
- Improved aesthetics

PARTNERS:

- Black Dog Watershed Management Organization

WATERSHED:

- Black Dog

INSTALLATION:

- Summer 2021

PROJECT: Installation of a 620 square foot Native Shoreline Planting

COST: Project materials cost estimated at \$1,096

FUNDING: Landowners receive a \$250 Landscaping for Clean Water grant as well as technical assistance provided by the Dakota County Soil and Water Conservation District

Black Dog
Watershed Management Organization

LOCATION:

Keller Lake Road
Burnsville



Appendix H
Buckthorn Management Guidelines

Buckthorn Management Guidelines

Goal: Restore native plant communities in designated natural areas and other park locations by controlling and removing non-native invasive species.

Buckthorn belongs to the *Rhamnaceae* family. It is native to Europe and Asia, first appearing in the U.S. in the late 1700s. Buckthorn quickly naturalized in the woodlands of the northeastern states. Today buckthorn flourishes in the understory of Minnesota woodlands and in brushy thickets along roadsides and fields. It has become a major plant pest in natural woodlands and wetlands.

Buckthorn can grow to 15-20 feet and has dark green elliptical or oval leaves. In the fall its leaves hang on late into the season and without much color change. It starts easily from seed and will tolerate almost any soil condition or location. In partial shade it will outstretch its neighbors toward the light.

Buckthorn removal is recommended for those areas where the native plant community has been displaced by buckthorn species and where there is a high likelihood that the native plant community can be enhanced and restored.

Restoration of the native communities is the overall intent of non-native eradication efforts.

Volunteer Considerations

Volunteers must be trained in species identification, removal techniques and other aspects related to the eradication/restoration efforts.

Identification of buckthorn by volunteers is best performed during the month of October.

Process

Buckthorn removal is a long-term process requiring several steps over a three- to four-year period. Pulling seedlings, cutting and removing mature plants, chemically treating stumps and replanting the site with native species are critical to the long-term success of restoration efforts.

Staff are responsible for cutting mature plants and chemically treating the stumps in areas designated for restoration. A 20%-25% solution of glyphosate (Roundup) with a dye is used to paint, chemically treat, and mark the stumps.

Volunteer procedures

1. Hand pulling allowed by volunteers with training or under the supervision of a "trained" volunteer supervisor.
2. Use of loppers allowed by volunteers.
3. No use of power tools or chemicals by volunteers; chemicals and power tool use only by staff or contractor.
4. Volunteers must sign waiver form.

Recommended chronology of restoration activities with volunteers

Year one

- Seedlings cut or pulled (September-November)
- Mature trees cut by staff and/or volunteers in late fall (October-December)
- Stumps or stems chemically treated by staff immediately after cutting
- Removal of brush to a chipping location (or pile on site for burning)

Year two

- Remove seedlings by hand pulling or cutting and treating (June-November)
- Follow-up cutting by staff and/or volunteers in late fall (October-December) and chemically treat stump and stems.

Year three

- Seedling removal by hand pulling or cutting and treating as necessary
- Plant native understory shrubs, trees, ferns, wildflowers and grasses to approximate prior native plant community.

Year four

- Continued monitoring and buckthorn seedling removal

Other removal techniques

Mechanical

- Prescribed fire for seedlings; prescribed burns in early spring and fall annually or biannually to control buckthorn may have to be continued for several years

Chemical

- Cut-stump and stem treatment with glyphosate; 20%-25% active ingredient cut-stump; or basal bark spray treatment around the stem with 25-50% a.i. triclopyr (Garlon) – consideration of worker safety issues will dictate chemical selection. Glyphosate products registered for wetland/aquatic use should be used on water bodies and wetlands. Sponge applicators can help prevent chemical spill or spread to workers.
- Fosamine, a non-selective bud inhibitor for woody species, can be applied as a basal bark treatment in the fall at 3% a.i. concentration in winter

Another technique is goat rental.

The method of buckthorn control should be selected based on the site, safety concerns, and opportunities for continued vegetation management.

Other Sources for Guidance

University of Minnesota:

<http://www.extension.umn.edu/environment/agroforestry/woody-vegetation-control.html>

University of Wisconsin:

http://mipncontroldatabase.wisc.edu/search?name=common_buckthorn&habitat=7&season=7

Minnesota Department of Natural Resources:

<http://www.dnr.state.mn.us/invasives/terrestrialplants/woody/buckthorn/control.html>

U.S. Department of Agriculture Natural Resources Conservation Service:

<https://efotg.sc.egov.usda.gov/references/public/MN/797Buckthorn.pdf>

See Buckthorn Control Quick Guide for a summary of control techniques.

Appendix I
Pollinators Brochure

How can YOU help pollinators?

A decline in pollinators affects us all. Reversing this trend is important to our ecosystem as well as to human health and well-being. Pollinators have evolved with plants over thousands of years, developing unique and interdependent relationships. We can all do our part to help pollinators rebound from the challenges they face.

1. **Plant a variety of native flowering plants in your home garden, agricultural or natural landscapes (with bloom times from April to October).**
2. **Provide a variety of natural habitats for nesting sites and clean water sources.**
3. **Avoid pesticide use and purchase pollinator plants (and seeds) that have not been treated with systemic pesticides.**
4. **Help increase awareness about the need to protect pollinators**



PROTECTING Minnesota's Pollinators

There is increasing evidence that insect pollinators are disappearing at alarming rates. Major factors include loss of forage plants and nesting habitat, disease, pesticide use, and pests.



Pollination causes plants to produce the seeds and fruits that sustain wildlife and humans, and provides important ecosystem services. More than 1/3rd of all plants or plant products consumed by humans are dependent on pollinators.



Many Minnesota-grown crop plants cannot produce seed without the help of insect pollinators.

These include:

- Apples
- Berries
- Sunflowers
- Clovers
- Beans
- Squash
- Cucumbers



More resources about pollinators can be found at
<http://www.bwsr.state.mn.us/practices/pollinator/index.html>

Minnesota Board of Water & Soil Resources
www.bwsr.state.mn.us



Minnesota's Pollinators & Pollinator Plants

When these critters visit a flower to consume nectar and/or pollen, some of the pollen grains stick to their bodies. Pollination occurs when this pollen is transferred from one plant to another.

Bees

With over 4000 species, bees are considered the most important pollinators in North America, around 500 of which are native to Minnesota and Wisconsin. Bee families include honey bees, bumble bees, mason bees, carpenter bees, and sweat bees.




Butterflies & Moths

Butterflies and moths are also important pollinators and many are in trouble. Milkweed is the host plant for monarch butterfly caterpillars, and the loss of this plant is drastically reducing monarch butterfly populations. The Poweshiek skipperling, Dakota skipper, and Karner Blue butterflies are threatened or endangered in Minnesota.



Beetles, Flies, Wasps & Midges

Beetles are considered to be important pollinators because of their large numbers. Beetles play an important role in controlling agricultural pests. Though less effective as pollinators, many flies, wasps, midges, and even mosquitos visit flowers and consume nectar as part of their diet.



Hummingbirds

Of the 20 hummingbirds in North America, only the Ruby-throated is regularly found in Minnesota. This charismatic pollinator is attracted to brightly colored tubular flowers like the columbine.



Goldenrod
Solidago spp.



Joe Pye Weed
Eupatorium spp.



Milkweed
Asclepias spp.



Black-eyed Susan
Rudbeckia spp.



Prairie Blazing Star
Liatris spp.



Columbine
Aquilegia spp.